

Enumerative vs. Symplectic Invariants and Obstruction Bundles

Aleksey Zinger *

February 1, 2008

Contents

1	Introduction	2
1.1	Background and Results	2
1.2	Summary	3
1.3	Fundamental Notation	4
2	Spaces of Bubble Maps	6
2.1	Bubble Trees	6
2.2	The Basic Gluing Construction	7
2.3	Curves with Marked Points	10
2.4	Bubble Maps	12
2.5	Stratums of Bubble Maps	13
3	The Gluing Construction and the Obstruction Bundle	17
3.1	Summary and Notation	17
3.2	The Basic Setup	20
3.3	Construction of Nearly Holomorphic Bubble Maps	24
3.4	Scale of Variations	26
3.5	Obstruction Bundle Setup	30
3.6	The Gluing Map	34
3.7	An Implicit Function Theorem	36
3.8	The Orientation of $\mathcal{M}_{\Sigma, \nu, \lambda^*}(\mu)$ and the Gluing Map	38
3.9	Gluing Maps for Spaces $\bar{\mathcal{U}}_{\mathcal{T}}^{(0)}(\mu)$ and Orientations	46
4	Technical Issues	49
4.1	Continuity of the Gluing Map	49
4.2	Injectivity of the Gluing Map	53
4.3	The Basic Gluing Map and the Space of Balanced Maps	56
4.4	Gromov Convergence and the L^p -norm of the Differential	60
4.5	Surjectivity of the Gluing Map	64

*Partially supported by NSF Graduate Research Fellowship and NSF grant DMS-9803166

5	Appendix	67
5.1	Properties of Smooth Families of Metrics on Σ	67
5.2	Sobolev Inequalities for the Metrics g_v	71
5.3	Elliptic Estimates for the Metrics g_v	72
5.4	Fiber-Uniform Inverse for the Operator D_v	75

1 Introduction

1.1 Background and Results

Suppose (Σ, j) is a nonsingular Riemann surface of genus $g \geq 2$ and (V, J, ω) is a Kahler manifold of complex dimension n . If $\lambda \in H_2(V; \mathbb{Z})$, denote by $\mathcal{H}_{\Sigma, \lambda}(V)$ the set of simple (J, j) -holomorphic maps u from Σ to V such that $u_*[\Sigma] = \lambda$. Let $\mu = (\mu_1, \dots, \mu_N)$ be an N -tuple of proper oriented submanifolds of V such that

$$\text{codim } \mu \equiv \sum_{l=1}^{l=N} \text{codim } \mu_l = 2(\langle c_1(V, J), \lambda \rangle - n(g-1) + N). \quad (1.1)$$

For many Kahler manifolds (V, J, ω) and choice of constraints μ , the cardinality of the set

$$\mathcal{H}_{\Sigma, \lambda}(\mu) = \{(\Sigma; y_1, \dots, y_N; u) : u \in \mathcal{H}_{\Sigma, \lambda}(V); y_l \in \Sigma, u(y_l) \in \mu_l \forall l = 1, \dots, N\} \quad (1.2)$$

is finite and depends only on the homology classes of $\mu_1, \dots, \mu_N$. The cardinality $|\mathcal{H}_{\Sigma, \lambda}(\mu)|$ of the set $\mathcal{H}_{\Sigma, \lambda}(\mu)$ is then an *enumerative* invariant of the complex manifold (V, J) . Such numbers for algebraic manifolds (V, J) , e.g. the complex projective spaces $\mathbb{P}^n$, have been of great interest in algebraic geometry for a long time.

If (V, ω, J) is a semipositive symplectic manifold, the *symplectic* invariant of (V, ω) ,

$$\text{RT}_{g, \lambda}(\mu) \equiv \text{RT}_{g, \lambda}(\mu_1, \dots, \mu_N)$$

of [RT], is a well-defined integer. Due to the two composition laws of [RT], this symplectic invariant is often more readily computable than the enumerative invariant $|\mathcal{H}_{\Sigma, \lambda}(\mu)|$. In fact, all such symplectic invariants of $\mathbb{P}^n$ are easily computable. It is also shown in [RT] that the appropriately defined genus-zero invariants of $\mathbb{P}^n$ agree with the corresponding symplectic invariants. On the other hand, even for $\mathbb{P}^2$ and for genus one, the two invariants are no longer equal. In [I], the difference

$$\text{RT}_{1, \lambda}(\mu_1; \mu_2, \dots, \mu_N) - |\mathcal{H}_{\Sigma, \lambda}(\mu)|$$

is commuted for genus-one surfaces Σ and all projective spaces using an obstruction-bundle approach, first introduced by [T] in a very different setting. In [Z2], the difference

$$\text{RT}_{2, \lambda}(\mu) - |\mathcal{H}_{\Sigma, \lambda}(\mu)|$$

is commuted for genus-two surfaces Σ for $\mathbb{P}^2$ and $\mathbb{P}^3$ using a similar approach.

The main results of this paper are *detailed* descriptions of gluing pseudoholomorphic maps in symplectic geometry. Theorem 3.29 concerns the adoption of the obstruction-bundle idea of [T] into

the symplectic setting. Theorem 3.33 describes the local structure of moduli spaces of holomorphic rational maps under certain regularity conditions. In addition to providing the necessary analytic justification for the computations in [I] and [Z2], the two theorems lay foundations for comparing enumerative and symplectic invariants in many Kahler manifolds, in particular, the complex homogeneous manifolds. Furthermore, in [Z2], the second theorem is used to enumerate cuspidal rational curves in $\mathbb{P}^2$ and two-component tacnodal rational curves in $\mathbb{P}^3$. The same arguments can be used to count curves with higher-order degeneracies.

The author is grateful to T. Mrowka for pointing out the paper [I], encouraging the author to work out all of the analytic issues arising in [I], and sharing some of his expertise in applications of global analysis over countless hours of conversations. The author also thanks G. Tian for first introducing him to Gromov's symplectic invariants and helping him understand [LT].

1.2 Summary

Let $\Lambda^{0,1}\pi_\Sigma^*T^*\Sigma \otimes \pi_V^*TV \longrightarrow \Sigma \times V$ denote the bundle of (J, j) -antilinear homomorphisms from $\pi_\Sigma^*T^*\Sigma$ to π_V^*TV . If

$$\nu \in \Gamma(\Sigma \times V; \Lambda^{0,1}\pi_\Sigma^*T^*\Sigma \otimes \pi_V^*TV),$$

denote by $\mathcal{M}_{\Sigma, \nu, \lambda}$ the set of all smooth maps u from Σ to $\mathbb{P}^n$ such that $u_*[\Sigma] = \lambda$ and $\bar{\partial}u|_z = \nu|_{(z, u(z))}$ for all $z \in \Sigma$. If μ is an N -tuple of constraints as above, put

$$\mathcal{M}_{\Sigma, \nu, \lambda}(\mu) = \{(\Sigma; y_1, \dots, y_N; u) : u \in \mathcal{M}_{\Sigma, \nu, \lambda}; y_i \in \Sigma, u(y_i) \in \mu_i \ \forall i = 1, \dots, N\}.$$

If (V, ω, J) is semipositive, for generic choices of ν and μ , $\mathcal{M}_{\Sigma, \nu, \lambda}$ is a smooth finite-dimensional oriented manifold, and $\mathcal{M}_{\Sigma, \nu, \lambda}(\mu)$ is a zero-dimensional finite submanifold of $\mathcal{M}_{\Sigma, \nu, \lambda} \times \Sigma^N$, whose cardinality (with sign) depends only the homology classes of $\lambda_1, \dots, \lambda_N$; see [RT]. The symplectic invariant $\text{RT}_{g, \lambda}(\mu)$ is the signed cardinality of the set $\mathcal{M}_{\Sigma, \nu, \lambda}(\mu)$.

If $\|\nu_i\|_{C^0} \longrightarrow 0$ and $(\Sigma; \underline{y}_i; u_i) \in \mathcal{M}_{\Sigma, \nu_i, \lambda}(\mu)$, then a subsequence of $\{(\Sigma; \underline{y}_i, u_i)\}_{i=1}^\infty$ must converge in the Gromov topology to one of the following:

- (1) an element of $\mathcal{H}_{\Sigma, \lambda}(\mu)$;
- (2) $(\Sigma_T; \underline{y}; u)$, where Σ_T is a bubble tree of S^2 's attached to Σ with marked points $y_1, \dots, y_N$, and $u: \Sigma_T \longrightarrow V$ is a holomorphic map such that $u(y_i) \in \mu_i$ for $i = 1, \dots, N$, and
 - (2a) $u|_{\Sigma_T}$ is simple and the tree contains at least one S^2 ;
 - (2b) $u|_{\Sigma_T}$ is multi-covered;
 - (2c) $u|_{\Sigma_T}$ is constant and the tree contains at least one S^2 .

This convergence statement says that for all t sufficiently small all the elements of $\mathcal{M}_{\Sigma, t\nu, \lambda}(\mu)$ lie near one of the spaces described by (1)-(2c). In many practical applications, it is easy to show that there is a bijection between the elements of $\mathcal{H}_{\Sigma, \lambda}(\mu)$ and the nearby elements of $\mathcal{M}_{\Sigma, t\nu, \lambda}(\mu)$; see Proposition 3.30. This is the case for all projective spaces, provided λ is a sufficiently high multiple of the line (depending on the genus g). In [I] and [Z2], Cases (2a) and (2b) do not occur, but they may have to be considered when dealing with higher-dimensional projective spaces or higher genera. Thus, the convergence statement implies that if the signed cardinality of $\mathcal{M}_{\Sigma, t\nu, \lambda}(\mu)$ is $\text{RT}_{g, \lambda}(\mu)$ for all $t > 0$ sufficiently small, the number of elements of $\mathcal{M}_{\Sigma, t\nu, \lambda}(\mu)$ that lie near the spaces described by (2) is exactly $\text{RT}_{2, \lambda}(\mu) - |\mathcal{H}_{\Sigma, \lambda}(\mu)|$. The goal of this paper is to describe this difference in terms of the spaces of holomorphic maps themselves, which can be viewed as an *enumerative* object, rather than a *symplectic* one. We do need to assume that certain spaces of

holomorphic maps are smooth, but they do not need to have the expected dimension.

While there is a very good understanding of what constitutes a stable map, there is little in a way of commonly accepted notation for stable maps and various spaces of stable maps. In Section 2, we recall the definition of bubble or stable maps as well as set up analytically convenient notation. Our notation for bubble maps evolved from that of D. McDuff's lectures at Harvard. We also define various spaces of bubble maps and restate the definition of the Gromov topology on the set of all bubble maps in our notation.

In Section 3, we describe a gluing construction and an obstruction bundle setup. One difficulty in the gluing procedure is that a certain operator has eigenvalues that tend to zero as the gluing parameter tends to zero, but then disappear as the gluing parameter hits zero. This is not really dealt with in [I], but there are now several standard approaches to this problem. We use the modified Sobolev norms of [LT], redefined in the notation of Section 2. The main goal of Section 3 is to describe a correspondence between the elements of $\mathcal{M}_{\Sigma, tv, \lambda}(\mu)$ lying near the maps of type (2) and the zero set of a map between two bundles; see Theorem 3.29. The domain and target bundles will be the bundle of gluing parameters and the obstruction bundle, respectively. We also give a local analytic description of spaces of stable rational maps into V under certain assumptions. If $V = \mathbb{P}^n$, these spaces are very familiar in algebraic geometry, but the description of Theorem 3.33 is used for a transition from the analysis of the gluing problem to the topology of moduli spaces of stable rational maps in [Z2].

Section 4 contains proofs of continuity, injectivity, and surjectivity of the gluing maps. These are usually omitted in the literature, but in the given case one has to choose the obstruction bundle setup carefully to ensure that these properties of the gluing map actually hold. In particular, Section 4 contains what [LT] may mean by “asymptotic analysis near the nodes,” which they omit. The Appendix deals with even more technical details of the analysis.

1.3 Fundamental Notation

In this subsection, we collect the most frequently used combinatorial and analytic notation.

Definition 1.1 *A finite partially ordered set I is a linearly ordered set if for all $i_1, i_2, h \in I$ such that $i_1, i_2 < h$, either $i_1 \leq i_2$ or $i_2 \leq i_1$.*

A linearly ordered set I is a rooted tree if I has a unique minimal element, i.e. there exists $\hat{0} \in I$ such that $\hat{0} \leq h$ for all $h \in I$.

If I and I' are linearly ordered sets, a bijection $\phi: I \rightarrow I'$ is an isomorphism of linearly ordered sets if for all $h, i \in I$, $i < h$ if and only if $\phi(i) < \phi(h)$.

Let I be a linearly ordered set. We denote the subset of the non-minimal elements of I by $\hat{I}$, i.e.

$$\hat{I} = \{h \in I : i < h \text{ for some } i \in I\}.$$

For every $h \in \hat{I}$, the set $\{i \in I : i < h\}$ has a unique maximal element ι_h , i.e.

$$\iota_h < h \quad \text{and} \quad i \leq \iota_h \text{ for all } i \in I \text{ s.t. } i < h.$$

For reasons made clear in Subsection 2.1, $\iota: \hat{I} \rightarrow I$ will be called the attaching map of I . It is clear from Definition 1.1 that I has a unique splitting $I = \bigsqcup_{k \in K} I_k$ such that $I_k \subset I$ is a rooted tree.

The attaching map of I restricts to the attaching map of each I_k , which will still be denoted by ι .

Let I be a rooted tree. We denote the unique minimal element of I by $\hat{0}_I$, or simply by $\hat{0}$ if there is no ambiguity. If I^* , I_* , and I^* are rooted trees, we will write $\hat{I}^*$, $\hat{I}_*$, and $\hat{I}^*$ for $\widehat{I^*}$, $\widehat{I_*}$, and $\widehat{I^*}$, respectively; here $*$ denotes any string of symbols. If $i \in I$, let

$$D_i I = \{h \in I : h > i\}, \quad \bar{D}_i I = D_i I \cup \{i\}.$$

Every rooted tree I has a number of subsets that are rooted trees; the subsets $\bar{D}_i I$ are one example. If H is a subset of I , the set

$$I^{(H)} \equiv \{i \in I : i \not\prec h \ \forall h \in H\}$$

is also a rooted tree. If $i \in I$, denote $I^{\{\hat{i}\}}$ by $I^{(i)}$. If H is a subset of $\hat{I}$, let

$$I^H = \{i \in I : i \not\prec h \ \forall h \in H\}, \quad I(H) = H \cup \{\hat{0}_I\}.$$

If $h \in \hat{I}$, denote $I^{\{h\}}$ by I^h .

If M_1 and M_2 are two sets, let $M_1 + M_2$ be the disjoint union of M_1 and M_2 . Finally, if N is a nonnegative integer, let $[N] = \{1, \dots, N\}$.

We now introduce some analytic notation. Let $\beta : \mathbb{R} \rightarrow [0, 1]$ be a smooth function such that

$$\beta(t) = \begin{cases} 0, & \text{if } t \leq 1; \\ 1, & \text{if } t \geq 2, \end{cases} \quad \text{and} \quad \beta'(t) > 0 \quad \text{if } t \in (1, 2). \quad (1.3)$$

If $r > 0$, let $\beta_r \in C^\infty(\mathbb{R}; \mathbb{R})$ be given by $\beta_r(t) = \beta(r^{-\frac{1}{2}}t)$. Note that

$$\text{supp}(\beta_r) = [r^{\frac{1}{2}}, 2r^{\frac{1}{2}}], \quad \|\beta'_r\|_{C^0} \leq C_\beta r^{-\frac{1}{2}}, \quad \text{and} \quad \|\beta''_r\|_{C^0} \leq C_\beta r^{-1}. \quad (1.4)$$

Throughout the paper, β and β_r will refer to these smooth bump functions.

Let $q_N, q_S : \mathbb{C} \rightarrow S^2 \subset \mathbb{R}^3$ be the stereographic projections mapping the origin in $\mathbb{C}$ to the north and south poles, respectively. Explicitly,

$$q_N(z) = \left(\frac{2z}{1+|z|^2}, \frac{1-|z|^2}{1+|z|^2} \right) \in \mathbb{C} \times \mathbb{R}, \quad q_S(z) = \left(\frac{2z}{1+|z|^2}, \frac{-1+|z|^2}{1+|z|^2} \right). \quad (1.5)$$

We denote the south pole of S^2 , i.e. the point $(0, 0, -1) \in \mathbb{R}^3$, by ∞ . Let

$$e_\infty = (0, 0, 1) = dq_S \Big|_0 \left(\frac{\partial}{\partial s} \right) \in T_\infty S^2, \quad (1.6)$$

where we write $z = s + it \in \mathbb{C}$. We identify $\mathbb{C}$ with $S^2 - \{\infty\}$ via the map q_N . If $x, z \in S^2 - \{\infty\}$, define $\phi_x z \in \mathbb{C}$ by

$$\phi_x z = z - x \equiv q_N^{-1}(z) - q_N^{-1}(x). \quad (1.7)$$

Note that the map $\phi_x : S^2 - \{\infty\} \rightarrow \mathbb{C}$ is biholomorphic. If g is a Riemannian metric on a Riemann surface (Σ, j) of positive genus, $x \in \Sigma$ and $v \in T_x \Sigma$, we write $\exp_{g,x} v \in \Sigma$ for the exponential

of v defined with respect to the Levi-Civita connection of g . Let $\text{inj}_g x$ denote the corresponding injectivity radius at x and d_g the distance function. If $x, z \in \Sigma$ are such that $d_g(x, z) < \text{inj}_g x$, define $\phi_{g,x} z \in T_x \Sigma$ by

$$\exp_{g,x} \phi_{g,x} z = z, \quad |\phi_{g,x} z|_{g,x} < \text{inj}_g x. \quad (1.8)$$

Note that if g is flat on a neighborhood U of x in Σ , then $\phi_{g,x}|_U$ is holomorphic.

Let g_V be the Kahler metric of (V, J, ω) . Denote the corresponding Levi-Civita connection, exponential map, and distance function by ∇^V , $\exp_V$ and d_V , respectively. For every $\lambda \in H_2(V; \mathbb{Z})$, let $|\lambda| = \langle \omega, \lambda \rangle$. The number $|\lambda|$ is the g_V -energy of any element of $\mathcal{H}_{\Sigma, \lambda}$; see [MS]. By rescaling ω , it can be assumed that $|\lambda| \geq 1$, whenever $\lambda \neq 0$ and $\mathcal{H}_{S^2, \lambda} \neq \emptyset$. If g is any Kahler metric on (V, J) , denote the corresponding Levi-Civita connection, exponential map, distance function, injectivity radius, and the parallel transport along the geodesic for $X \in TV$ by ∇^g , $\exp_g$, d_g , inj_g , and $\Pi_{g,X}$, respectively. If $q \in V$ and $\delta \in \mathbb{R}$, let $B_g(q, \delta) = \{q' \in V : d_g(q, q') \leq \delta\}$. In our construction, we allow g vary in a smooth family. Without causing any additional difficulty in the gluing construction, consideration of such families simplifies computations in specific cases such as in [Z2]. If (S, j) is a smooth Riemann surface and $u \in C^\infty(S; V)$, put

$$\begin{aligned} \Gamma(u) &= \Gamma(S; u^* TV), \quad \Gamma^1(u) = \Gamma(S; T^* S \otimes u^* TV); \\ \Gamma^{0,1}(u) &= \Gamma(S; \Lambda^{0,1} T^* S \otimes u^* TV), \quad \bar{\partial} u = \frac{1}{2} (du + J \circ du \circ j) \in \Gamma^{0,1}(u). \end{aligned}$$

We denote by D_V and D_g the linearization of $\bar{\partial}$ -operator with respect to the metrics g_V and g on V , respectively. Since both metrics are Kahler, D_V and D_g commute with J and have no zeroth-order term; see [Z1].

It should be mentioned that it is not essential for the main gluing construction described in this paper that (V, J, g) is Kahler or even symplectic. If (V, J, g) is not Kahler, we would need to choose an orientation on certain spaces of holomorphic maps and take the induced orientation on the cokernel bundle; see Subsection 3.2. Dropping the Kahler assumption would have almost no effect on the analysis, but would slightly complicate the notation.

2 Spaces of Bubble Maps

2.1 Bubble Trees

Let S be either the Riemann sphere S^2 or a smooth Riemann surface Σ of genus at least 2. Allowing the genus-one case would lead to somewhat more complicated notation, but would have no effect on the analysis done in Section 3. Denote by S^* the open subset $S - \{\infty\}$ if $S = S^2$ and S itself if $S = \Sigma$.

Definition 2.1 A bubble tree based on S is a tuple $\mathbb{T} = (S, I; x)$, where

- (1) I is a rooted tree and $x: \hat{I} \longrightarrow S \cup S^2$ is a map;
- (2) $x_h \in S^*$ if $\iota_h = \hat{0}$ and $x_h \in S^2 - \{\infty\}$ otherwise;
- (3) if $h_1 \neq h_2$ and $\iota_{h_1} = \iota_{h_2}$, $x_{h_1} \neq x_{h_2}$.

Given a bubble tree $\top$ as above, let $\Sigma_\top$ be the complex curve

$$\Sigma_\top = \left((\{\hat{0}\} \times S) \sqcup \bigsqcup_{h \in \hat{I}} (\{h\} \times S^2) \right) / \sim,$$

where $(h, \infty) \sim (\iota_h, x_h)$ for $h \in \hat{I}$. The subset $\Sigma_{\top, \hat{0}} \equiv \{\hat{0}\} \times S$ of $\Sigma_\top$ will be called the *principal component* of $\top$ or $\Sigma_\top$. If $h \in \hat{I}$, let $\Sigma_{\top, h} = \{h\} \times S^2$. For every $i \in I$, $\Sigma_{\top, i}$ will be called the *i*th *bubble component* of $\top$ or $\Sigma_\top$ or simply a *bubble component*. Let $\Sigma_{\top, i}^*$ and $\Sigma_\top^*$ denote the open subsets of smooth points of $\Sigma_{\top, i}$ and $\Sigma_\top$, respectively, i.e.

$$\Sigma_{\top, i}^* = \begin{cases} \Sigma_{\top, i} - \{(i, \infty)\} - \{(i, x_h) : \iota_h = i\}, & \text{if } i \in \hat{I}; \\ S - \{(\hat{0}, x_h) : \iota_h = \hat{0}\}, & \text{if } i = \hat{0}; \end{cases} \quad \Sigma_\top^* = \bigcup_{i \in I} \Sigma_{\top, i}^*.$$

The complement of $\Sigma_{\top, i}^*$ in $\Sigma_\top$ are the *singular points* of $\Sigma_\top$.

For every $i \in I$, let $\top^{(i)} = (S, I^{(i)}; x|_{\hat{I}^{(i)}})$. Similarly, for each $h \in \hat{I}$, let $\top^h = (S, I^h; x|_{\hat{I}^h})$. These tuples are again bubble trees based on S . The complex curve $\Sigma_{\top^{(i)}}$ is obtained from $\Sigma_\top$ by dropping all bubble components descendent from the *i*th bubble component. The curve $\Sigma_{\top^h}$ is obtained by dropping the *h*th bubble component along with all bubble components descendent from it.

If $S = S^2$ and $h \in \hat{I}$, we denote the map ϕ_{x_h} defined in (1.7) by $\phi_{\top, h}$. If $z \in \Sigma_{\top, i}$, put

$$r_{\top, h}(z) = \begin{cases} |\phi_{\top, h} z|, & \text{if } i = \iota_h \text{ and } z \neq \infty; \\ 100, & \text{otherwise.} \end{cases} \quad (2.1)$$

If $\delta > 0$, let $B_{\top, h}(\delta) = \{z \in \Sigma_\top : r_{\top, h}(z) < \delta\}$. Put

$$r_\top = \min_{h \in \hat{I}} \left(|q_S^{-1}(x_h)|, \min\{r_{\top, h}(\iota_l, x_l) : l \neq h\} \right). \quad (2.2)$$

If $S = \Sigma$ and $h \in \hat{I}$ is such that $\iota_h \in \hat{I}$, we again let ϕ_{x_h} denote the function ϕ_{x_h} of (1.7) and define $r_{\top, h}$ and $B_{\top, h}(\delta)$ as above. If g is a Riemannian metric on Σ , $\iota_h = \hat{0}$, and $z \in \Sigma_{\top, i}$, put

$$r_{\top, g, h}(z) = \begin{cases} d_g(x_h, z), & \text{if } i = \hat{0}; \\ 100, & \text{otherwise.} \end{cases} \quad (2.3)$$

We denote by $\phi_{\top, g, h}$ the function ϕ_{g, x_h} of (1.8) and by $B_{\top, g, h}(\delta)$ the ball $B_g(x_h, \delta)$. Put

$$r_\top g = \min \left(\min_{\iota_h = \hat{0}} \{r_{\top, g, h}(\iota_l, x_l) : l \neq h\}, \min_{\iota_h \neq \hat{0}} \left(|q_S^{-1}(x_h)|, \min\{r_{\top, h}(\iota_l, x_l) : l \neq h\} \right) \right). \quad (2.4)$$

We say g is a $\top$ -admissible Riemannian metric on Σ if there exists $\delta > 0$ such that for all $h \in \hat{I}$ with $\iota_h = \hat{0}$ the metric g is flat on $B_{\top, g, h}(\delta)$.

2.2 The Basic Gluing Construction

In this subsection, we describe a gluing construction on $\top$ which is the basis of all the other gluing constructions in this paper. Lemma 2.2 plays a very important role in the next section and in the

explicit computations of [Z2].

Let $\top = (S, I; x)$ be a bubble tree. If $h \in \hat{I}$, put

$$F_{h,\top}^{(0)} = \begin{cases} \mathbb{C}, & \text{if } x_h \in S^2; \\ T_{x_h} \Sigma, & \text{if } x_h \in \Sigma, \end{cases} \quad F_{\top}^{(0)} = \bigoplus_{h \in \hat{I}} F_{h,\top}^{(0)}. \quad (2.5)$$

If $S = S^2$, for any $\delta > 0$, put

$$F_{\top,\delta}^{(0)} = \left\{ v = (\top, v_{\hat{I}}) : v_{\hat{I}} \in F_{\top}^{(0)}, |v| \equiv \sum_{h \in \hat{I}} |v_h| < \delta \right\}.$$

Let $\delta_{\top} \in (0, 1)$ be such that $8\delta_{\top}^{\frac{1}{2}} < r_{\top}$. If $S = \Sigma$ and g is an admissible metric on Σ , put

$$F_{\top,g,\delta}^{(0)} = \left\{ v = (\top, v_{\hat{I}}) : v_{\hat{I}} \in F_{\top}^{(0)}, |v|_g \equiv \sum_{i_h = \hat{0}} |v_h|_g + \sum_{i_h \neq \hat{0}} |v_h| < \delta \right\},$$

where $|v_h|_g = |v_h|_{g,x_h}$. Let $\delta_{\top} g \in (0, 1)$ be such that $8(\delta_{\top} g)^{\frac{1}{2}} < r_{\top} g$ and the metric g is flat on $B_g(x_h, 4(\delta_{\top} g)^{\frac{1}{2}})$ for all $h \in \hat{I}$ with $i_h = \hat{0}$. We now construct a family of bubble trees parameterized by $F_{\top,\delta}^{(0)}$ with $\delta = \delta_{\top}$ if $S = S^2$ and by $F_{\top,g,\delta}^{(0)}$ with $\delta = \delta_{\top} g$ if $S = \Sigma$.

First, for every $h \in \hat{I}$ and $v_h \in F_{\top,h}^{(0)}$ with

$$|v_h| \in (0, \delta) \quad \text{if } x_h \in S^2 \quad \text{and} \quad |v_h|_g \in (0, \delta) \quad \text{if } x_h \in \Sigma,$$

we define local stretching maps

$$q_{h,(x_h,v_h)} : \Sigma_{\top(\iota_h)} \longrightarrow \Sigma_{\top(h)} \quad \text{if } x_h \in S^2 \quad \text{and} \quad q_{g,h,(x_h,v_h)} : \Sigma_{\top(\iota_h)} \longrightarrow \Sigma_{\top(h)} \quad \text{if } x_h \in \Sigma$$

as follows. If $x_h \in S^2$, let $p_{h,(x_h,v_h)} : B_{\top,h}(\delta^{\frac{1}{2}}) \longrightarrow \mathbb{C} \cup \infty$ be given by

$$p_{h,(x_h,v_h)}(z) = (1 - \beta_{|v_h|}(2|\phi_{\top,h} z|)) \overline{\left(\frac{v_h}{\phi_{\top,h} z} \right)}. \quad (2.6)$$

Define $q_{h,(x_h,v_h)} : \Sigma_{\top(h)} \longrightarrow \Sigma_{\top(h)}$ by

$$q_{h,(x_h,v_h)}(z) = \begin{cases} (h, q_S(p_{h,(x_h,v_h)}(z))), & \text{if } \frac{1}{2} \leq |v_h|^{-\frac{1}{2}} r_{\top,h}(z) \leq 1; \\ (\iota_h, \phi_{\top,h}^{-1}(\beta_{|v_h|}(|\phi_{\top,h} z|)(\phi_{\top,h} z))), & \text{if } 1 \leq |v_h|^{-\frac{1}{2}} r_{\top,h}(z) \leq 2; \\ z, & \text{otherwise.} \end{cases} \quad (2.7)$$

Note that $q_{h,(x_h,v_h)}$ is smooth and is a diffeomorphism, except on the circle $r_{\top,h}(z) = |v_h|^{\frac{1}{2}}$ in $\Sigma_{\top,\iota_h}$. The map $q_{h,(x_h,v_h)}$ stretches $B_{\top,h}(|v_h|^{\frac{1}{2}})$ around the sphere $\Sigma_{\top,h}$. If $x_h \in \Sigma$, similarly to the above, let $p_{g,h,(x_h,v_h)} : B_{\top,g,h}(\delta^{\frac{1}{2}}) \longrightarrow \mathbb{C} \cup \infty$ be given by

$$p_{g,h,(x_h,v_h)}(z) = (1 - \beta_{|v_h|_g}(2|\phi_{\top,g,h} z|_g)) \overline{\left(\frac{v_h}{\phi_{\top,g,h} z} \right)}. \quad (2.8)$$

Note that the ratio $v_h/\phi_{\top,g,h}z$ is well-defined as an extended complex number, since $T_{x_h}\Sigma$ is one-dimensional and $v_h \neq 0$. Define $q_{g,h,(x_h,v_h)}: \Sigma_{\top^h} \rightarrow \Sigma_{\top^{(h)}}$ by

$$q_{g,h,(x_h,v_h)}(z) = \begin{cases} (h, q_S(p_{g,h,(x_h,v_h)}(z))), & \text{if } \frac{1}{2} \leq |v_h|_g^{-\frac{1}{2}} r_{\top,g,h}(z) \leq 1; \\ (\iota_h, \phi_{g,\top,h}^{-1}(\beta_{|v_h|_g}(|\phi_{\top,g,h}z|_g)(\phi_{\top,g,h}z))), & \text{if } 1 \leq |v_h|_g^{-\frac{1}{2}} r_{\top,g,h}(z) \leq 2; \\ z, & \text{otherwise.} \end{cases} \quad (2.9)$$

Similarly to the case $x_h \in S^2$, $q_{g,h,(x_h,v_h)}$ is smooth and is a diffeomorphism, except on the circle $r_{\top,g,h}(z) = |v_h|_g^{\frac{1}{2}}$ in $\Sigma_{\top,i_h}$.

If $S = S^2$, for every $h \in I$ and $v \in F_{\top,\delta}^{(0)}$, we now define a bubble tree $\top_h(v)$ and a smooth map $q_{v,h}: \Sigma_{\top_h(v)} \rightarrow \Sigma_{\top^{(h)}}$. Choose an ordering of I consistent with its partial ordering. If $h = \hat{0}$, we take $I_h(v) = \{\hat{0}\}$, $\top_h(v) = (S, I_h(v);)$, and $q_{v,h} = Id_S$. Suppose $h \neq \hat{0}$ and

$$\top_{h-1}(v) = (S, \hat{I}_{h-1}(v); x_h(v))$$

with $I_{h-1}(v) \subset I$. If $v_h = 0$, put

$$I_h(v) = I_{h-1}(v) \cup \{h\}, \quad (\iota_{h,l}(v), x_{h,l}(v)) = \begin{cases} (\iota_{h-1,l}(v), x_{h-1,l}(v)), & \text{if } l \in I_{h-1}(v); \\ q_{v,\iota_h}^{-1}(\iota_h, x_h), & \text{otherwise.} \end{cases}$$

Let $q_{v,h}|_{\Sigma_{\top_{h-1}(v)}} = q_{v,h-1}$ and $q_{v,h}(h, z) = (h, z)$. If $v_h \neq 0$, let

$$I_h(v) = I_{h-1}(v), \quad (\iota_{h,l}(v), x_{h,l}(v)) = (\iota_{h-1,l}(v), x_{h-1,l}(v)).$$

Take $q_{v,h} = q_{h,(x_h,v_h)} \circ q_{v,h-1}$. Inductively this procedure defines a bubble tree $\top(v) = \top_{h^*}(v)$ based on S and a smooth map $q_v = q_{v,h^*}: \Sigma_{\top(v)} \rightarrow \Sigma_{\top}$, which is a diffeomorphism outside of $|I - I(v)|$ disjoint circles, where h^* is the largest element of I . The resulting bubble tree and map are independent of the choice of the extension of the partial ordering. While the domains of the maps $q_{v,h}$ do depend on such a choice, whenever we make use of the maps $q_{v,h}$ below, the result will also be independent of the choice. If $S = \Sigma$, for every $h \in I$ and $v \in F_{\top,g,\delta}^{(0)}$, we define bubble tree $\top_{g,h}(v)$ and maps $q_{g,v,h}: \Sigma_{\top_{g,h}(v)} \rightarrow \Sigma_{\top^{(h)}}$ similarly to the above, but replacing $q_{h,(x_h,v_h)}$ by $q_{g,h,(x_h,v_h)}$ whenever $\iota_h = \hat{0}$. We let $\top_g(v) = \top_{g,h^*}(v)$ and $q_{g,v} = q_{g,v,h^*}$. As before, $q_{g,v}$ is smooth and a diffeomorphism outside of $|I - I(v)|$ disjoint circles.

If $S = S^2$ and $v_h \neq 0$, put

$$\begin{aligned} A_{v,h}^+ &= q_{v,\iota_h}^{-1} \left(\{z \in \Sigma_{\top,\iota_h} : |v_h|^{\frac{1}{2}} \leq r_{\top,h}(z) \leq 2|v_h|^{\frac{1}{2}}\} \right); \\ A_{v,h}^- &= q_{v,\iota_h}^{-1} \left(\{z \in \Sigma_{\top,\iota_h} : \frac{1}{2}|v_h|^{\frac{1}{2}} \leq r_{\top,h}(z) \leq |v_h|^{\frac{1}{2}}\} \right). \end{aligned} \quad (2.10)$$

Note that $A_{v,h}^{\pm} \subset \Sigma_{\top(v), i_h^*(v)}$, where

$$i_h^*(v) = \min\{i \in I : i < h \text{ and } v_{h'} \neq 0 \text{ if } i < h' < h\} = \max\{i \in I(v) : i < h\}.$$

If $S = \Sigma$ and $v_h \neq 0$, we similarly define

$$\begin{aligned} A_{g,v,h}^+ &= q_{g,v,\iota_h}^{-1} \left(\{z \in \Sigma_{\top, \iota_h} : |v_h|_g^{\frac{1}{2}} \leq r_{\top, g, h}(z) \leq 2|v_h|_g^{\frac{1}{2}} \} \right); \\ A_{g,v,h}^- &= q_{g,v,\iota_h}^{-1} \left(\{z \in \Sigma_{\top, \iota_h} : \frac{1}{2}|v_h|_g^{\frac{1}{2}} \leq r_{\top, g, h}(z) \leq |v_h|_g^{\frac{1}{2}} \} \right), \end{aligned} \quad (2.11)$$

where $|v_h|_g$ and $r_{\top, g, h}$ denote $|v_h|$ and $r_{\top, h}$ if $\iota_h \in \hat{I}$.

Lemma 2.2 *If $S = S^2$, the map q_v is holomorphic outside of the annuli $A_{v,h}^\pm$ with $v_h \neq 0$. For such h ,*

$$\begin{aligned} \|dq_{h,(x_h,v_h)}\|_{C^0(q_{v,\iota_h}(A_{v,h}^\pm))} &\leq C; \\ \bar{\partial}(q_v \circ q_{v,\iota_h}^{-1})|_z &= -2|v_h|^{-\frac{1}{2}} \left(\frac{v_h}{\phi_{\top, h} z} \right) dq_S|_{p_{h,(x_h,v_h)} z} \circ \partial\beta|_{2|v_h|^{-\frac{1}{2}} \phi_{\top, h} z} \circ d\phi_{\top, h}|_z \quad \forall z \in q_{v,\iota_h}(A_{v,h}^-), \end{aligned}$$

where the norm is computed with respect to the standard metric on S^2 , and β is viewed as a function on $\mathbb{C}$ via the standard norm on $\mathbb{C}$. If $S = \Sigma$, the map $q_{g,v}$ is holomorphic outside of the annuli $A_{g,v,h}^\pm$ with $v_h \neq 0$. For such h ,

$$\begin{aligned} \|dq_{g,h,(x_h,v_h)}\|_{C^0(q_{g,v,\iota_h}(A_{g,v,h}^\pm))} &\leq C_g \quad \text{if } \iota_h = \hat{0}; \quad \|dq_{h,(x_h,v_h)}\|_{C^0(q_{g,v,\iota_h}(A_{g,v,h}^\pm))} \leq C \quad \text{if } \iota_h \neq \hat{0}; \\ \bar{\partial}(q_{g,v} \circ q_{g,v,\iota_h}^{-1})|_z &= -2|v_h|^{-\frac{1}{2}} \left(\frac{v_h}{\phi_{\top, h} z} \right) dq_S|_{p_{h,(x_h,v_h)} z} \circ \partial\beta|_{2|v_h|^{-\frac{1}{2}} \phi_{\top, h} z} \circ d\phi_{\top, h}|_z \quad \forall z \in q_{g,v,\iota_h}(A_{v,h}^-), \end{aligned}$$

where we regard β as a function on $T_{x_h}\Sigma$ via the metric g and denote $\phi_{g,\top,h}$ by $\phi_{\top, h}$ if $\iota_h = \hat{0}$.

Proof: The first statement in each of the two cases is immediate from the construction. The estimates on the differential of $q_{h,(x_h,v_h)}$ and $q_{g,h,(x_h,v_h)}$ follow from (1.4). Suppose $S = \Sigma$, $\iota_h = \hat{0}$, $v_h \neq 0$ and $z \in A_{g,v,h}^-$. Since $q_{g,v} = q_{g,h,(x_h,v_h)}$ on $A_{g,v,h}^-$ and q_S is anti-holomorphic, from (2.8) and (2.9) we obtain

$$\begin{aligned} \bar{\partial}q_{g,v}|_z &= dq_S|_{p_{g,h,(x_h,v_h)} z} \circ \partial p_{g,h,(x_h,v_h)}|_z \\ &= -2|v_h|_g^{-\frac{1}{2}} \left(\frac{v_h}{\phi_{\top, g, h} z} \right) dq_S|_{p_{g,h,(x_h,v_h)} z} \circ \partial\beta|_{2|v_h|_g^{-\frac{1}{2}} \phi_{\top, g, h} z} \circ d\phi_{\top, g, h}|_z. \end{aligned}$$

The other cases are proved similarly, since $q_{g,v} \circ q_{g,v,\iota_h}^{-1} = q_{h,(x_h,v_h)}$ on $q_{g,v,\iota_h}(A_{g,v,h}^-)$ and a similar statement holds in the case $S = S^2$.

2.3 Curves with Marked Points

Definition 2.3 *If M is a finite set, a curve with M -marked points based on S is a tuple*

$$\mathcal{C} = (S, M, I; x, (j, y)), \quad \text{where}$$

- (1) $\top_{\mathcal{C}} \equiv (S, I; x)$ is a bubble tree based on S , and $j: M \rightarrow I$ and $y: M \rightarrow S \cup S^2$ are maps;
- (2) $j_l \in I$, $(j_l, y_l) \in \Sigma_{\top_{\mathcal{C}}, j_l}^*$, and $y_l \neq \infty$ for all $l \in M$;
- (3) for any $l_1, l_2 \in M$ with $l_1 \neq l_2$ and $j_{l_1} = j_{l_2}$, $y_{l_1} \neq y_{l_2}$.

The curve $\mathcal{C}$ is stable if $|\{h: \iota_h = i\}| + |\{l: j_l = i\}| \geq 2$ for all $i \in \hat{I}$ if $S = \Sigma$ and all $i \in I$ if $S = S^2$.

Via the construction in Subsection 2.1, such a tuple $\mathcal{C}$ corresponds to a complex curve $\Sigma_{\mathcal{C}} \equiv \Sigma_{\top_{\mathcal{C}}}$ with marked points $\{(j_l, y_l)\}_{l \in M}$. For each $i \in I$, we denote by $\Sigma_{\mathcal{C}, i}$ and $\Sigma_{\mathcal{C}, i}^*$ the surfaces $\Sigma_{\top_{\mathcal{C}}, i}$ and $\Sigma_{\top_{\mathcal{C}}, i}^*$, respectively.

With notation as above, for every $h \in \hat{I}$, let $F_{h, \mathcal{C}}^{(0)}$ and $F_{\mathcal{C}}^{(0)}$ denote the spaces $F_{h, \top_{\mathcal{C}}}^{(0)}$ and $F_{\top_{\mathcal{C}}}^{(0)}$, respectively. If $S = S^2$, put

$$r_{\mathcal{C}} = \min \left(r_{\top_{\mathcal{C}}}, \min_{l \in M} (|q_S^{-1} y_l|, \min\{r_{\top_{\mathcal{C}}, h}(j_l, y_l) : h \in \hat{I}\}, \min\{|\phi_{y_l}^{-1} y_h| : h \neq l, j_h = j_l\}) \right). \quad (2.12)$$

Let $\delta_{\mathcal{C}} \in (0, 1)$ be such that $16(|I| + |M|)\delta_{\mathcal{C}}^{\frac{1}{2}} < r_{\mathcal{C}}$. If $v = (\mathcal{C}, v_{\hat{I}})$ with $v_{\hat{I}} \in F_{\mathcal{C}}^{(0)}$ and $|v| < \delta_{\mathcal{C}}$, we now construct a curve $\mathcal{C}(v)$ with M -marked points as follows. Let

$$\top(v) = (S, I(v); x(v)) \quad \text{and} \quad q_v : \Sigma_{\top(v)} \longrightarrow \Sigma_{\mathcal{C}}$$

be the bubble tree and the smooth map defined in Subsection 2.2. Then we take

$$\mathcal{C}(v) = (S, M, I(v); x(v), (j(v), y(v))),$$

where $(j_l(v), y_l(v)) \in \Sigma_{\top(v), j_l(v)}$ is defined by

$$q_v(j_l(v), y_l(v)) = (j_l, y_l).$$

Similarly, if $S = \Sigma$ and g is an admissible Riemannian metric on Σ , put

$$r_{\mathcal{C}}g = \min \left(r_{\top_{\mathcal{C}}}g, \min_{j_l = \hat{0}} \left(\min_{\iota_h = \hat{0}} \{r_{\top_{\mathcal{C}}, g, h}(j_l, y_l)\}, \min\{|\phi_{g, y_l}^{-1} y_h|_g : h \neq l, j_h = \hat{0}\} \right), \right. \\ \left. \min_{j_l \neq \hat{0}} (|q_S^{-1} y_l|, \min_{\iota_h \neq \hat{0}} \{r_{\top_{\mathcal{C}}, h}(j_l, y_l)\}, \min\{|\phi_{y_l}^{-1} y_h| : h \neq l, j_h = j_l\}) \right). \quad (2.13)$$

Let $\delta_{\mathcal{C}}g \in (0, 1)$ be such that $16(|I| + |M|)(\delta_{\mathcal{C}}g)^{\frac{1}{2}} < r_{\mathcal{C}}g$ and g is flat in $B_g(x_h, 8(\delta_{\mathcal{C}}g)^{\frac{1}{2}})$ for all $h \in \hat{I}$ with $\iota_h = \hat{0}$. If $v \in F_{\mathcal{C}}^{(0)}$ and $|v|_g < \delta_{\mathcal{C}}g$, we construct a curve $\mathcal{C}_g(v)$ with M -marked points in the same way as above, but replacing q_v and $\top(v)$ by $q_{g, v}$ and $\top_g(v)$.

Definition 2.4 *An isomorphism of curves with M -marked points $\mathcal{C} = (S, M, I; x, (j, y))$ and $\mathcal{C}' = (S, M, I'; x', (j', y'))$, is a tuple of maps,*

$$\phi_0 : I \longrightarrow I', \quad \phi_{1, \hat{0}} : S \longrightarrow S, \quad \phi_{1, h} : S^2 \longrightarrow S^2 \quad \text{for } h \in I, \quad \text{where}$$

- (1) ϕ_0 is an isomorphism of the linearly ordered sets I and I' and $\phi_0(j_l) = j'_l$ for all $l \in M$;
- (2) $\phi_{1, i}$ is a biholomorphic map for all $i \in I$ and $\phi_{1, \hat{0}}$ is the identity map if $S = \Sigma$;
- (3) $\phi_{1, i}(\infty) = \infty$ for all $i \in I$ if $S = S^2$ and for all $i \in \hat{I}$ if $S = \Sigma$;
- (4) $\phi_{1, \iota_h}(x_h) = x'_{\phi_0(h)}$ for $h \in \hat{I}$ and $\phi_{1, j_l}(y_l) = y'_l$ for all $l \in M$.

Such a set of maps corresponds to a continuous map $\Sigma_{\mathcal{C}} \longrightarrow \Sigma_{\mathcal{C}'}$ that maps the l th marked point (j_l, y_l) on $\Sigma_{\mathcal{C}}$ to the l th marked point (j'_l, y'_l) on $\Sigma_{\mathcal{C}'}$ and is biholomorphic on each component of $\Sigma_{\mathcal{C}}$. Note that if $\mathcal{C}$ is stable, $\mathcal{C}$ has no nontrivial automorphisms. Let $[\mathcal{C}]$ denote the equivalence class of $\mathcal{C}$ in the set of all curves based on S with marked points. Denote by $\overline{\mathcal{M}}_{S, M}$ the set of all equivalence classes of stable curves based on S with M -marked points. If $S = S^2$, $\overline{\mathcal{M}}_{S, M}$ can be

identified with the moduli space $\overline{\mathcal{M}}_{0,|M|+1}$ of all stable rational curves with $|M| + 1$ marked points, or more canonically with the space $\overline{\mathcal{M}}_{0,M+\{\hat{0}\}}$ of all stable rational curves with the marked points labeled by the set $M + \{\hat{0}\}$. If $S = \Sigma$ has genus bigger than two and is generic, $\overline{\mathcal{M}}_{S,M}$ is the closed subset of $\overline{\mathcal{M}}_{g,M}$ consisting of all stable curves of genus g with M -marked points that have a fixed complex structure on the principal component. If S has genus two, $\overline{\mathcal{M}}_{S,M}$ is a double cover of the corresponding set for $g = 2$, since any smooth genus-two curve has a holomorphic automorphism of order two; see [GH, p254]. The reason we require $\phi_{1,\hat{0}} = Id_\Sigma$ is that the symplectic invariant of [RT] disregards the automorphisms of Σ .

2.4 Bubble Maps

Definition 2.5 A V -valued bubble map is a tuple $b = (S, M, I; x, (j, y), u)$, where

- (1) I is a linearly ordered set, which is a rooted tree if $S = \Sigma$ and $u: I \rightarrow C^\infty(S; V) \cup C^\infty(S^2; V)$ is a map;
- (2) if $I = \bigsqcup_{k \in K} I_k$ is the splitting of I into rooted trees, then $M = \bigsqcup_{k \in K} M_k$ for some subsets M_k of M such that $\mathcal{C}_k = (S, M_k, I_k; x|_{\hat{I}_k}, (j, y)|_{M_k})$ is an M_k -marked curve based on S ;
- (3) $u_h: S \rightarrow V$ if $h \in I - \hat{I}$ and $u_h: S^2 \rightarrow V$ if $h \in \hat{I}$ is a smooth map such that $u_h(\infty) = u_{\iota_h}(x_h)$ for all $h \in \hat{I}$;
- (4) for all $i \in \hat{I}$ if $S = \Sigma$ and $i \in I$ if $S = S^2$,

$$|\{h \in \hat{I}: \iota_h = i\}| + |\{l \in M: j_l = i\}| < 2 \implies u_{i*}[S^2] \neq 0 \in H_2(V; \mathbb{Z}).$$

The bubble map b is simple if I is a rooted tree; b is holomorphic if $\bar{\partial}u_i = 0$ for all $i \in I$.

With notation as in Definition 2.5, every bubble map b corresponds to a continuous map

$$u_b: \Sigma_b \equiv \bigsqcup_{k \in K} \Sigma_{\mathcal{C}_k} \rightarrow V,$$

which is smooth on the components of $\Sigma_{\mathcal{C}_k}$. If $h \in \hat{I}_k$, let $F_{h,b}^{(0)} = F_{h,\mathcal{C}_k}^{(0)}$. Similarly, let

$$F_b^{(0)} = \bigoplus_{k \in K} F_{\mathcal{C}_k}^{(0)}, \quad \Sigma_b^* = \bigcup_{k \in K} \Sigma_{\mathcal{C}_k}^* \subset \Sigma_b, \quad \Sigma_{b,i}^* = \Sigma_{\mathcal{C}_k,i}^* \subset \Sigma_{b,i} \equiv \Sigma_{\mathcal{C}_k,i},$$

whenever $i \in I_k \subset I$. If b is simple, denote by $\top_b$ the bubble tree $\top_{\mathcal{C}_k}$ for the unique element $k \in K$.

Definition 2.6 An isomorphism of V -valued bubble maps $b = (S, M, I; x, (j, y), u)$ and $b' = (S, M, I'; x', (j', y'), u')$ is a tuple of maps

$$\phi_0: I \rightarrow I', \quad \phi_{1,i}: S \rightarrow S \text{ for } i \in I - \hat{I}, \quad \phi_{1,i}: S^2 \rightarrow S^2 \text{ for } i \in \hat{I}, \quad \text{where}$$

- (1) ϕ_0 is an isomorphism of the linearly ordered sets I and I' with $\phi_0(j_l) = j'_l$ for all $l \in M$;
- (2) $\phi_{1,i}$ is a biholomorphic map for all $i \in I$ and is the identity map if $S = \Sigma$ and $i \notin \hat{I}$;
- (3) $\phi_{1,i}(\infty) = \infty$ for all $i \in I$ if $S = S^2$ and for all $i \in \hat{I}$ if $S = \Sigma$;
- (4) $\phi_{1,\iota_h}(x_h) = x'_{\phi_0(h)}$ for all $h \in \hat{I}$ and $\phi_{1,j_l}(y_l) = y'_l$ for all $l \in M$;
- (5) $u'_{\phi_0(i)} \circ \phi_{1,i} = u_i$ for all $i \in I$.

Such a set of maps corresponds to a continuous map $\Sigma_b \longrightarrow \Sigma_{b'}$ that maps the marked points of b to the marked points of b' , intertwines the maps $u_b : \Sigma_b \longrightarrow V$ and $u_{b'} : \Sigma_{b'} \longrightarrow V$, and is biholomorphic on each component $\Sigma_{b,i}$ of Σ_b . Let G_b denote the group of automorphisms of the bubble map b . This group is necessarily finite by the stability condition (4) of Definition 2.5. If $\lambda \in H_2(V; \mathbb{Z})$, let

$$\bar{C}_{(\lambda;M)}^\infty(S;V) = \{b = (S, M, I; x, (j, y), u) \text{ is } V\text{-valued bubble map: } \sum_{i \in I} u_{i*}[\Sigma_{b,i}] = \lambda\} / \sim;$$

$$C_{(\lambda;M)}^\infty(S;V) = \{b = (S, \{\hat{0}\};, (\hat{0}, y), u_{\hat{0}}) \text{ is } V\text{-valued bubble map: } u_{\hat{0}*}[S] = \lambda\} / \sim,$$

where the equivalence relation is given by isomorphisms of V -valued bubble maps. If $\mu = \mu_{\tilde{M}}$ is an $\tilde{M}$ -tuple of submanifolds of V , let

$$\begin{aligned} \bar{C}_{(\lambda;M)}^\infty(S; \mu) &= \{b = [S, M, I; x, (j, y), u] \in \bar{C}_{(\lambda;M)}^\infty(S; V) : u_{j_l}(y_l) \in \mu_l \ \forall l \in M \cap \tilde{M}\}, \\ C_{(\lambda;M)}^\infty(S; \mu) &= \{b = [S, M, \{\hat{0}\};, (\hat{0}, y), u_{\hat{0}}] \in C_{(\lambda;M)}^\infty(S; V) : u_{\hat{0}}(y_l) \in \mu_l \ \forall l \in M \cap \tilde{M}\}. \end{aligned}$$

A topology on $\bar{C}_{(\lambda;M)}^\infty(S;V)$ and its subsets $C_{(\lambda;M)}^\infty(S;V)$, $\bar{C}_{(\lambda;M)}^\infty(S; \mu)$, $C_{(\lambda;M)}^\infty(S; \mu)$ is defined below.

Definition 2.7 Let $b^* = (S, M, I^*; x^*, (j^*, y^*), u^*)$ and $b_k = (S, M, I_k; x_k, (j_k, y_k), u_k)$ be simple bubble maps. If $S = S^2$, the sequence $\{b_k\}$ converges to b^* if for all k sufficiently large one can choose

- (a) M -marked curves $\mathcal{C}_k = (S, M, I^*; x'_k, (j^*, y^*))$, and
 - (b) vectors $(v_k)_{\hat{I}^*} \in F_{\mathcal{C}_k}^{(0)}$ with $16|v_k| < r_{\mathcal{C}_k}^2$,
- such that with $v_k = (\mathcal{C}_k, (v_k)_{\hat{I}^*})$,
- (1) $\lim_{k \rightarrow \infty} x'_{k,h} = x_h^*$ for all $h \in \hat{I}$, and $\lim_{k \rightarrow \infty} |v_k| = 0$;
 - 2) $\mathcal{C}(v_k) = (S, M, I_k; x_k, (j_k, y(v_k)))$,

$$\lim_{k \rightarrow \infty} q_{v_k}(j_{k,l}, y_{k,l}) = (j_l^*, y_l^*) \ \forall l \in M, \quad \text{and} \quad \lim_{k \rightarrow \infty} \sup_{z \in \Sigma_{\mathcal{C}(v_k)}} d_V(u_{b^*}(q_{v_k}(z)), u_{b_k}(z)) = 0.$$

If $S = \Sigma$, convergence is defined in the same way, but $|v_k|$ and $\mathcal{C}(v_k)$ are replaced by $|v_k|_g$ and $\mathcal{C}_g(v_k)$, respectively, for a $\mathbb{T}_{b^*}$ -admissible metric g on Σ .

This notion of convergence is independent of the choice of an admissible metric on Σ . Definition 2.7 induces a topology on the space $\bar{C}_{(\lambda;M)}^\infty(S;V)$, which will be referred to as the Gromov topology.

2.5 Stratums of Bubble Maps

In this subsection, we introduce the notion of a bubble type. We then define various spaces of holomorphic bubble maps indexed by bubble types and vector bundles over them.

Definition 2.8 A bubble type is a tuple $\mathcal{T} = (S, M, I; j, \lambda)$, such that

- (1) I is a linearly ordered set, and $j : M \longrightarrow I$ and $\lambda : I \longrightarrow H_2(S; \mathbb{Z})$ are maps;
- (2) for all $i \in \hat{I}$ if $S = \Sigma$ and all $i \in I$ if $S = S^2$, $\lambda_i \neq 0$ if $|\{h : \iota_h = i\}| + |\{l : j_l = i\}| < 2$.

Bubble type $\mathcal{T}$ is simple if I is a rooted tree; $\mathcal{T}$ is basic if $\hat{I} = \emptyset$.

Two bubble types $\mathcal{T} = (S, M, I; j, \lambda)$ and $\mathcal{T}' = (S, M, I'; j', \lambda')$ are equivalent if there exists an isomorphism of linearly ordered sets $\phi_0 : I \longrightarrow I'$ such that $\phi_0(j_l) = j'_l$ for all $l \in M$ and $\lambda'_{\phi_0(i)} = \lambda_i$

for all $i \in I$.

If $\mathcal{T}^* = (S, M, I^*; j^*, \lambda^*)$ and $\mathcal{T} = (S, M, I; j_M, \lambda)$ are two bubble types, $\underline{\mathcal{T}^*} < \underline{\mathcal{T}}$ if $I \subset I^*$,

$$j_l = \max\{i \in I : i \leq j_l^*\} \text{ for all } l \in M \quad \text{and} \quad \lambda_i = \sum_{i=\max\{i' \leq h : i' \in I\}} \lambda_h^*.$$

If $\mathcal{T} = (S, M, I; j, \lambda)$ is a bubble type, a $\mathcal{T}$ -bubble map is a bubble map $b = (S, M, I; x, (j, y), u)$ such that $u_{i*}[\Sigma_{b,i}] = \lambda_i \in H_2(V; \mathbb{Z})$ for all $i \in I$.

The splitting of I into rooted trees I_k induces a splitting of $\mathcal{T}$ into simple bubble types

$$\mathcal{T}_k = (S, M_k, I_k; j_k, \lambda_k),$$

where j_k and λ_k are the restrictions of j and λ to M_k and I_k , respectively. Similarly, each $\mathcal{T}$ -bubble map b corresponds to a K -tuple of bubble maps $b_K = (b_k)_{k \in K}$, where b_k is a $\mathcal{T}_k$ -bubble map.

We denote the equivalence class of the bubble type $\mathcal{T}$ by $[\mathcal{T}]$ and the groups of automorphisms of $\mathcal{T}$ by $\text{Aut}(\mathcal{T})$. The partial ordering on the set of bubble types induces a partial ordering on the set of their equivalence classes. If b and b' are $\mathcal{T}$ - and $\mathcal{T}'$ -bubble maps, respectively, such that $[b] = [b']$, then $[\mathcal{T}] = [\mathcal{T}']$. Furthermore, if $\{b_k\}$ is a sequence of $\mathcal{T}$ -bubble maps, b^* is $\mathcal{T}^*$ -bubble map, and $[b_k]$ converges to $[b^*]$ with respect to the Gromov topology, then $[\mathcal{T}^*] \leq [\mathcal{T}]$.

Let $\mathcal{T} = (S, M, I; j, \lambda)$ be a bubble type. We denote by $\langle \mathcal{T} \rangle$ the basic bubble type such that $\langle \mathcal{T} \rangle \geq \mathcal{T}$. It can be described explicitly as follows. Let $I = \bigsqcup_{k \in K} I_k$ be the splitting of I into rooted trees and $M = \bigsqcup_{k \in K} M_k$ the corresponding splitting of M ; see Definition 2.5. It can be assumed that $K = I - \hat{I}$ and k is the unique minimum element of I_k . For every $k \in K$ and $l \in M_k$, let

$$\lambda'_k = \sum_{i \in I_k} \lambda_i, \quad j'_l = k.$$

Then $\langle \mathcal{T} \rangle = (S, M, K; j', \lambda')$.

If $\mathcal{T} = (S, M, I; j, \lambda)$ is a simple bubble type and $i \in I$, let $D_i \mathcal{T} = D_i I$ and $\bar{D}_i \mathcal{T} = \bar{D}_i I$. If H is a subset of $\hat{I}$, let $\mathcal{T}(H) = (S, M, H \cup \hat{0}; j', \lambda')$, where

$$j'_l = \max\{i \in H \cup \hat{0} : i \leq j_l\} \quad \text{and} \quad \lambda'_i = \sum_{i_H^* \leq h \leq i} \lambda_h \quad \text{with} \quad i_H^* = \max\{i^* \in H \cup \hat{0} : i^* \leq i\}.$$

Then $\mathcal{T}(H)$ is again a bubble type. The bubble type $\mathcal{T}(H)$ is the bubble type obtained by gluing $\mathcal{T}$ -bubble maps with the parameter $v_{\hat{f}}$ such that $v_h = 0$ if and only if $h \in H$; see the next section. Finally, we will denote by $(\lambda; M)$ the equivalence class of the basic simple bubble type $(S^2, M, \{\hat{0}\}; \hat{0}, \lambda)$.

Given a bubble type $\mathcal{T} = (S, M, I; j, \lambda)$, let $d(\mathcal{T}) : I \rightarrow \mathbb{R}$ be given by

$$d_i(\mathcal{T}) = |\lambda_i| + |\{l \in M : j_l = i\}| + \sum_{\iota_h = i} d_h(\mathcal{T}) \quad \forall i \in I. \quad (2.14)$$

Since I is a linearly ordered set, the numbers $d_i(\mathcal{T})$ are uniquely defined by (2.14). If

$$b = (S, M, I; x, (j, y), u)$$

is a $\mathcal{T}$ -bubble map, b is $\mathcal{T}$ -balanced if for all $i \in \hat{I}$

$$(B1) \int_{\mathbb{C}} |du_i \circ q_N|^2 z + \sum_{\iota_h=i} d_h(\mathcal{T}) x_h + \sum_{j_l=i} y_l = 0;$$

$$(B2) \int_{\mathbb{C}} |du_i \circ q_N|^2 \beta(|z|) + \sum_{\iota_h=i} d_h(\mathcal{T}) \beta(|x_h|) + \sum_{j_l=i} \beta(|y_l|) = \frac{1}{2}.$$

The integrals above are computed with respect to the metric g_V on V . Recall that we consider $\mathbb{C}$ to be a subset of S^2 via the map q_N . Thus, x_h and y_l can be viewed as complex numbers, as done above. If $S = S^2$ and b is as above, b is *completely $\mathcal{T}$ -balanced* (or *cb*) if (B1) and (B2) hold for all $i \in I$.

Denote by $\mathcal{H}_{\mathcal{T}}$ the set of all holomorphic $\mathcal{T}$ -bubble maps. Let

$$PSL_2^{(0)} = \{g \in PSL_2 : g(\infty) = \infty\}, \quad \mathcal{G}_{\mathcal{T}} = \prod_{h \in \hat{I}} PSL_2^{(0)}.$$

The group $\mathcal{G}_{\mathcal{T}}$ acts on $\mathcal{H}_{\mathcal{T}}$ as follows. If

$$b = (S, M, I; x, (j, y), u) \in \mathcal{H}_{\mathcal{T}} \quad \text{and} \quad g = g_{\hat{I}} \in \mathcal{G}_{\mathcal{T}},$$

define $gb = (S, M, I; gx, (j, gy), (gu))$ by

$$(gx)_h = \begin{cases} g_{\iota_h} x_h, & \text{if } \iota_h \in \hat{I}; \\ x_h, & \text{if } \iota_h \notin \hat{I}; \end{cases} \quad (gy)_l = \begin{cases} g_{j_l} y_l, & \text{if } j_l \in \hat{I}; \\ y_l, & \text{if } j_l \notin \hat{I}; \end{cases} \quad (gu)_i = \begin{cases} g_i \cdot u_i, & \text{if } i \in \hat{I}; \\ u_i, & \text{if } i \notin \hat{I}, \end{cases}$$

where for any map $f : S^2 \rightarrow V$ and $g \in PSL_2$, we define

$$g \cdot f : S^2 \rightarrow V \quad \text{by} \quad \{g \cdot f\}(z) = f(g^{-1}z).$$

Let $\mathcal{M}_{\mathcal{T}}^{(0)} \subset \mathcal{H}_{\mathcal{T}}$ denote the subset of $\mathcal{T}$ -balanced holomorphic maps. The group $G_{\mathcal{T}} \times \text{Aut}(\mathcal{T})$, where

$$G_{\mathcal{T}} \equiv \prod_{h \in \hat{I}} S^1 \subset \mathcal{G}_{\mathcal{T}},$$

acts on $\mathcal{M}_{\mathcal{T}}^{(0)}$ and all the stabilizers are finite. Denote the quotient by $\mathcal{M}_{\mathcal{T}}$, and let

$$\bar{\mathcal{M}}_{\mathcal{T}} = \bigcup_{\mathcal{T}' \leq \mathcal{T}} \mathcal{M}_{\mathcal{T}'}$$

If $\text{Aut}(\mathcal{T}) = \{1\}$, corresponding to the quotient $\mathcal{M}_{\mathcal{T}} = \mathcal{M}_{\mathcal{T}}^{(0)} / G_{\mathcal{T}}$, we obtain $|\hat{I}|$ line (orbi)-bundles

$$\{L_h \mathcal{T} \rightarrow \mathcal{M}_{\mathcal{T}} : h \in \hat{I}\},$$

which carry natural norms:

$$|[b, c_h]| = |c_h| \quad \text{if} \quad b \in \mathcal{M}_{\mathcal{T}}^{(0)} \quad \text{and} \quad c_h \in \mathbb{C}.$$

If $\text{Aut}(\mathcal{T}) \neq \{1\}$, the fiber products and connect sums of the above line bundles taken over each orbit of $\text{Aut}(\mathcal{T})$ are well-defined. Let $F_h^{(0)}\mathcal{T} \longrightarrow \mathcal{M}_{\mathcal{T}}^{(0)}$ be the bundle with the fiber $F_{h,b}^{(0)}$ at $b \in \mathcal{M}_{\mathcal{T}}^{(0)}$, i.e.

$$F_h^{(0)}\mathcal{T} = \begin{cases} \mathcal{M}_{\mathcal{T}}^{(0)} \times \mathbb{C}, & \text{if } x_h \in S^2; \\ \pi_h^* T\Sigma, & \text{if } x_h \in \Sigma, \end{cases} \quad \text{where } \pi_h(b) = x_h,$$

with notation as above. The action of $G_{\mathcal{T}}$ on $\mathcal{M}_{\mathcal{T}}^{(0)}$ lifts to an action on each bundle $F_h^{(0)}\mathcal{T}$ by

$$g \cdot (b, v_h) = \begin{cases} (g \cdot b, g_{\iota_h} g_h^{-1} v_h), & \text{if } \iota_h \in \hat{I}; \\ (g \cdot b, g_h^{-1} v_h), & \text{if } \iota_h \notin \hat{I}. \end{cases}$$

Here and in the rest of the paper, we identify S^1 with the unit complex numbers in the usual way. Let $F_h\mathcal{T}$ be the line orbi-bundle over $\mathcal{M}_{\mathcal{T}}$ given by

$$F_h\mathcal{T} = F_h^{(0)}\mathcal{T} / (G_{\mathcal{T}} \times \text{Aut}(\mathcal{T})).$$

This bundle has a natural norm unless $\iota_h = \hat{0}$ and $S = \Sigma$. In such a case, any metric g on Σ induces a norm on $F_h\mathcal{T}$. Let

$$F^{(0)}\mathcal{T} = \bigoplus_{h \in \hat{I}} F_h^{(0)}\mathcal{T}, \quad F_b^{(0)}\mathcal{T} = F^{(0)}\mathcal{T}|b; \quad F\mathcal{T} = \bigoplus_{h \in \hat{I}} F_h\mathcal{T}, \quad F_{[b]}^{(0)}\mathcal{T} = F^{(0)}\mathcal{T}|[b].$$

Note that if $\mathcal{T}^* < \mathcal{T}$, $G_{\mathcal{T}}$ is naturally a subgroup of $G_{\mathcal{T}^*}$ and thus acts on $\mathcal{M}_{\mathcal{T}^*}^{(0)}$ and the line bundles $F_h^{(0)}\mathcal{T}^*$. Furthermore, there is a natural decomposition $G_{\mathcal{T}^*} = G_{\mathcal{T}} \times G$ for a certain group G . In particular, the $G_{\mathcal{T}}$ -action on $\mathcal{M}_{\mathcal{T}}^{(0)}$ and $F_h^{(0)}\mathcal{T}$ induces an action of $G_{\mathcal{T}^*}$.

If $S = S^2$, let

$$\mathcal{B}_{\mathcal{T}} = \{b = (S, M, I; x, (j, y), u) \in \mathcal{H}_{\mathcal{T}} : b \text{ is cb}; u_{i_1}(\infty) = u_{i_2}(\infty) \forall i_1, i_2 \in I - \hat{I}\}.$$

Denote by $\mathcal{U}_{\mathcal{T}}^{(0)} \subset \mathcal{M}_{\mathcal{T}}$ the quotient $\mathcal{B}_{\mathcal{T}} / (G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$. The group

$$G_{\mathcal{T}}^* \equiv \prod_{i \in I - \hat{I}} S^1$$

acts on $\mathcal{U}_{\mathcal{T}}^{(0)}$ and $\mathcal{M}_{\mathcal{T}}$ as follows. If

$$[b] = [(S^2, M, I; x, (j, y), u)] \in \mathcal{M}_{\mathcal{T}} \quad \text{and} \quad g = (g_i)_{i \in I - \hat{I}} \in G_{\mathcal{T}}^*,$$

define $g[b] = [(S^2, M, I; gx, (j, gy), gu)]$ by

$$(gx)_h = \begin{cases} x_h, & \text{if } \iota_h \in \hat{I}; \\ g_{\iota_h} x_h, & \text{if } \iota_h \notin \hat{I}; \end{cases} \quad (gy)_l = \begin{cases} y_l, & \text{if } j_l \in I; \\ g_{j_l} y_l, & \text{if } j_l \notin I; \end{cases} \quad (gu)_i = \begin{cases} u_i, & \text{if } i \in \hat{I}; \\ g_i \cdot u_i, & \text{if } i \notin \hat{I}. \end{cases}$$

As in the previous paragraph, all the stabilizers are finite. Furthermore, this $G_{\mathcal{T}^*}$ -action on $\mathcal{M}_{\mathcal{T}}$ naturally lifts to an action on $\mathcal{M}_{\mathcal{T}}^{(0)}$ and along with the $G_{\mathcal{T}}$ -action on $\mathcal{M}_{\mathcal{T}}^{(0)}$ induces an action of $\tilde{G}_{\mathcal{T}} \equiv G_{\mathcal{T}}^* \times G_{\mathcal{T}}$ on $\mathcal{M}_{\mathcal{T}}^{(0)}$ as well as on $F_h^{(0)}\mathcal{T}$ by

$$(g^*, g) \cdot (b, v_h) = \begin{cases} ((g^*, g) \cdot b, g_{\iota_h} g_h^{-1} v_h), & \text{if } \iota_h \in \hat{I}; \\ ((g^*, g) \cdot b, g_{\iota_h}^* g_h^{-1} v_h), & \text{if } \iota_h \notin \hat{I}. \end{cases}$$

Note that $G_{\mathcal{T}'} = G_{\mathcal{T}}$ whenever $\mathcal{T}' \leq \mathcal{T}$. Let

$$\mathcal{U}_{\mathcal{T}} = \mathcal{U}_{\mathcal{T}}^{(0)} / G_{\mathcal{T}}^*, \quad \overline{\mathcal{U}}_{\mathcal{T}}^{(0)} = \bigcup_{\mathcal{T}' \leq \mathcal{T}} \mathcal{U}_{\mathcal{T}'}^{(0)}, \quad \overline{\mathcal{U}}_{\mathcal{T}} = \bigcup_{\mathcal{T}' \leq \mathcal{T}} \mathcal{U}_{\mathcal{T}'}.$$

With respect to the Gromov topology, the space $\overline{\mathcal{U}}_{\mathcal{T}}^{(0)}$ is Hausdorff and compact if (V, ω, J) is semipositive; see [RT]. Furthermore, $G_{\mathcal{T}}^*$ acts continuously on $\mathcal{U}_{\mathcal{T}'}^{(0)}$ as can be easily seen from Definition 2.7. It follows that $\overline{\mathcal{U}}_{\mathcal{T}}$ is also Hausdorff and compact if (V, ω, J) is semipositive in the quotient topology. Denote by $\{L_i \mathcal{T} \longrightarrow \overline{\mathcal{U}}_{\mathcal{T}} : i \in I - \hat{I}\}$ the line orbi-bundles corresponding to the quotient $\overline{\mathcal{U}}_{\mathcal{T}} = \overline{\mathcal{U}}_{\mathcal{T}}^{(0)} / G_{\mathcal{T}}^*$. Let

$$\mathcal{F}_h \mathcal{T} = \left(F_h \mathcal{T}^{(0)} | \mathcal{B}_{\mathcal{T}} \right) / \tilde{G}_{\mathcal{T}} \longrightarrow \mathcal{U}_{\mathcal{T}}, \quad \mathcal{F}_{h,[b]} \mathcal{T} = \mathcal{F}_h \mathcal{T} | [b]; \quad \mathcal{F} \mathcal{T} = \bigoplus_{h \in \hat{I}} \mathcal{F}_h \mathcal{T}, \quad \mathcal{F}_{[b]} \mathcal{T} = \mathcal{F} \mathcal{T} | [b].$$

The line bundles $\mathcal{F}_h \mathcal{T}$ have natural norms, defined as in the previous paragraph.

If $\mathcal{T} = (S, M, I; j, \lambda)$ is a bubble type and $b = (S, M, I; x, (j, y), u)$ is a $\mathcal{T}$ -bubble map, for any $l \in M$ let $\text{ev}_l : \mathcal{H}_{\mathcal{T}} \longrightarrow V$ be the map given by

$$\text{ev}_l((S, M, I; x, (j, y), u)) = u_{j_l}(y_l).$$

This map descends to the quotients defined above and induces continuous maps on the spaces $\overline{\mathcal{M}}_{\mathcal{T}}$, $\overline{\mathcal{U}}_{\mathcal{T}}^{(0)}$, and $\overline{\mathcal{U}}_{\mathcal{T}}$. If $\mu = \mu_{\tilde{M}}$ is an $\tilde{M}$ -tuple of submanifolds in V , where $\tilde{M}$ is possibly different from M , put

$$\mathcal{H}_{\mathcal{T}}(\mu) = \{b \in \mathcal{H}_{\mathcal{T}} : \text{ev}_l(b) \in \mu_l \ \forall l \in M \cap \tilde{M}\}.$$

Define spaces $\mathcal{M}_{\mathcal{T}}^{(0)}(\mu)$, $\mathcal{M}_{\mathcal{T}}(\mu)$, $\overline{\mathcal{M}}_{\mathcal{T}}(\mu)$, etc. similarly. If $S = S^2$, we define another evaluation map,

$$\text{ev} : \mathcal{B}_{\mathcal{T}} \longrightarrow V \quad \text{by} \quad \text{ev}((S^2, M, I; x, (j, y), u)) = u_{\hat{0}}(\infty),$$

where $\hat{0}$ is any minimal element of I . This map induces continuous maps on the spaces $\overline{\mathcal{U}}_{\mathcal{T}}^{(0)}$ and $\overline{\mathcal{U}}_{\mathcal{T}}$.

3 The Gluing Construction and the Obstruction Bundle

3.1 Summary and Notation

We now present a gluing construction on the spaces $\mathcal{M}_{\mathcal{T}}(\mu)$ such that $\mathcal{H}_{\mathcal{T}}$ is a smooth manifold with the tangent bundle isomorphic to the kernel of the linearization of the $\bar{\partial}$ -operator, as defined below. This is well-known to be the case if the linearization of the $\bar{\partial}$ -operator is surjective; see [MS]. However, surjectivity of the linearization is not a necessary condition; see [Z2] for examples. In fact, there are two main cases of primarily interest to us. The first is when $\mathcal{T} = (S^2, M; j_M, \lambda)$ and the linearization of the $\bar{\partial}$ -operator is indeed surjective. In this case, we give an analytic description of a neighborhood of $\mathcal{U}_{\mathcal{T}}(\mu)$ in $\overline{\mathcal{U}}_{\langle \mathcal{T} \rangle}(\mu)$ for a generic set of constraints μ . The second case is when $\mathcal{T} = (\Sigma, M, I; j, \lambda)$ and the cokernels of the linearization of the $\bar{\partial}$ -operator form a vector bundle over $\mathcal{H}_{\mathcal{T}}$, which will be the analogue of Taubes's obstruction bundle of [T] in the gluing construction below. Using the same analysis as in the first case, we describe any sufficiently nice element of $C_{(\lambda^*; M)}^{\infty}(\Sigma; \mu)$ lying near $\mathcal{M}_{\mathcal{T}}(\mu)$, where $\lambda^* = \sum \lambda_i$. The elements of $\mathcal{M}_{\Sigma, \nu, \lambda^*}(\mu)$ lying near $\mathcal{M}_{\mathcal{T}}(\mu)$

will correspond to the zero set of a certain section of the obstruction bundle.

For our gluing construction, we fix a smooth family $\{g_{V,b} : b \in \mathcal{M}_{\mathcal{T}}\}$ of Kahler metrics on (V, J) . We assume that this family is $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant if $S = \Sigma$ and $(\tilde{G}_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant if $S = S^2$. If $b \in \mathcal{M}_{\mathcal{T}}$, $X, Y \in T_q V$, and $u : (D, j) \longrightarrow V$ is a smooth map from a one-dimensional complex manifold, let

$$\exp_{b,q} X = \exp_{g_{b,q}} X, \quad \nabla^b = \nabla^{g_b}, \quad \Pi_{b,X} Y = \Pi_{g_b,X} Y, \quad D_{b,u} = D_{g_b,u};$$

see Subsection 1.3 for more details. If $S = \Sigma$, we also choose a smooth family

$$\{g_{\mathcal{T},x} : x = (x)_{\{h:\iota_h=\hat{0}\}}; \ x_h \in \Sigma; \ x_{h_1} \neq x_{h_2} \text{ if } h_1 \neq h_2\}$$

of Riemannian metrics on Σ such that each metric $g_{\mathcal{T},x}$ is flat on a neighborhood of x_h in Σ for all $h \in \hat{I}$ with $\iota_h = \hat{0}$. Existence of such a family of metrics is shown in [FO]. It can be assumed that all these metrics have the same volume as the standard metric on S^2 , i.e. 4π . If

$$b = (\Sigma, M, I; x, (j, y), u) \in \mathcal{H}_{\mathcal{T}},$$

let $g_{b,\hat{0}}$ denote the metric $g_{\mathcal{T},(x)_{\{h:\iota_h=\hat{0}\}}}$ on Σ . If $i \in \hat{I}$, we write $g_{b,i}$ for the standard metric on S^2 . Similarly, if $S = S^2$, for all $i \in I$, we write $g_{b,i}$ for the standard metric on S^2 .

If $b = (S, M, I; x, (j, y), u) \in \mathcal{H}_{\mathcal{T}}$, let

$$\begin{aligned} \Gamma'(b) &= \bigoplus_{i \in I} \Gamma(u_i); \quad \Gamma(b) = \Gamma(u_b) = \{\xi_I \in \Gamma'(b) : \xi_h(\infty) = \xi_{\iota_h}(x_h) \ \forall h \in \hat{I}\}; \\ \Gamma^1(b) &= \Gamma^1(u_b) = \bigoplus_{i \in I} \Gamma^1(u_i); \quad \Gamma^{0,1}(b) = \Gamma^{0,1}(u_b) = \bigoplus_{i \in I} \Gamma^{0,1}(u_i). \end{aligned}$$

Define $D_b : \Gamma(b) \longrightarrow \Gamma^{0,1}(b)$ by

$$(D_b \xi_I)_i = D_{b,u_i} \xi_i \quad \forall i \in I.$$

We denote the kernel of operator D_b on $\Gamma(b)$ by $\Gamma_-(b)$. If $\xi \in \Gamma(u_i)$ or $\xi \in \Gamma^1(u_i)$, let $\|\xi\|_{b,C^k}$ and $\|\xi\|_{b,2}$ denote the C^k - and L^2 -norms of ξ computed with respect to the metrics $g_{V,b}$ on V and $g_{b,i}$ on $\Sigma_{b,i}$. If $\xi = \xi_I \in \Gamma'(b)$ or $\xi \in \Gamma^1(b)$, put

$$\|\xi\|_{b,C^k} = \sum_{i \in I} \|\xi_i\|_{b,C^k}, \quad \|\xi\|_{b,2} = \sum_{i \in I} \|\xi_i\|_{b,2}.$$

Let $\pi_{b,-} : \Gamma(b) \longrightarrow \Gamma_-(b)$ be the (L^2, b) -orthogonal projection map.

The space $\mathcal{P}_b \mathcal{T}$ of perturbations of bubble map b is the collection of tuples $\sigma = (\xi_{\hat{I}}; w_{\hat{I}+M})$, where

$$\xi_i \in \Gamma(u_i) \ \forall i \in I, \quad w_h \in F_{h,b}^{(0)} \ \forall h \in \hat{I}, \quad w_l \in \begin{cases} \mathbb{C}, & \text{if } l \in M \ \& \ \Sigma_{b,j_l} = S^2; \\ T_{y_l} \Sigma, & \text{if } l \in M \ \& \ \Sigma_{b,j_l} = \Sigma. \end{cases}$$

If σ is sufficiently small, we define $\exp_b \sigma = (S, M, I; x(\sigma), (j, y(\sigma)), u_{\sigma})$ by

$$x_h(\sigma) = \begin{cases} x_h + w_h, & \text{if } \Sigma_{b,i_h} = S^2; \\ \exp_{g_{b,\hat{0}},x_h} w_h, & \text{if } \Sigma_{b,i_h} = \Sigma; \end{cases} \quad y_l(\sigma) = \begin{cases} y_l + w_l, & \text{if } \Sigma_{b,j_l} = S^2; \\ \exp_{g_{b,\hat{0}},y_l} w_l, & \text{if } \Sigma_{b,j_l} = \Sigma; \end{cases}$$

and $u_{\sigma,i} = \exp_{b,u_i} \xi_i$. If $z \in \Sigma$, let $|v|_b = |v|_{g_{b,\hat{0},x}}$. For consistency, if $v \in \mathbb{C}$, let $|v|_b = |v|$. Along with the (L^2, b) -norm on the vector fields defined above, we obtain an inner-product on the space of tuples σ as above.

In order to get a good description of the spaces $\mathcal{M}_{\mathcal{T}}^{(0)}$ as submanifolds of $\mathcal{H}_{\mathcal{T}}$, we describe an action of an open subset of 0 in $(\mathbb{C} \oplus \mathbb{R})^{\hat{I}}$ on bubble maps and distinguished elements $\sigma_{(b,i)}^{(k)} \in \mathcal{P}_b \mathcal{T}$ that correspond to this action. If $(c, r, \theta) = (c, r, \theta)_{\hat{I}} \in (\mathbb{C} \times \mathbb{R} \times \mathbb{R})^{\hat{I}}$ and b is a bubble map as above, we define

$$(c, r, \theta) \cdot b = (S, M, I; (c, r, \theta)x, (j, (c, r, \theta)y), (c, r, \theta)u)$$

by setting

$$\begin{aligned} ((c, r, \theta)x)_h &= e^{i\theta_{\iota_h}}(1 + r_{\iota_h})(x_h + c_{\iota_h}), & ((c, r, \theta)y)_l &= e^{i\theta_{j_l}}(1 + r_{j_l})(y_l + c_{j_l}), \\ ((c, r, \theta)u)_i(q_N(z)) &= u_i(q_N((1 + r_i)^{-1}e^{-i\theta_i}z - c_i)). \end{aligned}$$

If (c, r, θ) is sufficiently small, $(c, r, \theta) \cdot b$ is again a bubble map, i.e. the maps into V still agree at the nodes, and the nodes and the marked points are still all distinct. In fact, the values of the maps at the nodes or the marked points do not change, i.e.

$$((c, r, \theta)u)_{\iota_h}(((c, r, \theta)x)_h) = u_{\iota_h}(x_h), \quad ((c, r, \theta)u)_h(\infty) = u_h(\infty), \quad ((c, r, \theta)u)_{j_l}(((c, r, \theta)y)_l) = u_{j_l}(y_l).$$

Furthermore, if $b \in \mathcal{H}_{\mathcal{T}}$, $(c, r, \theta) \cdot b \in \mathcal{H}_{\mathcal{T}}$. If b is of type $\mathcal{T}$, the above describes the action of a neighborhood of the identity in $\mathcal{G}_{\mathcal{T}}$ on the space of stable maps of type $\mathcal{T}$. The action by $\mathbb{C}$ corresponds to the translations of $\mathbb{C}$, by the first $\mathbb{R}$ -component to dilations about the origin, and by the last $\mathbb{R}$ -component to rotations about the origin. In addition, if $S = S^2$, $\mathcal{T}$ is simple, and $(c, r, \theta) \in \mathbb{C} \times \mathbb{R} \times \mathbb{R}$ is sufficiently small, we can define $(c, r, \theta) \cdot b$ similarly by viewing (c, r, θ) as $(c_{\hat{0}}, r_{\hat{0}}, \theta_{\hat{0}})$ in the above definition of the action.

If $u \in C^\infty(S^2; V)$, define $\xi_u^{(1)}, \dots, \xi_u^{(4)} \in \Gamma(u)$ by:

$$\begin{aligned} \xi_u^{(1)}(q_N(z)) &= -d(u \circ q_N) \Big|_z \frac{\partial}{\partial s}, & \xi_u^{(2)}(q_N(z)) &= -d(u \circ q_N) \Big|_z \frac{\partial}{\partial t} \\ \xi_u^{(3)}(q_N(z)) &= -d(u \circ q_N) \Big|_z \left(s \frac{\partial}{\partial s} + t \frac{\partial}{\partial t} \right) = -rd(u \circ q_N) \Big|_z \frac{\partial}{\partial r}, \\ \xi_u^{(3)}(q_N(z)) &= d(u \circ q_N) \Big|_z \left(t \frac{\partial}{\partial s} - s \frac{\partial}{\partial t} \right) = -d(u \circ q_N) \Big|_z \frac{\partial}{\partial \theta}. \end{aligned}$$

where we write $z = s + it \in \mathbb{C}$ and $r = \sqrt{s^2 + t^2}$. These vector fields extend smoothly by zero over the south pole. For any $x \in S^2 - \{\infty\}$, let $w_x^{(1)}, \dots, w_x^{(4)} \in \mathbb{C}$ be given by

$$w_x^{(1)} = 1, \quad w_x^{(2)} = i, \quad w_x^{(3)} = x, \quad w_x^{(4)} = ix.$$

If b is a bubble map as above, $k = 1, \dots, 4$, $i^* \in \hat{I}$ if $S = \Sigma$ and $i^* \in \hat{I}$ if $S = S^2$, let

$$\sigma_{(b,i^*)}^{(k)} = ((\xi_{(b,i^*)}^{(k)})_I, (w_{(b,i^*)}^{(k)})_{\hat{I}+M})$$

be given by

$$\xi_{(b,i^*),i}^{(k)} = \begin{cases} \xi_{u_i}^{(k)}, & \text{if } i = i^*; \\ 0, & \text{if } i \neq i^*; \end{cases} \quad w_{(b,i^*),h}^{(k)} = \begin{cases} w_{x_h}^{(k)}, & \iota_h = i^*; \\ 0, & \iota_h \neq i^*; \end{cases} \quad w_{(b,i^*),l}^{(k)} = \begin{cases} w_{y_l}^{(k)}, & j_l = i^*; \\ 0, & j_l \neq i^*. \end{cases}$$

The tuples $\sigma_{(b,i^*)}^{(k)}$ correspond to the infinitesimal action of $\mathcal{G}_{\mathcal{T}}$ on the space of stable maps of type $\mathcal{T}$.

Finally, if X is any space, $F \longrightarrow X$ a normed vector bundle, and $\delta: X \longrightarrow \mathbb{R}$ is any function, let

$$F_{\delta} = \{(b, v) \in F : |v|_b < \delta(b)\}.$$

Similarly, if Ω is a subset of F , let $\Omega_{\delta} = F_{\delta} \cap \Omega$. If $v = (b, v) \in F$, denote by b_v the image of v under the bundle projection map, i.e. b in this case.

3.2 The Basic Setup

In this subsection, we describe our assumptions on the smooth structure of $\mathcal{H}_{\mathcal{T}}$ and state some of their implications.

Definition 3.1 *Bubble type $\mathcal{T} = (S^2, M, I; j_M, \lambda)$ is (V, J) -regular if for all*

$$b = (S, M, I; x, (j, y), u) \in \mathcal{H}_{\mathcal{T}}$$

- (1) $D_{b, u_i}: \Gamma(u_i) \longrightarrow \Gamma^{0,1}(u_i)$ is onto for all $i \in I$;
- (2) $\ker D_{b, u_i} \longrightarrow T_{u_i(\infty)}V$, $\xi \longrightarrow \xi(\infty)$, is onto for all $i \in I$.

Definition 3.2 *Simple bubble type $\mathcal{T} = (S, M, I; j, \lambda)$ is (V, J) -semiregular if*

- (1) *the space $\mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}})}$ is a complex manifold of the same dimension as $\ker D_{g_V, b}$ for all $b \in \mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}})}$, and there exist $\delta, C \in C^{\infty}(\mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}); \mathbb{R}^+)$ and for each $b = (S, \emptyset, \{\hat{0}\};, u_0) \in \mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}})}$ a map*

$$h_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}), b}: \{\xi \in \ker D_{g_V, b} : \|\xi\|_{g_V, C^0} < \delta(b)\} \longrightarrow \Gamma(u_0)$$

such that

$$\|h_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}), b}(\xi)\|_{g_V, b} \leq C(b) \|\xi\|_{g_V, b}^2, \quad \|h_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}), b}(\xi) - h_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}), b}(\xi')\|_{g_V, C^0} \leq C(b) \|\xi - \xi'\|_{g_V, C^0},$$

for all $\xi, \xi' \in \ker D_{g_V, b}$ with $\|\xi\|_{g_V, C^0}, \|\xi'\|_{g_V, C^0} < \delta(b)$ and the map

$$H_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}), b}: \{\xi \in \ker D_{g_V, b} : \|\xi\|_{g_V, C^0} < \delta(b)\} \longrightarrow \mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}), b}, \quad \xi \longrightarrow \exp_{g_V, u_0}(\xi + h_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}), b}(\xi))$$

is an orientation-preserving diffeomorphism onto an open neighborhood of b in $\mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}})}$. Furthermore, the family of maps $\{H_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}}), b} : b \in \mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}})}\}$ is continuous.

- (2) for all $b = (S, M, I; x, (j, y), u) \in \mathcal{H}_{\mathcal{T}}$

- (2a) $D_{b, u_h}: \Gamma(u_h) \longrightarrow \Gamma^{0,1}(u_h)$ is onto for all $h \in \hat{I}$;
- (2b) $\ker D_{b, u_h} \longrightarrow T_{u_h(\infty)}V$, $\xi \longrightarrow \xi(\infty)$, is onto for all $h \in \hat{I}$.

Remarks: (1) All conditions in both definitions above are independent of the choice of metric on V .

(2) Condition (1) of Definition 3.2 says that $\mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}})}$ is a smooth manifold modeled on $\ker D_b$ for $b \in \mathcal{H}_{(S, \emptyset, \{\hat{0}\};, \lambda_{\hat{0}})}$, as would be the case if $D_b: \Gamma(u_b) \longrightarrow \Gamma^{0,1}(u_b)$ were surjective.

(3) The conditions of Definitions 3.1 and 3.2 insure that $\mathcal{H}_{\mathcal{T}}$ is a smooth manifold; see Proposition 3.3 below. However, (2) of Definition 3.1 and (2b) of Definition 3.2 are somewhat stronger than necessary to show that $\mathcal{H}_{\mathcal{T}}$ is smooth. They allow us to obtain the second part of (1) in

Proposition 3.3, which is used in the proof of surjectivity of the gluing map; see Subsection 4.3. These two conditions hold for all complex homogeneous manifolds; see [RT].

Note that if $\mathcal{T}$ is semiregular, the homotopy invariance of the index implies that the vector spaces

$$\Gamma_-(b) \equiv \text{coker } D_b \approx \ker D_b^* \subset \Gamma^{0,1}(b), \quad b \in \mathcal{H}_{\mathcal{T}},$$

form a vector bundle over $\mathcal{H}_{\mathcal{T}}$. Here D_b^* denotes the formal adjoint of D_b with respect to a metric g on S ; it is a J -linear operator. The space $\ker D_b^*$ is independent of a conformal choice of the metric g . The bundle $\Gamma_- \rightarrow \mathcal{H}_{\mathcal{T}}$ will be called the $\mathcal{T}$ -cokernel bundle. It is $G_{\mathcal{T}}$ -equivariant, and thus descends to a bundle $\Gamma_- \rightarrow \mathcal{M}_{\mathcal{T}}$, which will be the analogue of Taubes's obstruction in our gluing setting.

Let $\mathcal{T} = (S, M, I; j, \lambda)$ be a bubble type. If $b = (S, M, I; x, (j, y), u) \in \mathcal{H}_{\mathcal{T}}$, put

$$\begin{aligned} \mathcal{K}_b \mathcal{T} = \left\{ \sigma = (\xi, w_{\hat{I}+M}) \in \mathcal{P}_b \mathcal{T} : \xi_i \in \ker(D_{b, u_i}) \ \forall i \in I; \ \langle \sigma, \sigma_{(b, h)}^{(k)} \rangle = 0 \ \forall h \in \hat{I}, k \in [4]; \right. \\ \left. \xi_h(\infty) = \xi_{\iota_h}(x_h) + du_{\iota_h}|_{x_h} w_h \ \forall h \in \hat{I} \right\}. \end{aligned}$$

If $\sigma = (\xi, w_{\hat{I}+M}) \in \mathcal{K}_b \mathcal{T}$, let

$$\|\sigma\|_{b, C^k} = \|\xi\|_{b, C^k} + \sum_{h \in \hat{I}} |w_h|_b + \sum_{l \in M} |w_l|_b.$$

We take the default norm on $\mathcal{K}_b \mathcal{T}$ be given by $\|\cdot\|_{b, C^0}$. If b is as above, $b = (S, M, I; x', (j, y'), u)$, and $\delta > 0$, we say $d(b, b') < \delta$ if there exists $\sigma \in \mathcal{P}_b \mathcal{T}$ such that $\exp_b \sigma = b'$ and $\|\sigma\|_{b, C^0} \leq \delta$.

Proposition 3.3 (1) If $\mathcal{T} = (S, M, I; j, \lambda)$ is a regular or semiregular bubble type, $\mathcal{H}_{\mathcal{T}}$ is a complex manifold and there exists $\epsilon_{\mathcal{T}}, C_{\mathcal{T}} \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ with the following property. If $b^* \in \mathcal{H}_{\mathcal{T}}$ and

$$b = (S, M, I; x, (j, y), u) \quad \text{is s.t.} \quad d(b^*, b) < \epsilon_{\mathcal{T}}(b^*) \quad \text{and} \quad \bar{\partial} u_i = 0 \ \forall i \in I,$$

there exist $\xi_i \in \Gamma(u_i)$ for $i \in \hat{I}$ such

$$\|\xi_i\|_{g_V, C^0} \leq C_{\mathcal{T}}(b^*) \sum_{h \in \hat{I}} d_V(u_{\iota_h}(x_h), u_h(\infty)) \quad \text{and} \quad b' = (S, M, I; x, (j, y), u') \in \mathcal{H}_{\mathcal{T}},$$

where $u'_0 = u_0$ and $u'_i = \exp_{g_V, u_i} \xi_i$ if $i \in \hat{I}$.

(2) The space $\mathcal{M}_{\mathcal{T}}^{(0)}$ is a smooth oriented manifold on which the group $G_{\mathcal{T}}$ acts smoothly. The maps

$$\begin{aligned} \text{ev} : \mathcal{M}_{\mathcal{T}}^{(0)} &\longrightarrow V, & \text{ev}(S, M, I; x, (j, y), u) &= u_0(\infty), \\ \text{ev}_l : \mathcal{M}_{\mathcal{T}}^{(0)} &\longrightarrow V, & \text{ev}_l(S, M, I; x, (j, y), u) &= u_{j_l}(y_l), \\ du_i|_z : \mathcal{M}_{\mathcal{T}}^{(0)} &\longrightarrow T^* \Sigma_{\mathcal{T}, i} \otimes u_i^* TV, & du_i|_z(S, M, I; x, (j, y), u) &= du_i|_z, \end{aligned}$$

are smooth. In particular, $u_i \rightarrow \|du_i\|_{b, C^0}$ defines a continuous function on $\mathcal{M}_{\mathcal{T}}^{(0)}$.

(3) There exist $\delta_{\mathcal{T}}, C_{\mathcal{T}} \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ and smooth maps

$$h_{\mathcal{T}, b} = h_{\mathcal{T}, b}^{(1)} \oplus h_{\mathcal{T}, b}^{(2)} : \mathcal{K}_b \mathcal{T}_{\delta_{\mathcal{T}}(b)} \longrightarrow \Gamma'(b) \oplus (\mathbb{C} \oplus \mathbb{R})^{\hat{I}},$$

such that $\|h_{\mathcal{T},b}(\sigma)\|_{b,C^0} \leq C_{\mathcal{T}}(b)\|\sigma\|_{b,C^0}^2$,

$$\|h_{\mathcal{T},b}(\sigma) - h_{\mathcal{T},b}(\sigma')\|_{b,C^0} \leq C_{\mathcal{T}}(b) (\|\sigma\|_{b,C^0} + \|\sigma'\|_{b,C^0}) \|\sigma - \sigma'\|_{b,C^0},$$

and each map

$$\begin{aligned} H_{\mathcal{T},b}^{(0)}: \{(\sigma, \theta) \in \mathcal{K}_b \mathcal{T}_{\delta_{\mathcal{T}}(b)} \times \mathbb{R}^{\hat{I}}: |\theta| < \pi\} &\longrightarrow \mathcal{M}_{\mathcal{T}}^{(0)}, \\ H_{\mathcal{T},b}^{(0)}(b, \sigma, \theta) &= (h_{\mathcal{T},b}^{(2)}(\sigma), \theta) \cdot \exp_b(\sigma + h_{\mathcal{T},b}^{(1)}(\sigma)), \end{aligned}$$

is orientation-preserving diffeomorphism onto an open neighborhood of b in $\mathcal{M}_{\mathcal{T}}^{(0)}$.

Proof: (1) Let $\mathcal{T}_i = (\Sigma_{\mathcal{T},i}, \{l: j_l = i\} + \{h: \iota_h = i\}, \{\hat{0}\}; \hat{0}, \lambda_i)$. By (1) of Definition 3.1, and (1) and (2a) of Definition 3.2, $\mathcal{H}_{\mathcal{T}_i}$ is a complex manifold for all $i \in I$. Let

$$\Delta_V^{\hat{I}} = \left\{ (q, q)_{\hat{I}} \in \prod_{\hat{I}} (V \times V): q_h \in V \right\}.$$

The submanifold $\Delta_V^{\hat{I}}$ is the $\hat{I}$ -product of the diagonal in $V \times V$. Since V is oriented, so is the normal bundle of $\Delta_V^{\hat{I}}$. Claim (1) of the proposition follows by applying the Implicit Function Theorem, (2) of Definition 3.1 and (2b) of Definition 3.2 to the smooth map

$$\text{ev}_{\hat{I}}: \prod_{i \in I} \mathcal{H}_{\mathcal{T}_i} \longrightarrow \prod_{\hat{I}} (V \times V), \quad \text{ev}_h((S, M, I; x, (j, y), u)) = (u_h(\infty), u_{\iota_h}(x_h)).$$

Note that $\mathcal{H}_{\mathcal{T}} = \text{ev}_{\hat{I}}^{-1}(\Delta_V^{\hat{I}})$.

(2) For any $u \in C^\infty(S^2; V)$, define $\tilde{\Psi}u \in \mathbb{C}$, $\Psi^{(3)}u \in \mathbb{R}$, and $\Psi u \in \mathbb{C} \times \mathbb{R}$ by

$$\Psi u = (\tilde{\Psi}u, \Psi^{(3)}u) = \left(\int_{\mathbb{C}} |du \circ q_N|^2 z, \int_{\mathbb{C}} |du \circ q_N|^2 \beta(|z|) - \frac{1}{2} \right),$$

where the integrals are computed using the metric g_V . For $i^* \in \hat{I}$ if $S = \Sigma$ and $i^* \in I$ if $S = S^2$, we define maps

$$\begin{aligned} \Psi_{\mathcal{T},i^*}: \prod_{i \in I} \mathcal{H}_{\mathcal{T}_i} &\longrightarrow \mathbb{C} \times \mathbb{R} \quad \text{by} \\ \Psi_{\mathcal{T},i^*}(S, M, I; x, (j, y), u) &= \left(\tilde{\Psi}u_{i^*} + \sum_{\iota_h=i^*} d_h(\mathcal{T})x_h + \sum_{j_l=i^*} y_l, \Psi^{(3)}u_{i^*} + \sum_{\iota_h=i^*} d_h(\mathcal{T})\beta(|x_h|) + \sum_{j_l=i^*} \beta(|y_l|) \right). \end{aligned}$$

These maps $\Psi_{\mathcal{T},i^*}$ are smooth, since the smooth structure on all $\mathcal{H}_{\mathcal{T}_i}$ is described similarly to (1) of Definition 3.2. Furthermore, if $b \in \mathcal{M}_{\mathcal{T}}^{(0)}$, $i^* \in \hat{I}$, and $k^* = 1, 2, 3$, since $\Psi_{\mathcal{T},i}(b) = 0$ for all i and β' does not change sign, by Lemma 3.4,

$$d\Psi_{\mathcal{T},i^*}^{(k^*)} \Big|_b \sigma_{(b,i)}^{(k)} \begin{cases} = 0, & \text{if } i \neq i^*; \\ \neq 0, & \text{if } i = i^*, k = k^*; \\ = 0, & \text{if } k \neq k^* \neq 3 \end{cases}$$

where $k = 1, 2, 3$. By (2) of Definition 3.1 and (2b) of Definition 3.2, it follows that the map

$$\prod_{i \in I} \mathcal{H}_{\mathcal{T}_i} \longrightarrow (\mathbb{C} \times \mathbb{R})^{\hat{I}} \times \prod_{\hat{I}} (V \times V), \quad b \longrightarrow \left((\Psi_{\mathcal{T}, i}(b))_{i \in \hat{I}}, \text{ev}_{\hat{I}}(b) \right)$$

is transversal to the submanifold $\{0\} \times \Delta_{\hat{V}}^{\hat{I}}$. The preimage of this submanifold is precisely the space $\mathcal{M}_{\mathcal{T}}^{(0)}$. Thus, $\mathcal{M}_{\mathcal{T}}^{(0)}$ is a smooth oriented manifold by the Implicit Function Theorem.

Lemma 3.4 *For any $k \in [4]$ and $u \in C^\infty(S^2; V)$, $\xi^{(k)}(\infty) = 0$. Furthermore,*

$$\tilde{\Psi}((c, r, \theta) \cdot u) = (1 + r)(\tilde{\Psi}u + c\|du\|_2^2) \quad \forall (c, r) \in \mathbb{C} \times \mathbb{R}; \quad (3.1)$$

$$\left. \frac{d}{dr} \Psi^{(3)}((0, r, \theta) \cdot u) \right|_{r=0} = \int_{\mathbb{C}} |d(u \circ q_N)|^2 \beta'(|z|)|z|, \quad (3.2)$$

where $(c, r) \cdot u$ is defined as in Section 3.1. Finally, $D_u \xi_u^{(k)} = 0$ if $\bar{\partial}u = 0$.

Proof: The first and last statements are immediate. To prove (3.1), we use the change of $z \longrightarrow (1 + r)^{-1}z - c$.

$$\begin{aligned} \int_{\mathbb{C}} |d(((c, r) \cdot u) \circ q_N)|^2 z &= \int_{\mathbb{C}} (1 + r)^{-2} |d(u \circ q_N)|_{(1+r)^{-1}z - c}^2 \\ &= (1 + r) \int_{\mathbb{C}} |d(u \circ q_N)|^2 (z + c) = (1 + r)(\tilde{\Psi}u + c\|du\|_2^2), \end{aligned}$$

Similarly,

$$\begin{aligned} \left. \frac{d}{dr} \int_{\mathbb{C}} |d((r \cdot u) \circ q_N)|^2 \beta(|z|) \right|_{r=0} &= \left. \frac{d}{dr} \int_{\mathbb{C}} |d(u \circ q_N)|^2 \beta((1 + r)|z|) \right|_{r=0} \\ &= \int_{\mathbb{C}} |d(u \circ q_N)|^2 \beta'(|z|)|z|. \end{aligned}$$

The lemma is now proved, since the action by the θ -component does not change $\tilde{\Psi}$.

If $\mathcal{T} = (S^2, M, I; j, \lambda)$ is a regular bubble type, with notation as above, let

$$\tilde{\mathcal{K}}_b \mathcal{T} = \{ \sigma = (\xi_I, w_{M+\hat{I}}) \in \mathcal{K}_b \mathcal{T} : \langle \sigma, \sigma_{(b, \hat{0})}^k \rangle = 0 \quad \forall k \in [4], \quad \xi_{i_1}(\infty) = \xi_{i_1}(\infty) \quad \forall i_1, i_2 \in I - \hat{I} \}.$$

By (3) of Definition 3.1 and the same argument as in the proof of Proposition 3.3, we can construct smooth maps $h_{\mathcal{T}, b}^{(1)} \times h_{\mathcal{T}, b}^{(2)} : \tilde{\mathcal{K}}_b \mathcal{T}_{\delta(b)} \longrightarrow \Gamma'(b) \times (\mathbb{C} \times \mathbb{R})^I$ such that each map

$$\begin{aligned} H_{\mathcal{T}, b}^{(0)} : \{ (b, \sigma, \theta) \in \mathcal{K}_b \tilde{\mathcal{T}}_{\delta(b)} \times \mathbb{R}^I : |\theta| < \pi \} &\longrightarrow \mathcal{B}_{\mathcal{T}}, \\ H_{\mathcal{T}, b}^{(0)}(\sigma, \theta) &= (h_{\mathcal{T}, b}^{(2)}(\sigma), \theta) \cdot \exp_b(\sigma + h_{\mathcal{T}, b}^{(1)}(\sigma)), \end{aligned}$$

is orientation-preserving diffeomorphism onto an open neighborhood of b in $\mathcal{B}_{\mathcal{T}}$.

3.3 Construction of Nearly Holomorphic Bubble Maps

Let $\mathcal{T} = (S, M, I; j, \lambda)$ be a simple bubble type. In this subsection, for all $b \in \mathcal{M}_{\mathcal{T}}^{(0)}$ and $v = (b, v_{\hat{I}})$ with $v_{\hat{I}} \in F_b^{(0)}\mathcal{T}$ sufficiently small, we construct a bubble map $b(v)$ with domain Σ_v , where Σ_v is as Subsection 2.2. The map $u_{b(v)}$ will be just the composite $u_b \circ q_v$. We then define a Riemannian metric $g_{v,i}$ and a nonnegative function $\rho_{v,i}$ on each component $\Sigma_{v,i}$ of Σ_v . The metrics will be such that the C^0 -norm of the differential of q_v is bounded independently of $v_{\hat{I}}$. The nonnegative functions are used to modify the Sobolev norms, in such a way that the norm of the inverse of the operator $D_{b(v)}$ on certain subspaces of $\Gamma(b(v))$ is bounded independently of $v_{\hat{I}}$.

By Proposition 3.3, $\mathcal{M}_{\mathcal{T}}^{(0)}$ is a smooth manifold. If $S = S^2$, let $\delta_{\mathcal{T}} \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ be a $\tilde{G}_{\mathcal{T}}$ -invariant function such that $\delta_{\mathcal{T}}(b) < r_{\top_b}$ for all $b \in \mathcal{M}_{\mathcal{T}}^{(0)}$. If $S = \Sigma$, let $\delta_{\mathcal{T}} \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ be a $G_{\mathcal{T}}$ -invariant function such that for all

$$b = (\Sigma, M, I; x, (j, y), u) \in \mathcal{M}_{\mathcal{T}}^{(0)}$$

(A1) $4\delta_{\mathcal{T}}$ is smaller than the function δ of Lemma 5.1;

(A2) $4\delta_{\mathcal{T}}(b) < r_{\mathcal{C}_b} g_{b, \hat{0}}$.

In both cases, it can be assumed that $\delta_{\mathcal{T}}$ does not exceed $\frac{1}{4}$.

If H is a subset of $\hat{I}$, put

$$\begin{aligned} F^{(H)}\mathcal{T} &= \{v = (b, v_{\hat{I}}) \in F^{(0)}\mathcal{T} : v_h = 0 \text{ if and if } h \in H\}, \\ F^H\mathcal{T} &= \{v = [b, v_{\hat{I}}] \in F\mathcal{T} : v_h = 0 \text{ if and if } h \in H\}. \end{aligned}$$

For any $v = (b, v_{\hat{I}}) \in F^{(0)}\mathcal{T}$, let $|v|$ denote $|v|_{g_b}$ if $S = \Sigma$. From now on, we assume that $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ is a $G_{\mathcal{T}}$ -equivariant function if $S = \Sigma$ and a $\tilde{G}_{\mathcal{T}}$ -equivariant function if $S = S^2$ such that $8\delta^{\frac{1}{2}} \leq \delta_{\mathcal{T}}$. If

$$v = (b_v, v_{\hat{I}}) = ((S, M, I; x, (j, y), u), v_{\hat{I}}) \in F^{(0)}\mathcal{T}_{\delta},$$

let $q_v : \Sigma_v \longrightarrow \Sigma_{b_v}$ be the smooth map defined in Subsection 2.2 for

$$v = (\mathcal{C}, v_{\hat{I}}) = ((S, M, I; x, (j, y)), v_{\hat{I}}),$$

using the metric $g_{b_v, \hat{0}}$ on Σ if $S = \Sigma$. Let $u_v = u_{b_v} \circ q_v$ and $b(v) = (\mathcal{C}(v), u_v)$.

We now define a Riemannian metric $g_{v,i}$ on $\Sigma_{v,i}$ for each $i \in I(v) \subset I$. Along the way, we construct a metric $g_{v,i}$ on $\Sigma_{b_v,i}$ for each $i \in I$. Suppose $i \in I$ and for all $h \in \hat{I}$ such that $\iota_h = i$, we have constructed a metric $g_{v,h}$ on $\Sigma_{b_v,h}$. For each $h \in \hat{I}$ such that $\iota_h = i$ and $v_h \neq 0$, let $\tilde{g}_{v,i,h}$ denote the metric on $B_{b_v,h}(2\delta(b_v)^{\frac{1}{2}})$ which is the pullback of the metric $g_{v,h}$ by the map

$$z \longrightarrow q_N\left(\frac{\phi_{b_v,h}}{v_h}\right), \quad \text{where} \quad \phi_{b_v,h} = \begin{cases} \phi_{\top_{b_v},h}, & \text{if } x_h \in S^2; \\ \phi_{\top_{b_v},g_b,h}, & \text{if } x_h \in \Sigma. \end{cases}$$

This metric is conformal with the original metric $g_{b_v,i}$ on $\Sigma_{b_v,i}$, because the maps $\phi_{b,h}$ are holomorphic on the set $\{r_{b,h} \leq \delta_{\mathcal{T}}(b)\}$ and the metric $g_{v,h}$ is conformal with the standard metric on $\mathbb{C}$. Thus,

there exists a smooth positive function $\lambda_{v,i,h}$ such that $\tilde{g}_{v,i,h} = \lambda_{v,i,h}^2 g_{b_v,i}$. Let $\lambda_{v,i} \in C^\infty(\Sigma_{b_v,i}; \mathbb{R}^+)$ be given by

$$\lambda_{v,i}(z) = \begin{cases} \lambda_{v,i,h}(z) + \beta_{|v_h|}(r_{b_v,h}(z))(1 - \lambda_{v,i,h}(z)), & \text{if } \iota_h = i \text{ and } r_{b_v,h}(z) \leq 2|v_h|^{\frac{1}{2}}; \\ 1, & \text{if } r_{b_v,h}(z) \geq 2|v_h|^{\frac{1}{2}} \quad \forall h \in \hat{I}. \end{cases}$$

Since I is a rooted tree, this procedure defines metrics $g_{v,i}$ for each $i \in I(v)$.

In addition, we define a smooth nonnegative function $\rho_{v,i}$ on $\Sigma_{v,i}$ for each $i \in I(v)$. As in the previous paragraph, along the way we define a function $\rho_{v,i}$ for each $i \in I$. Suppose $i \in I$ and for all $h \in \hat{I}$ such that $\iota_h = i$, we have constructed a smooth function $\rho_{v,h}$ on $\Sigma_{b_v,h}$. For $h \in \hat{I}$ with $\iota_h = i$ and $z \in \Sigma_{b_v,i}$ with $|z|_h \equiv r_{b_v,h}(z) \leq 2\delta_{\mathcal{T}}(b_v)$, put

$$\rho_{v,i}(z) = \begin{cases} \rho_{v,h}(q_{h,v_h}z) + \beta\left(\frac{\delta_{\mathcal{T}}(b_v)|z|_h}{|v_h|}\right)\left\{\left(|z|_h^2 + \frac{|v_h|^2}{|z|_h^2}\right) - \tilde{\rho}_{v,h}(q_{h,v_h}z)\right\}, & \text{if } |z|_h \leq \delta_{\mathcal{T}}(b_v); \\ \left(|z|_h^2 + \frac{|v_h|^2}{|z|_h^2}\right) + \beta\left(\frac{|z|_h}{\delta_{\mathcal{T}}(b_v)}\right)\left\{1 - \left(|z|_h^2 + \frac{|v_h|^2}{|z|_h^2}\right)\right\}, & \text{if } |z|_h \geq \delta_{\mathcal{T}}(b_v), \end{cases}$$

if $v_h \neq 0$, where q_{h,v_h} is defined as in Section 2.3, using the metric $g_{b_v,\hat{0}}$ on Σ if $S = \Sigma$. If $v_h = 0$ and z is as above, let

$$\rho_{v,i}(z) = |z|_h^2 + \beta\left(\frac{|z|_h}{\delta_{\mathcal{T}}(b_v)}\right)\{1 - |z|_h^2\}.$$

If $|z|_h \geq 2\delta_{\mathcal{T}}(b_v)$ for all $h \in \hat{I}$ with $\iota_h = i$ and $v_i \neq 0$ if $i > 0$, set $\rho_{v,i}(z) = 1$. Otherwise, let

$$\rho_{v,i}(z) = |q_S^{-1}(z)|^2 + \beta(\delta_{\mathcal{T}}(b_v)|q_S^{-1}(z)|)\{1 - |q_S^{-1}(z)|^2\}.$$

This construction defines nonnegative functions $\rho_{v,i}$ on $\Sigma_{v,i}$ for all $i \in I(v)$.

We finally define norms on the spaces $\Gamma(u_v)$ and $\Gamma^1(u_v)$. If $\eta_i \in \Gamma^1(u_{v,i})$, put

$$2\|\eta_i\|_{v,p;i} = \left(\int_{\Sigma_{v,i}} |\eta_i|^p\right)^{\frac{1}{p}} + \left(\int_{\Sigma_{v,i}} \rho_{v,i}^{-\frac{p-2}{p}} |\eta_i|^2\right)^{\frac{1}{2}}, \quad (3.3)$$

where $|\eta_i|$ and the integrals are computed with respect to the metric $g_{v,i}$ on $\Sigma_{v,i}$ and g_{V,b_v} on V . Denote by $\|\eta_i\|_{v,C^0;i}$ the C^0 -norm of η_i with respect to these metrics. If $\eta = \eta_I \in \Gamma^1(u_v)$, let

$$\|\eta\|_{v,p} = \sum_{i \in I(v)} \|\eta_i\|_{v,p;i}, \quad \|\eta\|_{v,C^0} = \sum_{i \in I(v)} \|\eta_i\|_{v,C^0;i}.$$

Similarly, for any $\xi_i \in \Gamma(u_{v,i})$, put

$$2\|\xi_i\|_{v,p;i} = \left(\int_{\Sigma_{v,i}} |\xi_i|^p\right)^{\frac{1}{p}} + \left(\int_{\Sigma_{v,i}} \rho_{v,i}^{-\frac{p-2}{p}} |\xi_i|^2\right)^{\frac{1}{2}}; \quad \|\xi\|_{v,p,1;i} = \|\xi_i\|_{v,p;i} + \|\nabla \xi_i\|_{v,p;i}, \quad (3.4)$$

where we again use the metrics $g_{v,i}$ on $\Sigma_{v,i}$ and g_{V,b_v} on V as in (3.3). Denote by $\|\xi_i\|_{v,C^0;i}$ the C^0 -norm of ξ_i with respect to the metric g_{V,b_v} on V . If $\xi = \xi_I \in \Gamma(u_v)$, let

$$\|\xi\|_{v,p} = \sum_{i \in I(v)} \|\xi_i\|_{v,p;i}, \quad \|\xi\|_{v,p,1} = \sum_{i \in I(v)} \|\xi_i\|_{v,p,1;i}, \quad \|\xi\|_{v,C^0} = \sum_{i \in I(v)} \|\xi_i\|_{v,C^0;i}.$$

Note that even though the functions $\rho_{v,i}^{-\frac{p-2}{p}}$ have poles at the singular points of Σ_v , all smooth one-forms and vector fields have finite norms defined by (3.3) and (3.4), respectively, since $\frac{p-2}{p} < 1$. We denote by $L_1^p(v)$ the completion of $\Gamma(u_v)$ with respect to the $(v, p, 1)$ -norm and by $L^p(v)$ the completion of $\Gamma^{0,1}(u_v)$ with respect to the (v, p) -norm. Finally, let

$$D_v : \Gamma(u_v) \longrightarrow \Gamma^{0,1}(u_v)$$

denote the linearization of the $\bar{\partial}$ -operator at u_v with respect to the metric g_{V,b_v} on V .

Lemma 3.5 *If $\mathcal{T}$ is a simple bubble type and $p > 2$, there exist $\delta, C \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$,*

- (1) $\|du_v\|_{v,C^0} \leq C(b_v)$ and $\|\bar{\partial}u_v\|_{v,p} \leq C(b_v)|v|^{\frac{1}{p}};$
- (2) $\|D_v\xi\|_{v,p} \leq C(b_v)\|\xi\|_{v,p,1}$ for all $\xi \in \Gamma(u_v);$
- (3) $\|\xi\|_{v,C^0} \leq C(b_v)\|\xi\|_{v,p,1}$ for all $\xi \in \Gamma(u_v);$
- (4) $\|\xi\|_{v,p,1} \leq C(b_v)(\|D_v\xi\|_{v,p} + \|\xi\|_{v,p})$ for all $\xi \in \Gamma(u_v).$

Proof: If $h \in I - I(v)$ and $S = S^2$, let $A_{v,h}^\pm$ be the annulus as in Subsection 2.2. If $S = \Sigma$, let $A_{v,h}^\pm$ denote $A_{g_{b_v},v,h}^\pm$. By definition of the norms, q_v is an isometry outside of such annuli, and by Lemma 2.2 the C^0 -norm of dq_v is bounded on such annuli independently of v_f . Thus, the first part of (1) follows from (2) of Proposition 3.3. Since $\rho_v \geq |v_h|$ on $A_{v,h}$, the second part of (1) follows from Lemma 2.2. Statement (2) of the lemma is immediate from the definition of the norms. The last two claims are proved in the Appendix; see Proposition 5.7 and 5.11. In fact, the C^0 -norm of ξ is bounded by the usual L_1^p -norm of ξ .

3.4 Scale of Variations

In Subsection 3.6, we consider perturbations of the bubble maps $\{b(v)\}$ in directions “away” from the space of such bubble maps. More precisely, we look at replacing u_v by $\exp_{b_v, u_v} \xi$ with ξ lying in a certain subspace of $L_1^p(v)$ complementary to “the tangent space” of the space of maps $\{b(v)\}$. If $\mathcal{T}$ is regular, one obvious candidate for such a subspace is the (L^2, v) -orthogonal complement of the kernel of D_v . While the construction in Subsection 3.6 would go through, we would run into significant difficulty showing injectivity and surjectivity of the gluing map; see Subsections 4.2 and 4.5. In this subsection, we start by describing a choice of the complementary subspace which will work for the purposes of Subsections 3.6, 4.2, and 4.5. We then describe norms on the tangent spaces to $F\mathcal{T}$ and the properties of our setup that are sufficient to show injectivity and surjectivity of the gluing map.

Suppose $v = ((S, M, I; x, (j, y), u), v) \in F^{(0)}\mathcal{T}_\delta$, where $\mathcal{T}$ is a simple bubble type as before. For any $\xi \in \Gamma(b_v)$, define $R_v\xi \in L_1^p(v)$ by

$$\{R_v\xi\}(z) = \xi(q_v(z)).$$

Note that $R_v\xi$ is smooth outside of the $|I - I(v)|$ circles mapped by q_v to the nodes of Σ_{v_b} and is continuous everywhere, since $\Gamma(b_v)$ is the set of smooth vector fields on the components of Σ_{b_v} that agree at the nodes. It follows that $R_v\xi$ is indeed of class L_1^p . Let $\Gamma_-(v)$ be the image of $\ker(D_{b_v})$ under the map R_v . This space models the “tangent bundle” to the space of maps $\{b(v)\}$. Denote by $\Gamma_+(v)$ its (L^2, g_v) -orthogonal complement in $L_1^p(v)$. Let $\pi_{v,-}$ and $\pi_{v,+}$ be the (L^2, g_v) -orthogonal

projections onto $\Gamma_-(v)$ and $\Gamma_+(v)$, respectively.

With $H \subset \hat{I}$ and $v \in F^{(H)}\mathcal{T}_\delta$, let

$$\begin{aligned} T_v F^H \mathcal{T} &= \{ \varpi = (\xi, w_{\hat{I}+M}, \theta_{\hat{I}}, r_{\hat{I}-H}) : (\xi, w_{\hat{I}+M}) \in \mathcal{K}_{b_v} \mathcal{T}; \theta_h, r_h \in \mathbb{R} \}; \\ \tilde{T}_v F^H \mathcal{T} &= \{ (\xi, w_{\hat{I}+M}, \theta_{\hat{I}}, r_{\hat{I}-H}) \in T_v F^H \mathcal{T} : w_h = 0 \ \forall h \in H \}. \end{aligned}$$

Given ϖ as above, put

$$\|\varpi\| = \|\xi\|_{b_v, C^0} + \sum_{h \in \hat{I}} |w_h|_{b_v} + \sum_{l \in M} |w_l|_{b_v} + \sum_{h \in \hat{I}} |\theta_h| + \sum_{h \in \hat{I}-H} |r_h|.$$

If $\delta_{\mathcal{T}}$ and $H_{\mathcal{T}, b_v}^{(0)}$ are as in Proposition 3.3 and $\|\varpi\| < \delta_{\mathcal{T}}(b_v)$, put

$$\begin{aligned} b_\varpi &\equiv (S, M, I; x(\varpi), (j, y(\varpi)), u(\varpi)) = H_{\mathcal{T}, b_v}^{(0)}(\xi, w_{\hat{I}+M}; \theta_{\hat{I}}) \in \mathcal{M}_{\mathcal{T}}^{(0)}, \\ v_h(\varpi) &= \begin{cases} (1+r_h) \begin{cases} v_h, & \text{if } x_h \in S^2; \\ d\phi_{b_v, h}^{-1} | \phi_{b_v, h} x_h(\varpi) v_h, & \text{if } x_h \in \Sigma; \end{cases} & \text{if } h \notin H; \\ 0, & \text{if } h \in H; \end{cases} \quad v(\varpi) \equiv (b_\varpi, (v(\varpi))_{\hat{I}}). \end{aligned}$$

Then $v(\varpi) \in F^{(H)}\mathcal{T}_{2\delta}$ if $\|\varpi\| < \delta(b_v)$ for some $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ sufficiently small. If $H = \emptyset$, $T_v F^H \mathcal{T} = \tilde{T}_v F^H \mathcal{T}$ models the tangent space of $[v]$ in $F^H \mathcal{T}$. If $H \neq \emptyset$, the bundle $F^H \mathcal{T}$ and the construction in the previous subsection lift to a bundle ${}^H F \mathcal{T}$ over

$$\mathcal{M}_{\mathcal{T}}^H \equiv \mathcal{M}_{\mathcal{T}}^{(0)} / \{ g_{\hat{I}} \in G_{\mathcal{T}} : g_h = 1 \ \forall h \in H \}.$$

Then $T_v F^H \mathcal{T}$ models the tangent space of $[v]$ in ${}^H F \mathcal{T}$. On the other hand, $\tilde{T}_v F^H \mathcal{T}$ models the tangent space of $[v]$ in the restriction of ${}^H F \mathcal{T}$ to the subspace

$$\{ [b' = (S, M, I; x', (j, y'), u)] \in \mathcal{M}_{\mathcal{T}}^H : x'_h = x_h \ \forall h \in H \}.$$

The reason for defining subspaces $\tilde{T}_v F^H \mathcal{T}$ is that if $x'_h \neq x_h$ for some $h' \in H$, $b(v)$ and $b(v')$ do not have the same singular points for all $v \in F_b^{(H)} \mathcal{T}$ and $v \in F_{b'}^{(H)} \mathcal{T}$. Since the perturbation construction of Subsection 3.6 does not change the singular points of $b(v)$ and $b(v')$, the resulting bubble maps $\tilde{b}(v)$ and $\tilde{b}(v')$ will necessarily be different.

We now define norms on $T_v F^H \mathcal{T}$, which make the estimates in Lemma 3.6 dependent only on b_v . If $h \in \hat{I} - H$, let

$$w'_h = \phi_{b_v, h} q_{v(\varpi), \iota_h} (q_{v, \iota_h}^{-1}(\iota_h, x_h)) \in F_{h, b_v}^{(0)} \quad \text{if } q_{v(\varpi), \iota_h} (q_{v, \iota_h}^{-1}(\iota_h, x_h)) \in \Sigma_{b_v, \iota_h}.$$

In such a case, let $\|\varpi\|_{v, h} = \left| \frac{w'_h + w_h}{v_h} \right|$. Otherwise, put $\|\varpi\|_{v, h} = 1$. Let $\|\varpi\|_v = \|\varpi\| + \sum_{h \in \hat{I}-H} \|\varpi\|_{v, h}$.

In order to simplify notation, we replace $v(\varpi)$ by ϖ whenever there is no ambiguity. If $\|\varpi\|_v$ is sufficiently small, define $\zeta_\varpi \in \Gamma'(u_v)$ by

$$\exp_{b_v, u_v} \zeta_\varpi = u_\varpi, \quad \|\zeta_\varpi\|_{b_v, C^0} < \inf g_{V, b_v}.$$

Similarly, for $h \in H$ and $l \in M$, define $w_h(\varpi) \in T_{x_h(v)} \Sigma_{v, \iota_h(v)}$ and $w_l(\varpi) \in T_{y_l(v)} \Sigma_{v, j_l(v)}$ by

$$\begin{aligned} \exp_{g_v, x_h(v)} w_h(\varpi) &= x_h(v(\varpi)), & |w_h(\varpi)| &\equiv |w_h(\varpi)|_{g_v} < \text{inj}_{x_h(v)} g_v; \\ \exp_{g_v, y_l(v)} w_l(\varpi) &= y_l(v(\varpi)), & |w_l(\varpi)| &\equiv |w_l(\varpi)|_{g_v} < \text{inj}_{y_l(v)} g_v. \end{aligned}$$

If $\varpi \in \tilde{T}F_v^H \mathcal{T}$ and $\xi \in \Gamma(u_v)$, let $R_\varpi \xi \in \Gamma(u_\varpi)$ be the vector field given by

$$R_\varpi \xi(z) = \Pi_{b_v, \zeta_\varpi(z)} \xi(z).$$

Note that since $b(v)$ and $b(\varpi)$ have the same singular points whenever $\varpi \in \tilde{T}F_v^H \mathcal{T}$, Π_{b_v, ζ_ϖ} does indeed map $\Gamma(u_v)$ to $\Gamma(u_\varpi)$. If $\eta \in \Gamma^1(u_v)$, we define $R_\varpi \eta \in \Gamma^1(u_\varpi)$ similarly. Let S_ϖ denote the inverse of R_ϖ .

Lemma 3.6 *There exist $\delta, C \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R}^+)$ such that for all $v \in F^{(H)} \mathcal{T}_\delta$ and $\varpi \in \tilde{T}_v F^H \mathcal{T}_\delta$,*

- (1) $C(b_v)^{-1} \|\varpi\|_v \leq \|\zeta_\varpi\|_{v,p,1} + \sum_{h \in H} |w_h(\varpi)|_{g_v} + \sum_{l \in M} |w_l(\varpi)|_{g_v} \leq C(b_v) \|\varpi\|_v$;
- (2) $\left\| \frac{g_{V, b_\varpi}}{g_{V, b_v}} - 1 \right\|_{C^3} \leq C(b_v) \|\varpi\|$, $\left\| \frac{g_\varpi}{g_v} - 1 \right\|_{C^0} \leq C(b_v) \|\varpi\|_v$ and $\left\| \frac{\rho_\varpi}{\rho_v} - 1 \right\|_{C^0} \leq C(b_v) \|\varpi\|_v$;
- (3) $\|S_\varpi du_\varpi - du_v\|_{v,p} \leq C(b_v) \|\varpi\|_v$ and $\|S_\varpi \bar{\partial} u_\varpi - \bar{\partial} u_v\|_{v,p} \leq C(b_v) |v|^{\frac{1}{p}} \|\varpi\|_v$;
- (4) $\|S_\varpi \nu - \nu\|_{v,p} \leq C(b_v) \|\varpi\|_v$;
- (5) $\|S_\varpi D_\varpi R_\varpi \xi - D_v \xi\|_{v,p} \leq C(b_v) \|\varpi\|_v \|\xi\|_{v,p,1}$ and $\|S_\varpi \pi_{\varpi, \pm} R_\varpi \xi - \pi_{v, \pm} \xi\|_{v,p,1} \leq C(b_v) \|\varpi\|_v \|\xi\|_{v,p,1}$ for all $\xi \in \Gamma(u_v)$.

Proof: The first statement of (2) is clear. Proofs of (1), the last two claims of (2), (3), and the last claim of (4) are direct, though lengthy, computations, all of the same nature. The statement of (4) is immediate from (1). The first claim of (5) follows from (2) and basic Riemannian geometry estimates as in [Z1].

Remark: The second of Claim (5) above is proved by choosing an orthonormal basis $\{\xi_{b,i}\}$ for the kernel of D_b for b lying near b_v in $\mathcal{M}_T^{(0)}$, so that each $\xi_{b,i}$ varies smoothly with b . Then the claim follows immediately from an estimate on $S_\varpi R_{v(\varpi)} \xi_{b_\varpi, i} - R_v \xi_{b,i}$, since the projection maps can be expressed in terms of inner-products with $\xi_{b,i}$. Note that if we had defined $\Gamma_-(v)$ to be the kernel of D_v in the case $\mathcal{T}$ is regular, this claim, if true, would have been much harder to prove because of the presence of small eigenvalues of $D_v^* D_v$; see Subsection 3.6 for more details.

If $v \in F^{(0)} \mathcal{T}_\delta$, (Σ_v, g_v) can be viewed as a connected sum of the surfaces $\{(\Sigma_{\mathcal{T}, i}, g_{b_v, i})\}$ with very thin necks. If $\varpi \in \mathcal{K}_{b_v} \mathcal{T} \subset T_v F^\emptyset \mathcal{T}$ is as above and $|w_h| \geq 2|v_h|^{\frac{1}{2}}$, the maps $u_v : \Sigma \rightarrow V$ and $u_\varpi : \Sigma \rightarrow V$ are very far apart in the C^0 -norm even if $\|\varpi\|$ is small. However, we can still compare the two maps and the various objects of Lemma 3.6, appropriately defined, on the corresponding direct summands. If the gluing map of Subsection 3.6 is defined only on $F^{(0)} \mathcal{T}_\delta$, and not on $F \mathcal{T}_\delta$, we need to be able to do such comparisons in order to adjust the gluing map in the presence of constraints μ ; see Subsection 3.8.

In order to state an analogue of Lemma 3.6 with $\|\varpi\|_v$ for $\varpi \in \tilde{T}_\varpi F^H \mathcal{T}$ replaced by $\|\varpi\|$ for

$$\varpi \in \mathcal{K}_{b_v} \mathcal{T} \subset \tilde{T}_\varpi F^\emptyset \mathcal{T},$$

for each $\varpi \in \mathcal{K}_{b_v} \mathcal{T}_{\delta(b)}$, with δ sufficiently small, we construct a smooth map $\tilde{q}_\varpi : (\Sigma_v, g_v) \longrightarrow (\Sigma_\varpi, g_\varpi)$, which is almost an isometry. The map will depend only on the elements $w_h \in F_{b,h}^{(0)}$. The structure of the construction is similar to the construction of the map q_v in Subsection 2.2. For each $h \in \hat{I}$ with $\iota_h = \hat{0}$, let $\tilde{p}_{h,\varpi} : B_{b_v,h}(4\delta_{\mathcal{T}}(b_v)) \longrightarrow \Sigma$ be the (holomorphic) $(g_{b_v,\hat{0}}, g_{b_\varpi,\hat{0}})$ -isometry provided by Lemma 5.1. Define $\tilde{q}_{h,\varpi} : \Sigma \longrightarrow \Sigma$ by

$$\tilde{q}_{h,\varpi}(z) = \begin{cases} \phi_{b_v,h}^{-1} \left\{ \phi_{b_v,h} \tilde{p}_{h,\varpi}(z) + \beta_{\delta_{\mathcal{T}}^2(b_v)}(r_{b_v,h}(z)) (\phi_{b_v,h}(z) - \phi_{b_v,h} \tilde{p}_{h,\varpi}(z)) \right\}, & \text{if } r_{b_v,h}(z) \leq 2\delta_{\mathcal{T}}(b_v); \\ z, & \text{if } r_{b_v,h}(z) \geq 2\delta_{\mathcal{T}}(b_v). \end{cases}$$

If $h \in \hat{I}$ and $\iota_h \neq \hat{0}$, we similarly define $\tilde{q}_{h,(x_h,w_h)} : \Sigma_{b,\iota_h} \longrightarrow \Sigma_{b,\iota_h}$ by

$$\tilde{q}_{h,\varpi}(z) = \begin{cases} \phi_{b_v,h}^{-1} \left\{ \phi_{b_v,h}(z) + w_h - \beta_{\delta_{\mathcal{T}}^2(b_v)}(r_{b_v,h}(z)) w_h \right\}, & \text{if } r_{b_v,h}(z) \leq 2\delta_{\mathcal{T}}(b_v); \\ z, & \text{if } r_{b_v,h}(z) \geq 2\delta_{\mathcal{T}}(b_v). \end{cases}$$

Let $\tilde{q}_{\varpi,\hat{0}} = Id_\Sigma$. If $h \in \hat{I}$ and $\tilde{q}_{\varpi,\iota_h} : \Sigma \longrightarrow \Sigma$ has been constructed, let

$$\tilde{q}_{\varpi,h}(z) = \begin{cases} q_{\varpi,\iota_h}^{-1} (\tilde{p}_{h,\varpi}(z) (q_{\varpi,\iota_h}(\tilde{q}_{\varpi,\iota_h}(z)))) , & \text{if } r_{b_v,h}(q_{\varpi,\iota_h}(z)) \leq 2\delta_{\mathcal{T}}(b_v); \\ \tilde{q}_{\varpi,\iota_h}(z), & \text{if } r_{b_v,h}(q_{\varpi,\iota_h}(z)) \geq 2\delta_{\mathcal{T}}(b_v). \end{cases}$$

Going through all of I , we obtain a map $\tilde{q}_\varpi : \Sigma \longrightarrow \Sigma$, which shifts the connect-summands of (Σ, g_v) to the connect-summands of (Σ, g_ϖ) . The important properties of such maps $\tilde{q}_\varpi$ as summarized below.

Lemma 3.7 *There exist $\delta, C \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R})$ and a smooth family of maps*

$$\{\tilde{q}_\varpi : \Sigma \longrightarrow \Sigma \mid \varpi \in \mathcal{K}_{b_v} \mathcal{T}_{\delta(b_v)} \subset T_v F_v^{(\emptyset)} \mathcal{T}, v \in F_{b_v}^{(\emptyset)} \mathcal{T}_{\delta(b_v)}\}, \quad \text{such that}$$

(1) $\tilde{q}_0 = Id_\Sigma$ and $q_v = q_\varpi \circ \tilde{q}_\varpi \in \Sigma_{b_v,i}^* = \Sigma_{b_\varpi,i}^*$ outside of the annuli

$$A_\varpi = \tilde{q}_{\varpi,\iota_h}^{-1} q_{\varpi,\iota_h}^{-1} \left(\{z \in \Sigma_{b_v,\iota_h} : \delta_{\mathcal{T}}(b_v) \leq r_{b,h}(z) \leq 2\delta_{\mathcal{T}}(b_v)\} \right),$$

which contain no marked points of $b(v)$ or $b(\varpi)$.

(2) $\left| \frac{\tilde{q}_\varpi^* g_\varpi}{g_v} - \frac{\tilde{q}_{\varpi'}^* g_{\varpi'}}{g_v} \right| \leq C(b_v) \|\varpi - \varpi'\|$ for all $\varpi, \varpi' \in \mathcal{K}_{b_v} \mathcal{T}_{\delta(b_v)}$.

These maps q_ϖ allow us to compare operators on vector fields and one-forms on (Σ_v, u_v) and $(\Sigma_\varpi, u_\varpi)$ whenever $\|\varpi\|$ is sufficiently small. Define $\zeta'_\varpi \in \Gamma(u_v)$ by

$$\exp_{b_v, u_v} \zeta'_\varpi = u_\varpi \circ \tilde{q}_\varpi, \quad \|\zeta'_\varpi\|_{b_v, C^0} \leq \text{inj } g_{b_v}.$$

For $\xi \in \Gamma(u_v)$, let $R'_\varpi \xi \in \Gamma(u_\varpi)$ be given by

$$\{R'_\varpi \xi\}(z) = \Pi_{b_v, \zeta'_\varpi(\tilde{q}_\varpi^{-1}(z))} \xi(\tilde{q}_\varpi^{-1}(z)).$$

Similarly, of $\eta \in \Gamma^{0,1}(u_v)$, let $R'_\varpi \eta \in \Gamma^{0,1}(u_\varpi)$ be given by

$$\{R'_\varpi \eta\}|_z = \Pi_{b_v, \zeta'_\varpi(\tilde{q}_\varpi^{-1}(z))} \circ \eta|_{\tilde{q}_\varpi^{-1}(z)} \circ \partial \tilde{q}_\varpi^{-1}|_z.$$

Denote by S'_ϖ the inverse of R'_ϖ . Similarly to Lemma 3.6, we have

Lemma 3.8 *There exist $\delta, C \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R}^+)$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$ and $\varpi \in \mathcal{K}_{b_v}\mathcal{T} \subset \tilde{T}_v F^{(0)}\mathcal{T}_\delta$,*

- (1) $C(b_v)^{-1}\|\varpi\| \leq \|\zeta'_\varpi\|_{v,p,1} + \sum_{l \in M} |w_l(\varpi)|_{g_v} \leq C(b_v)\|\varpi\|$;
- (2) $\|S'_\varpi du_\varpi - du_v\|_{v,p} \leq C(b_v)\|\varpi\|$ and $\|S'_\varpi \bar{\partial} u_\varpi - \bar{\partial} u_v\|_{v,p} \leq C(b_v)|v|^{\frac{1}{p}}\|\varpi\|$;
- (3) $\|S'_\varpi \nu - \nu\|_{v,p} \leq C(b_v)\|\varpi\|$;
- (4) $\|S'_\varpi D_\varpi R'_\varpi \xi - D_v \xi\|_{v,p} \leq C(b_v)\|\varpi\|\|\xi\|_{v,p,1}$ and $\|S'_\varpi \pi_{\varpi, \pm} R'_\varpi \xi - \pi_{v, \pm} \xi\|_{v,p,1} \leq C(b_v)\|\varpi\|\|\xi\|_{v,p,1}$ for all $\xi \in \Gamma(u_v)$.

3.5 Obstruction Bundle Setup

In the next subsection, we look for solution of the equation $\bar{\partial} \exp_{b_v, u_v} \xi = t\nu$ with ξ lying in a fixed complement of $\Gamma_-(v)$. If t is sufficiently small, we are able to solve this equation up to an element of a vector bundle of the same rank as the dimension of $\Gamma_-(b_v)$, called obstruction bundle. This element is the obstruction to solving the equation. There are choices to be made for this obstruction bundle as well as for the subspace complementary to $\Gamma_-(v)$. We describe in this subsection what conditions these choices must satisfy for the gluing construction to work properly.

If $b^* = (S, M, I; x^*, (j, y^*), u^*) \in \mathcal{M}_T^{(0)}$ and $b = (S, M, I; x, (j, y), u_I) = H_{\mathcal{T}, b^*}(\sigma, \theta)$ for some $\sigma \in \mathcal{K}_{b^*}\mathcal{T}$ and $\theta \in \mathbb{R}^{\hat{I}}$, let $\xi_{b^*, b} = \xi_{b^*, b, I} \in \Gamma'(b)$ be given by

$$\exp_{b^*, u_i^*} \xi_{b^*, b, i} = u_i, \quad \|\xi_{b^*, b, i}\|_{C^0} < \text{inj } g_{V, b^*}.$$

Let $\Pi_{b^*, b} = \Pi_{b^*, \xi_{b^*, b}}$.

Definition 3.9 *Suppose $b^* = (S, M, I; x^*, (j, y^*), u^*)$, $b_k = (S, M, I; x_k, (j, y_k), u_k) \in \mathcal{M}_T^{(0)}$ and $v_k = (b_k, v_k) \in F^{(0)}\mathcal{T}$ are such that the sequences $\{b_k\}$ and $\{v_k|_{b_k}\}$ converge to $b^* \in \mathcal{M}_T^{(0)}$ and $0 \in \mathbb{R}$, respectively.*

- (1) *The sequence $\{\xi_k \in L_1^p(u_{v_k})\}$ C^0 -converges to $\xi^* \in \Gamma'(b^*)$ if*
 - (1a) *the sequence $\{\Pi_{b^*, b_k}^{-1}(\xi_k \circ q_{v_k}^{-1})\}$ C^0 -converges to ξ^* on compact subsets of $\Sigma_{b^*}^*$;*
 - (1b) *there exists $C > 0$ such that $\|\xi_k\|_{v_k, p, 1} < C$ for all k .*
- (2) *The sequence of subspaces $\{V_k \subset \Gamma(u_{v_k})\}$ C^0 -converges to subspace $V^* \subset \Gamma(b^*)$ if there exists a sequence of bases $\{\{\xi_{k,i}\}_{i=1}^{i=N} \subset V_k\}$ such that*
 - (2a) *for each i fixed, the sequence $\{\xi_{k,i}\}$ C^0 -converges to some $\xi_i^* \in V^*$;*
 - (2b) *the set $\{\xi_i^*\}$ has cardinality N and is basis for V^* .*

Lemma 3.10 *If the sequence $\{v_k\} \subset F^{(0)}\mathcal{T}$ converges to $b^* \in \mathcal{M}_T^{(0)}$ and the sequences $\{\xi_k \in L_1^p(v_k)\}$ and $\{\tilde{\xi}_k \in L_1^p(v_k)\}$ converge to $\xi^* \in \Gamma'(b^*)$ and $\tilde{\xi}^* \in \Gamma'(b^*)$, respectively,*

$$\lim_{k \rightarrow \infty} \langle \xi_k, \tilde{\xi}_k \rangle_{v_k, 2} = \langle \xi^*, \tilde{\xi}^* \rangle_{v^*, 2}.$$

Proof: If $v_k \rightarrow b^*$, the metrics $g_{V, b_{v_k}}$ on V and $g_{b_{v_k}, i}$ on $\Sigma_{\mathcal{T}, i}$ C^0 -converge to g_{V, b^*} and $g_{b^*, i}$, respectively. On the other hand, by (1b) of Definition 3.9 and (2) of Lemma 3.5, there exists $C > 0$ such that

$$\|\xi_k\|_{v_k, C^0}, \|\tilde{\xi}_k\|_{v_k, C^0} < C \quad \forall k.$$

Thus, the claim follows from (1a) of Definition 3.9.

Definition 3.11 Suppose Ω is an open subset of $F^{(0)}\mathcal{T}$ such that $b(v)$ is defined for all $v \in \Omega$. A $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant smooth complex subbundle $\tilde{\Gamma}_- \rightarrow \Omega$ of the Banach bundle $L_1^p \rightarrow \Omega$ is a tangent-space model over Ω if

- (1) for every sequence $\{v_k\} \subset \Omega$ converging to $b^* \in \mathcal{M}_{\mathcal{T}}^{(0)}$, a subsequence of $\{\tilde{\Gamma}_-(v_k)\}$ C^0 -converges to a subspace $V^* \subset \Gamma(b)$ such that $\pi_{b,-}: V^* \rightarrow \Gamma_-(b^*)$ is an isomorphism;
- (2) if $\tilde{\pi}_{v,-}: L_1^p(v) \rightarrow \tilde{\Gamma}_-(v)$ is the (L^2, v) -orthogonal projection, there exist $\delta, C \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for all $v \in \Omega_\delta$ and all $\xi \in \Gamma(u_v)$,
 - (2a) $\|S_\varpi \tilde{\pi}_{\varpi,-} R_\varpi \xi - \tilde{\pi}_{v,-} \xi\|_{v,2} \leq C(b_v) \|\varpi\|_v \|\xi\|_{v,p,1}$ for all $\varpi \in T_v F^{(0)} \mathcal{T}_{\delta(b_v)}$;
 - (2b) $\|S'_\varpi \tilde{\pi}_{\varpi,-} R'_\varpi \xi - \tilde{\pi}_{v,-} \xi\|_{v,2} \leq C(b_v) \|\varpi\| \|\xi\|_{v,p,1}$ for all $\varpi \in \mathcal{K}_{b_v} \mathcal{T} \subset T_v F^{(0)} \mathcal{T}_{\delta(b_v)}$.

One example of a tangent-space model is $\{\Gamma_-(v): v \in F^{(0)}\mathcal{T}_\delta\}$. In such a case, the limit V^* in (1) of Definition 3.11 is $\Gamma_-(b^*)$ and thus depends only on b^* , and not on the sequence $\{v_k\}$. However, for computational reasons, it is sometimes advantageous to work with other choices. With the choices in [Z2], the limit V^* in (1) of Definition 3.11 in fact depends on the sequence unless $|\hat{I}| = 1$. The following lemma collects some of the implications of (1) of Definition 3.11. Condition (2) is needed in Subsections 4.2 and 4.5. For any tangent space model over Ω and $v \in \Omega$, we denote the (L^2, v) -orthogonal complement of $\tilde{\Gamma}_-(v)$ by $\tilde{\Gamma}_+(v)$. Write $\tilde{\Gamma}_+^{0,1}(v)$ for the image of $\tilde{\Gamma}_+(v)$ under the operator $\tilde{D}_v$.

Lemma 3.12 Let $\tilde{\Gamma}_- \rightarrow \Omega$ be a tangent-space model. Then there exist $C, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R})$ such that for all $v \in \Omega_\delta$

- (1a) $\|\xi\|_{v,p,1} \leq C(b_v) \|\xi\|_{v,2}$ for all $\xi \in \tilde{\Gamma}_-(v)$;
- (1b) $\|\tilde{\pi}_{v,-} \xi\|_{v,p,1} \leq C(b_v) \|\xi\|_{v,p,1}$ for all $\xi \in \Gamma(u_v)$;
- (2a) $L_1^p(v) = \Gamma_-(v) \oplus \tilde{\Gamma}_+(v)$;
- (2b) if $\tilde{\pi}_-$ and $\tilde{\pi}_+$ are the projection maps corresponding to the above decomposition,

$$\|\tilde{\pi}_{v,\pm} \xi\|_{v,p,1} \leq C(b_v) \|\xi\|_{v,p,1} \quad \forall \xi \in \Gamma(u_v).$$

Proof: (1) Suppose there exists a sequence $\{v_k \in \Omega\}$ converging to $b^* \in \mathcal{M}_{\mathcal{T}}^{(0)}$ and a sequence $\{\xi_k \in \tilde{\Gamma}_-(v_k)\}$ such that $\|\xi_k\|_{v_k,p,1} = 1$, while $\|\xi_k\|_{v_k,2} \rightarrow 0$. Since $\|\xi_k\|_{v_k,p,1} = 1$, by (2) of Lemma 3.16 and (1) of Definition 3.11, a subsequence of $\{\xi_k\}$ C^0 -converges to some nonzero $\xi^* \in \Gamma(b^*)$. However, since $\|\xi_k\|_{v_k,2} \rightarrow 0$, $\|\xi^*\|_{b^*,2} = 0$ by Lemma 3.10. This is a contradiction, and thus (1a) holds. Claim (1b) is an immediate consequence of (1a) and (2) of Lemma 3.16.

(2) Claim (2a) is equivalent to saying that no nonzero element of $\tilde{\Gamma}_-(v)$ is orthogonal to $\Gamma_-(v)$. So, suppose $v_k \rightarrow b^* \in \mathcal{M}_{\mathcal{T}}^{(0)}$ and $\{\xi_k \in \tilde{\Gamma}_-(v_k)\}$ is such that ξ_k is orthogonal to $\Gamma_-(v)$ and $\|\xi_k\|_{v_k,p,1} = 1$. Since $\xi_k \in \tilde{\Gamma}_-(v_k)$ and $\|\xi_k\|_{v_k,p,1} = 1$, by (1) of Definition 3.11, a subsequence of $\{\xi_k\}$ converges to some nonzero $\xi^* \in \Gamma(b^*)$. By Lemma 3.10, ξ^* is orthogonal to $\Gamma_-(v)$. However, this contradicts the second part of (1) of Definition 3.11.

(3) Due to (1b), Claim (2b) is equivalent to saying that there exist $C, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}}; \mathbb{R})$ such that

$$\|\xi\|_{v,p,1} \leq C(b_v) \|\tilde{\pi}_{v,-} \xi\|_{v,p,1} \quad \forall v \in \Omega_\delta \text{ and } \xi \in \Gamma_-(v).$$

Suppose there exists a sequence $\{v_k\} \subset \Omega$ converging to some $b^* \in \mathcal{M}_{\mathcal{T}}^{(0)}$ and a sequence $\{\xi_k \in \Gamma_-(v_k)\}$ such that $\|\tilde{\pi}_{v_k,-} \xi_k\|_{v_k,2} \rightarrow 0$, while $\|\xi_k\|_{v_k,p,1} = 1$. By Definition 3.11, a subsequence of $\{\tilde{\Gamma}_-(v_k)\}$ converges to a subspace $V \subset \Gamma(b)$. On the other hand, a subsequence of $\{\xi_k\}$ C^0 -converges to a nonzero element $\xi^* \in \Gamma_-(b^*)$, which must be orthogonal to V by Lemma 3.10. This contradicts the second part of (1) of Definition 3.11.

Definition 3.13 Suppose Ω is an open subset of $F^{(0)}\mathcal{T}$ such that $b(v)$ is defined for all $v \in \Omega$. A $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant smooth complex subbundle $\Gamma_-^{0,1}(v) \longrightarrow \Omega$ of the Banach bundle $L^p \longrightarrow \Omega$ with the same rank as $\Gamma_-^{0,1} \longrightarrow \mathcal{M}_{\mathcal{T}}^{(0)}$ is an obstruction bundle if

(1) there exists $C \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R})$ such that

$$\|\eta\|_{v,p} \leq C(b_v)\|\eta\|_2 \quad \text{and} \quad \|D_v^* \eta\|_{v,1} \leq C(b_v)|v|^{\frac{1}{p}} \quad \forall v \in \Omega, \quad \eta \in \Gamma_-^{0,1}(v);$$

(2) if $\pi_{v,-}^{0,1} : L^p(v) \longrightarrow \Gamma_-^{0,1}(v)$ is the (L^2, v) -orthogonal projection, there exists $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for all $v \in \Omega_\delta$ and all $\eta \in \Gamma_-^{0,1}(u_v)$,

(2a) $\|S_\varpi \pi_{\varpi,-}^{0,1} R_\varpi \eta - \pi_{v,-}^{0,1} \eta\|_{v,2} \leq C(b_v)\|\varpi\|_v \|\xi\|_{v,p}$ for all $\varpi \in T_v F^{(0)}\mathcal{T}_{\delta(b_v)}$;

(2b) $\|S'_\varpi \pi_{\varpi,-}^{0,1} R'_\varpi \eta - \pi_{v,-}^{0,1} \eta\|_{v,2} \leq C(b_v)\|\varpi\| \|\xi\|_{v,p}$ for all $\varpi \in \mathcal{K}_{b_v} \mathcal{T} \subset T_v F^{(0)}\mathcal{T}_{\delta(b_v)}$.

Such an obstruction bundle is related to the cokernel bundle $\Gamma_-^{0,1} \longrightarrow \mathcal{M}_{\mathcal{T}}^{(0)}$. However, if $\hat{I} \neq \emptyset$, the low-eigenspaces of $D_v D_v^*$ are too large to form an obstruction bundle; see Remark below. Examples of bundles that satisfy Definition 3.13 can be found in [Z2]. Given such an obstruction bundle, we denote by $\pi_{v,-}^{0,1}$ and $\pi_{v,+}^{0,1}$ the (L^2, v) -orthogonal projection maps of $L^p(v)$ onto $\Gamma_-^{0,1}(v)$ and its (L^2, v) -orthogonal complement $\Gamma_+^{0,1}(v)$, respectively. The following lemma is clear from (1) of Definition 3.13.

Lemma 3.14 If $\Gamma_-^{0,1} \longrightarrow \Omega$ is an obstruction bundle, there exists $C \in C^\infty(\mathcal{M}_{\mathcal{T}}; \mathbb{R})$ such that

$$\|\pi_{v,\pm}^{0,1} \eta\|_{v,p} \leq C(b_v)\|\eta\|_{v,p} \quad \forall v \in \Omega, \quad \eta \in \Gamma_-^{0,1}(u_v).$$

Definition 3.15 If $\mathcal{T}$ is a semiregular bubble type, an obstruction bundle setup for $(V, J, \mathcal{T})$ is a tuple $(\delta, \tilde{\Gamma}_-, \Gamma_-^{0,1}, R)$, where

(1) $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ is $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant and $b(v)$ is defined for all $v \in F^{(0)}\mathcal{T}_\delta$;

(2) $\tilde{\Gamma}_- \longrightarrow F^{(0)}\mathcal{T}_\delta$ and $\Gamma_-^{0,1} \longrightarrow F^{(0)}\mathcal{T}_\delta$ are a tangent-bundle model and an obstruction bundle, respectively;

(3) $R : \pi^* \Gamma_-^{0,1} \longrightarrow \Gamma_-^{0,1}$ is a smooth oriented $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -equivariant bundle isomorphism over $F^{(0)}\mathcal{T}_\delta$, where $\pi : F^{(0)}\mathcal{T}_\delta \longrightarrow \mathcal{M}_{\mathcal{T}}^{(0)}$ is the bundle projection map.

For the rest of the paper, we fix such an obstruction bundle setup. However, whenever we refer to $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}}; \mathbb{R}^+)$, we will mean any function smaller than the function δ in Definition 3.15. The following lemma states some of the consequences of our setup that are crucial for the construction of the next subsection. If $\mathcal{T}$ is a regular bubble type, we take $\tilde{\Gamma}_-(v)$ and $\Gamma_-^{0,1}(v)$ to be $\Gamma_-(v)$ and $\{0\}$, respectively, and define the other bundles and the projection maps in the same way.

Lemma 3.16 If $\mathcal{T}$ is a simple bubble type, there exist $\delta, C \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for any $v \in F^{(0)}\mathcal{T}_\delta$ if $\mathcal{T}$ is regular and any $v \in F^{(0)}\mathcal{T}_\delta$ if $\mathcal{T}$ is semiregular,

(1) $\|\xi\|_{v,p,1} \leq C(b_v)\|D_v \xi\|_{v,p}$ for all $\xi \in \Gamma_+(v)$ and all $\xi \in \tilde{\Gamma}_+(v)$;

(2) $\|\pi_{v,-}^{0,1} \eta\|_{v,p} \leq C(b_v)|v|^{\frac{1}{p}}\|\eta\|_{v,p}$ for all $\eta \in \tilde{\Gamma}_+^{0,1}(v)$;

(3) $\pi_{v,+}^{0,1} : \tilde{\Gamma}_+^{0,1}(v) \longrightarrow \Gamma_+^{0,1}(v)$ is an isomorphism with the norm of the inverse bounded by $C(b_v)$.

Proof: (1) The first statement of the lemma is proved in the Appendix; see Proposition 5.13. It is consequence of (2) and (4) of Lemma 3.5 and of (1) of Definition 3.11. The second claim is immediate from (1) of Definition 3.13 and the first claim.

(2) Let W be the (L^2, g_v) -orthogonal complement of $\pi_{v,+}^{0,1}(\tilde{\Gamma}_+^{0,1}(v))$ in $\Gamma_+^{0,1}(v)$. The second claim implies that

$$L^p(v) = (\Gamma_-^{0,1}(v) \oplus W) \oplus \tilde{\Gamma}_+^{0,1}(v). \quad (3.5)$$

Since $\tilde{\Gamma}_+^{0,1}(v)$ is the image of $\tilde{\Gamma}_+(v)$ under D_v , with respect to the decompositions (3.5) and $L_1^p(v) = \Gamma_-(v) \oplus \tilde{\Gamma}_+(v)$,

$$D_v = \begin{vmatrix} D_v^{(--)} & 0 \\ D_v^{(+-)} & D_v^{(++)} \end{vmatrix}.$$

Since $D_v^{(++)}$ is an isomorphism by (1) of the lemma,

$$\begin{aligned} \text{ind } D_v &= \text{ind } D_v^{(--)} = \dim \Gamma_-(v) - (\dim \Gamma_-^{0,1}(v) + \dim W) \\ &= (\dim \Gamma_-(b_v) - (\dim \Gamma_-^{0,1}(b_v))) + \dim W = \text{ind } D_{b_v} + \dim W. \end{aligned} \quad (3.6)$$

On the other hand, by the Index Theorem, with $n = \dim_{\mathbb{C}} V$,

$$\begin{aligned} \text{ind } D_v &= 2 \left(\sum_{h \in \hat{I}(v)} (\langle c_1(V, J), \lambda_i(v) \rangle - n(g(\Sigma_{\mathcal{T}, i}) - 1)) - n(|\hat{I}(v)| - 1) \right) \\ &= 2 \left(\sum_{h \in \hat{I}(v)} \langle c_1(V, J), \lambda_i \rangle - n(g(S) - 1) \right) = \text{ind } D_{b_v}. \end{aligned} \quad (3.7)$$

By equations (3.6) and (3.7), $W = \{0\}$, and the last claim of the lemma follows from the second one.

Remark: It is essential for claim (1) of Lemma 3.16 that $p > 2$. The operator $D_v^* D_v$ has at least $|\hat{I}|(\dim V)$ eigenvalues that tend to 0 as $|v| \rightarrow 0$. The corresponding eigenfunctions converge to vector fields on the components of Σ_b that do not agree at the nodes. If $\mathcal{T}$ is semiregular, the operator D_{b_v} has cokernel $\Gamma_-^{0,1}(b)$. In such a case, the number of low eigenvalues of $D_v^* D_v$, including 0, is $(\dim \Gamma_-^{0,1}(b)) + |\hat{I}|(\dim V)$.

Let $\tilde{\pi}_{v,+}^{0,1} : \Gamma_+^{0,1}(v) \rightarrow \tilde{\Gamma}_+^{0,1}(v)$ denote the inverse of $\pi_{v,+}^{0,1} : \tilde{\Gamma}_+^{0,1}(v) \rightarrow \Gamma_+^{0,1}(v)$. We extend $\tilde{\pi}_{v,+}^{0,1}$ to all of $L^p(v)$ by taking it to be $\tilde{\pi}_{v,+}^{0,1} \circ \pi_{v,+}^{0,1}$. If $\eta \in \tilde{\Gamma}_+^{0,1}(v)$, let $P_v \eta \in \tilde{\Gamma}_+(v)$ be the unique element such that $D_v P_v \eta = \eta$. We extend P_v to all of $L^p(v)$ by taking it to be $P_v \circ \tilde{\pi}_{v,+}^{0,1}$. From Lemma 3.16, we immediately obtain

Corollary 3.17 *If $\mathcal{T}$ is a simple bubble type, there exist $\delta, C \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$ if $\mathcal{T}$ is regular and $v \in F^{(0)}\mathcal{T}_\delta$ if $\mathcal{T}$ is semiregular,*

- (1) $\|\tilde{\pi}_{v,+}^{0,1} \eta\|_{v,p} \leq C(b_v) \|\eta\|_{v,p}$ for all $\eta \in \Gamma_+^{0,1}(v)$;
- (2) $\|P_v \eta\|_{v,p,1} \leq C(b_v) \|\eta\|_{v,p}$ for all $\eta \in \Gamma_+^{0,1}(v)$.

3.6 The Gluing Map

In this subsection, we look for small vector fields $\xi \in \tilde{\Gamma}_+(v)$ such that $\exp_{b_v, u_v} \xi$ is holomorphic if $\mathcal{T}$ is regular and lies in $\mathcal{M}_{\Sigma, tv, \lambda^*}$ if $\mathcal{T}$ is semiregular. In Subsection 4.5, we show that all holomorphic maps if $\mathcal{T}$ is regular and all maps in $\mathcal{M}_{\Sigma, tv, \lambda^*} \times \Sigma^L$ if $\mathcal{T}$ is semiregular that lie near $\mathcal{M}_{\mathcal{T}}$ with respect to the Gromov topology can be obtained in this way.

If $\xi \in \Gamma(u_v)$, define $\exp_v \xi: \Sigma_v \longrightarrow V$ and $\bar{\partial}_v \xi \in \Gamma^{0,1}(u_v)$ by

$$\{\exp_v \xi\}(z) = \exp_{b_v, u_v(z)} \xi(z), \quad \{\bar{\partial}_v \xi\}|_z = \Pi_{b_v, \xi(z)}^{-1} \circ \bar{\partial}\{\exp_v \xi\}|_z.$$

If $S = \Sigma$ and $\nu \in \Gamma(\Sigma; \Lambda^{0,1} \pi_{\Sigma}^* T^* \Sigma \otimes \pi_V^* TV)$, let $\nu_{v, \xi} \in \Gamma^{0,1}(u_v)$ be given by

$$\nu_{v, \xi}|_z = \Pi_{b_v, \xi(z)}^{-1} \circ \nu|_{(z, \{\exp_v \xi\}z)}.$$

Then,

$$\bar{\partial}\{\exp_v \xi\}(\cdot) = t\nu|_{(\cdot, \{\exp_v \xi\}(\cdot))} \iff \bar{\partial}_v \xi = t\nu_{v, \xi}. \quad (3.8)$$

Write

$$\bar{\partial}_v \xi = \bar{\partial} u_v + D_v \xi + N_v \xi \quad \text{and} \quad \nu_{v, \xi}|_z = \nu|_{(z, u_v(z))} + L_{v, v} \xi|_z. \quad (3.9)$$

Then the second equation in (3.8) is equivalent to

$$D_v \xi + N_{v, tv} \xi = t\nu - \bar{\partial} u_v, \quad (3.10)$$

and by Proposition ?? in [Z1] and (1) of Lemma 3.5, there exist $C_{\bar{\partial}}, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for any $v \in F^{(0)} \mathcal{T}_{\delta}$ and $\xi_1, \xi_2 \in \Gamma(u_v)$,

$$\|N_{v, tv} \xi_1 - N_{v, tv} \xi_2\|_{v, p} \leq C_{\bar{\partial}}(b_v) (\|\xi_1\|_{v, p, 1} + \|\xi_2\|_{v, p, 1} + t) \|\xi_1 - \xi_2\|_{v, p, 1}. \quad (3.11)$$

If $\mathcal{T}$ is semiregular, the term ν will be fixed, and we will be looking for solutions of (3.10) with $t > 0$ very small for $v \in F^{(\emptyset)} \mathcal{T}_{\delta}$. If $\mathcal{T}$ is regular, we will consider (3.10) with $t = 0$ and $v \in F^{(0)} \mathcal{T}_{\delta}$. In both cases, we will consider only solutions ξ of (3.10) that lie in the subspace $\tilde{\Gamma}_+(v)$ of $L_1^p(v)$, since the subspace $\Gamma_-(v)$ corresponds to moving along the image of the pregluing map $v \longrightarrow b(v)$.

Vector field $\xi = P_v \eta$ with $\eta \in \Gamma_+^{0,1}(v)$ solves equation (3.10) if and only if

$$\eta + \pi_{v, +}^{0,1} N_{v, tv} P_v \eta = \pi_{v, +}^{0,1} (t\nu - \bar{\partial} u_v) \quad (3.12)$$

$$\text{and} \quad \pi_{v, -}^{0,1} \left(t\nu - \bar{\partial} u_v - \tilde{\pi}_{v, +}^{0,1} \eta - N_{v, tv} P_v \eta \right) = 0. \quad (3.13)$$

Denote the map $\eta \longrightarrow \pi_{v, +}^{0,1} N_{v, tv} P_v \eta$ by $N_{v, tv}^+$. By Corollary 3.17 and equation (3.11), there exist $\tilde{C}_{\bar{\partial}}, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for any $v \in F^{(0)} \mathcal{T}_{\delta}$ if $\mathcal{T}$ is regular and $v \in F^{(\emptyset)} \mathcal{T}_{\delta}$ if $\mathcal{T}$ is semiregular,

$$\|N_{v, tv}^+ \eta_1 - N_{v, tv}^+ \eta_2\|_{v, p} \leq \tilde{C}_{\bar{\partial}}(b_v) (\|\eta_1\|_{v, p} + \|\eta_2\|_{v, p} + t) \|\eta_1 - \eta_2\|_{v, p} \quad \forall \eta_1, \eta_2 \in \Gamma_+^{0,1}(v). \quad (3.14)$$

Lemma 3.18 *There exist $\epsilon, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$ and $t=0$ if $\mathcal{T}$ is regular, $v \in F^{(0)}\mathcal{T}_\delta$ and $t \in [0; \delta(b_v)]$ if $\mathcal{T}$ is semiregular, and $\alpha \in \Gamma_+^{0,1}(v)$ with $\|\alpha\|_{v,p} < \epsilon(b_v)$, the equation*

$$\eta + N_{v,t\nu}^+ \eta = \alpha$$

has a unique solution η_α in $\Gamma_+^{0,1}(v)$ such that $\|\eta_\alpha\|_{v,p} \leq 2\epsilon(b_v)$. Furthermore, such a solution satisfies $\|\eta_\alpha\|_{v,p} \leq 2\|\alpha\|_{v,p}$.

Proof: Put $\epsilon(b) = (6\tilde{C}_{\bar{\delta}}(b))^{-1}$, where $\tilde{C}_{\bar{\delta}}$ is as in (3.14). Define

$$\Psi_\alpha: \{\eta \in \Gamma_+^{0,1}(v) : \|\eta\|_{v,p} \leq 2\|\alpha\|_{v,p}\} \longrightarrow \Gamma_+^{0,1}(v)$$

by $\Psi_\alpha(\eta) = \alpha - N_{v,t\nu}^+ \eta$. By equation (3.14),

$$\begin{aligned} \|\Psi_\alpha(\eta)\|_{v,p} &\leq \|\alpha\|_{v,p} + \tilde{C}_{\bar{\delta}}(b_v)(\|\eta\|_{v,p} + t)\|\eta\|_{v,p} \leq 2\|\alpha\|_{v,p}; \\ \|\Psi_\alpha(\eta_1) - \Psi_\alpha(\eta_2)\|_{v,p} &\leq \tilde{C}_{\bar{\delta}}(b_v)(\|\eta_1\|_{v,p} + \|\eta_2\|_{v,p} + t)\|\eta_1 - \eta_2\|_{v,p} \leq \frac{5}{6}\|\eta_1 - \eta_2\|_{v,p}. \end{aligned}$$

It follows that Ψ_α is a contracting operator, and thus has a unique fixed point η_α , i.e.

$$\eta_\alpha + N_{v,t\nu}^+ \eta_\alpha = \alpha, \quad \text{and} \quad \|\eta_\alpha\|_{v,p} \leq 2\|\alpha\|_{v,p}.$$

The uniqueness claim follows immediately by taking the difference of the corresponding equations.

Corollary 3.19 *If $\mathcal{T}$ is a simple bubble type, there exist $\delta, \epsilon, C \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$ and $t=0$ if $\mathcal{T}$ is regular and $v \in F^{(0)}\mathcal{T}_\delta$ and $t \in [0; \delta(b_v)]$ if $\mathcal{T}$ is semiregular, there exists a unique $\eta_{v,t\nu} \in \Gamma^{0,1}(v)$ such that $\eta_{v,t\nu}$ satisfies equation (3.12) and $\|\eta_{v,t\nu}\|_{v,p} \leq \epsilon(b_v)$. Furthermore,*

$$\|\eta_{v,t\nu}\|_{v,p} \leq C(b_v)(t + |v|^{\frac{1}{p}}).$$

Proof: This corollary follows from Lemmas 3.18 and 3.5.

We now put $\xi_{v,t\nu} = P_v \eta_{v,t\nu}$ and $\tilde{u}_{v,t\nu} = \exp_v \xi_{v,t\nu}$. Replacing u_v in $b(v)$ by $\tilde{u}_{v,t\nu}$, we obtain a new bubble map that will be called $\tilde{b}_{t\nu}(v)$. If $\mathcal{T}$ is regular (and thus $t=0$), we will write $\tilde{u}_v$ and $\tilde{b}(v)$ for $\tilde{u}_{v,0}$ and $\tilde{b}_0(v)$, respectively. We can assume that the functions δ, ϵ and C of Corollary 3.19 are $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant. For $\mathcal{T}$ regular, we have thus constructed a *gluing* map

$$\tilde{\gamma}_{\mathcal{T}}^{(0)}: F^{(0)}\mathcal{T}_\delta \longrightarrow \bar{\mathcal{M}}_{\langle \mathcal{T} \rangle}, \quad v \longrightarrow \tilde{b}(v).$$

Since this map is $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant, as can be seen from the construction, $\tilde{\gamma}_{\mathcal{T}}^{(0)}$ induces a map on the quotient

$$\tilde{\gamma}_{\mathcal{T}}: F\mathcal{T}_\delta \longrightarrow \bar{\mathcal{M}}_{\langle \mathcal{T} \rangle}. \tag{3.15}$$

By the smooth dependence of solutions of (3.12), the restrictions

$$\tilde{\gamma}_{\mathcal{T}}^{(0)}: F^{(H)}\mathcal{T}_\delta \longrightarrow \mathcal{M}_{\mathcal{T}(H)}^{(0)}$$

are smooth. However, continuity of $\tilde{\gamma}_{\mathcal{T}}$ on all of $F\mathcal{T}_{\delta}$ is not immediate. In the next section, we show the map $\tilde{\gamma}_{\mathcal{T}}$ is a homeomorphism onto a neighborhood of $\mathcal{M}_{\mathcal{T}}$ in $\bar{\mathcal{M}}_{\langle \mathcal{T} \rangle}$. If $\mathcal{T}$ is semiregular and $t > 0$, we have constructed a map

$$\tilde{\gamma}_{\mathcal{T}, t\nu}^{(0)} : F^{(\emptyset)}\mathcal{T}_{\delta} \Big| \epsilon^{-1}(-t, t) \longrightarrow C_{(\lambda^*; M)}^{\infty}(\Sigma; V),$$

which again is $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant and thus descends to a map

$$\tilde{\gamma}_{\mathcal{T}, t\nu} : F^{\emptyset}\mathcal{T}_{\delta} \Big| (\epsilon^{-1}(-t, t)/G_{\mathcal{T}}) \longrightarrow C_{(\lambda^*; M)}^{\infty}(\Sigma; V). \quad (3.16)$$

The map $u_{\tilde{b}t\nu(v)}$ lies in $\mathcal{M}_{\Sigma, t\nu, \lambda^*}$ if and only if equation (3.13) is satisfied, i.e.

$$R_v \psi_{\mathcal{T}, t\nu}(v) \equiv t\nu - \bar{\partial}u_v - \tilde{\pi}_{v,+}^{0,1} \eta_{v, t\nu} - N_{v, t\nu} P_v \eta_{v, t\nu} = 0 \in \Gamma_{-}^{0,1}(v), \quad (3.17)$$

since $\eta_{v, t\nu}$ satisfies equation (3.12).

3.7 An Implicit Function Theorem

In this subsection, we prove a refined version of the Implicit Function Theorem. It will be used in the rest of this section to modify the gluing maps of Subsection 3.6 for the spaces $\mathcal{M}_{\mathcal{T}}(\mu)$, $\mathcal{U}_{\mathcal{T}}(\mu)$, etc.

Let $\mathcal{S}$ be a smooth oriented manifold, and $\mathcal{NS}$, $\mathcal{N}^{\mu}$, and F oriented Riemannian vector bundles over $\mathcal{S}$. We denote by b , $(b, \vec{n})$, (b, σ) , and (b, v) general elements of $\mathcal{S}$, $\mathcal{NS}$, $\mathcal{N}^{\mu}$, and F , respectively. If Ω is any subset of F and $\delta > 0$, let

$$\Omega(\delta) = \{(b, \vec{n}, v) \in \mathcal{NS} \oplus F : (b, v) \in \Omega; \ |\vec{n}|, |v| < \delta\}.$$

Let U be an open neighborhood of $\mathcal{S}$ in $\mathcal{NS} \oplus \mathcal{N}^{\mu} \oplus F$ and $h : U \longrightarrow \mathbb{R}^k$ a smooth map such that

$$h(b, \vec{n}, \sigma, v) = h(b, \vec{n}, \sigma, 0), \quad h|_{\mathcal{S}} = 0, \quad \text{and} \quad d(h : \mathcal{N}_b^{\mu} \longrightarrow \mathbb{R}^n)_{(b, 0)} : \mathcal{N}_b^{\mu} \longrightarrow \mathbb{R}^k$$

is an orientation-preserving isomorphism for all $b \in \mathcal{S}$. Let $\tilde{U}$ be a subset of U such that $\tilde{U}$ is the fiber product along $\mathcal{S}$ of an open neighborhood of $\mathcal{S}$ in $\mathcal{NS} \oplus \mathcal{N}^{\mu}$ and an open subset Ω of F . Suppose $\delta_{\mathcal{S}} > 0$, $C \in C^{\infty}(\mathcal{S}; \mathbb{R}^+)$, and $\tilde{h}_t : \tilde{U} \longrightarrow \mathbb{R}^k$ is a family of smooth functions with $t \in [0, \delta_{\mathcal{S}}]$ such that

$$|\tilde{h}_t - h|_{(b, \vec{n}, \sigma, v)}, \left| \frac{\partial \tilde{h}_t}{\partial \sigma} - \frac{\partial h}{\partial \sigma} \right|_{(b, \vec{n}, \sigma, v)} \leq C(b)(|v|^{\frac{1}{p}} + t) \quad \forall t \in (0, \delta_{\mathcal{S}}), \ (b, \vec{n}, \sigma, v) \in \tilde{U},$$

where $\frac{\partial h}{\partial \sigma}$ denotes the differential of h along the fibers of $\mathcal{N}^{\mu}$.

Lemma 3.20 *Let B be an open ball about $0 \in \mathbb{R}^k$. If $f : B \longrightarrow \mathbb{R}^k$ is a smooth function and*

$$k|Df|_z - Df|_0| < |(Df|_0)^{-1}|^{-1} \quad \forall z \in B,$$

then f is injective on B .

Proof: Let f_i denote the i th component of f . By the Mean Value Theorem, for all $x, y \in B$, there exists $z_i(x, y) \in B$ such that

$$|f_i(x) - f_i(y)| = |Df_i|_{z_i(x, y)}|x - y|.$$

Adding up these equations over all i , we obtain

$$\begin{aligned} \sum_{i=1}^{i=k} |f_i(x) - f_i(y)| &\geq \sum_{i=1}^{i=k} |Df_i|_0 |x - y| - k \sup_{z \in B} |Df|_z - Df|_0 |x - y| \\ &\geq \left(|(Df|_0)^{-1}|^{-1} - k \sup_{z \in B} |Df|_z - Df|_0| \right) |x - y|. \end{aligned}$$

Lemma 3.21 *For every precompact subset K of $\mathcal{S}$, there exists $\epsilon > 0$ such that for all $t \in (0, \epsilon)$ and $(b, \vec{n}, v) \in \Omega(\epsilon)|K$, the map*

$$\{(b, \sigma) \in \mathcal{N}^\mu : |\sigma| < \epsilon\} \longrightarrow \tilde{h}_t(b, \vec{n}, \sigma, v)$$

is defined and injective, and its differential defines an orientation-preserving isomorphism between $\mathcal{N}_b^\mu$ and $\mathbb{R}^k$.

Proof: The map above is defined as long as

$$\{(b, \vec{n}, \sigma, v) \in \mathcal{NS} \oplus \mathcal{N}_b^\mu \oplus F : b \in K, (b, \vec{n}, v) \in \Omega(\epsilon), |\sigma| < \epsilon\} \subset \tilde{U}.$$

Since K is precompact, existence of $\delta > 0$ such that the last inclusion holds is trivial. The other two statements follow from the third property of h and the second property of $\tilde{h}_t$ (see above); Lemma 3.20 is needed to prove the injectivity. Note that the variation of $\frac{\partial \tilde{h}_t}{\partial \sigma}$ over K can be bounded from the variation $\frac{\partial h}{\partial \sigma}$ and the second property of $\tilde{h}_t$.

Lemma 3.22 *For every precompact subset K of $\mathcal{S}$ and $\epsilon > 0$ sufficiently small, there exists $\delta > 0$ such that for all $t \in (0, \delta)$ and $(b, \vec{n}, v) \in \Omega(\delta)|K$, the image of the map*

$$\{(b, \sigma) \in \mathcal{N}^\mu : |\sigma| < \epsilon\} \longrightarrow \tilde{h}_t(b, \vec{n}, \sigma, v)$$

contains $0 \in \mathbb{R}^k$.

Proof: We assume $\epsilon > 0$ does not exceed the number provided by Lemma 3.21. Then by precompactness of K and the proof of Lemma 3.21,

$$\varepsilon \equiv \min \left\{ |h(b, \vec{n}, \sigma, v)| : (b, \vec{n}, v) \in \Omega_\epsilon|K, (b, \sigma) \in \mathcal{N}^\mu, |\sigma| = \frac{1}{2}\epsilon \right\} > 0. \quad (3.18)$$

Since for each $(b, \vec{n}, v) \in \Omega(\epsilon)|K$, the image of the map

$$\{(b, \sigma) \in \mathcal{N}^\mu : |\sigma| < \epsilon\} \longrightarrow h(b, \vec{n}, \sigma, v)$$

contains a neighborhood of 0 in $\mathbb{R}^k$ and $\tilde{h}_t$ is continuous, the claim follows from the first property of $\tilde{h}_t$ along with equation (3.18).

Corollary 3.23 *For every precompact open subset K of $\mathcal{S}$, there exist $\delta, C > 0$ with the following property. For all $t \in (0, \delta)$, there exists a smooth section*

$$\varphi_t \in \Gamma(\Omega(\delta)|K; \pi^* \mathcal{N}^\mu),$$

where $\pi : \Omega(\delta)|K \longrightarrow K$ is the bundle projection map, such that

$$\Omega(\delta)|K \longrightarrow \tilde{h}_t^{-1}(0), \quad (b, \vec{n}, v) \longrightarrow (b, \vec{n}, \varphi_t(b, \vec{n}, v), v),$$

is an orientation-preserving diffeomorphism. Furthermore,

$$|\varphi_t(b, \vec{n}, v)| \leq C(|v|^{\frac{1}{p}} + t + |\vec{n}|) \quad \forall (b, \vec{n}, v) \in \Omega(\delta)|K.$$

Finally, if G is a group that acts on the space $\mathcal{S}$ and bundles $\mathcal{NS}$, $\mathcal{N}^\mu$, and F , and preserves h , $\tilde{h}_t$, Ω , and K , then φ_t is G -equivariant.

Proof: With ϵ as provided by Lemma 3.21, let $\delta > 0$ be as provided by Lemma 3.22. Then

$$F_t : \{(b, \vec{n}, \sigma, v) : (b, \vec{n}, v) \in \Omega(\delta)|K, |\sigma| < \epsilon\} \longrightarrow \Omega(\delta) \times \mathbb{R}^k, \quad F_t(b, \vec{n}, \sigma, v) = (b, \vec{n}, v, \tilde{h}_t(b, \vec{n}, v))$$

is a diffeomorphism onto an open subset W of the target space. The inverse of F_t must have the form

$$F_t^{-1}(b, \vec{n}, v, \tilde{\sigma}) = (b, \vec{n}, \phi_t(b, \vec{n}, v, \tilde{\sigma}), v)$$

for some smooth function ϕ_t . By Lemma 3.22, $(\Omega(\delta)|K) \times \{0\} \subset W$. Thus,

$$\varphi_t \in \Gamma(\Omega(\delta)|K; \pi^* \mathcal{N}^\mu), \quad \varphi_t(b, \vec{n}, v) = \phi_t(b, \vec{n}, v, 0),$$

is a well-defined section, and by definition of ϕ_t ,

$$\Omega(\delta)|K \longrightarrow \tilde{h}_t^{-1}(0), \quad (b, \vec{n}, v) \longrightarrow (b, \vec{n}, \varphi_t(b, \vec{n}, v), v),$$

is a diffeomorphism. It is orientation-preserving by Lemma 3.21. The estimate on φ_t follows from the three properties of h , the first property of $\tilde{h}_t$, and the proof of Lemma 3.20. The final statement of the lemma is clear, since our construction commutes with the G -action.

3.8 The Orientation of $\mathcal{M}_{\Sigma, tv, \lambda^*}(\mu)$ and the Gluing Map

At this point, our treatments of regular and semiregular cases diverge. In this subsection, we assume that $\mathcal{T} = (\Sigma, [N], I; j, \lambda)$ is a semiregular bubble type and μ is an N -tuple of constraints in general position as defined below. We recall how each element of $\mathcal{M}_{\Sigma, tv, \lambda^*}(\mu)$ is assigned a sign and then specialize to the elements $\tilde{b}_{tv}(v) \in \mathcal{M}_{\Sigma, tv, \lambda^*}(\mu)$. We conclude this subsection with Theorem 3.29 that describes the elements of $\mathcal{M}_{\Sigma, tv, \lambda^*}(\mu)$ lying near the space $\mathcal{M}_{\mathcal{T}}(\mu)$.

Definition 3.24 (1) *Section $\nu \in \Gamma^{0,1}(\Sigma \times V; \Lambda_{j,j}^{0,1} \pi_\Sigma^* T^* \Sigma \otimes \pi_V^* TV)$ is λ^* -regular if for all $t \in (0, 1)$ and $u \in \mathcal{M}_{\Sigma, tv, \lambda^*}$, the operator $D_{V,u} : \Gamma(u) \longrightarrow \Gamma^{0,1}(u)$ is surjective.*

(2) *If ν is λ^* -regular, tuple μ of oriented submanifolds of V is ν -regular if for all $t \in (0, 1)$,*

$$\bigoplus_{l \in [N]} T_{u_{\hat{0}}(y_l)} V = \text{Im } dev_{[N]}|_b + \bigoplus_{l \in [N]} T_{u_{\hat{0}}(y_l)} \mu_l \quad \forall b = (\Sigma, [N], \{\hat{0}\}; (\hat{0}, y), u_{\hat{0}}) \in \mathcal{M}_{\Sigma, tv, \lambda^*}(\mu),$$

$$\text{where } dev_{[N]}|_b : \ker D_{V, u_{\hat{0}}} \oplus \bigoplus_{l \in [N]} T_{y_l} \Sigma \longrightarrow \bigoplus_{l \in [N]} T_{u_{\hat{0}}(y_l)} V, \quad dev_l|_b(\xi, w_{[N]}) = \xi(y_l) + du_{\hat{0}}|_{y_l} w_l.$$

(3) If $\mathcal{T}$ is a (V, J) -semiregular bubble type, tuple μ of oriented submanifolds of V is $\mathcal{T}$ -regular if

$$\bigoplus_{l \in [N]} T_{u_{j_l}(y_l)} V = \text{Im } \text{dev}_{[N]}|_b + \bigoplus_{l \in [N]} T_{u_{j_l}(y_l)} \mu_l \quad \forall b = (\Sigma, [N], I; x, (j, y), u) \in \mathcal{H}_{\mathcal{T}}(\mu);$$

where $\text{dev}_{[N]}|_b: \mathcal{K}_b \mathcal{T} \longrightarrow \bigoplus_{l \in [N]} T_{u_{j_l}(y_l)} V, \quad \text{dev}_l|_b(\xi_I, w_{\hat{I}+[N]}) = \xi_{j_l}(y_l) + du_{j_l}|_{y_l} w_l.$

(4) If $\mathcal{T}$ is a (V, J) -semiregular bubble type, $\mathcal{S} \subset \mathcal{M}_{\mathcal{T}}$ is a smooth submanifold, and $\tilde{\mathcal{S}} \subset \mathcal{M}_{\mathcal{T}}^{(0)}$ is the preimage of $\mathcal{S}$ under the quotient projection map, tuple μ of oriented submanifolds of V is $\mathcal{S}$ -regular if

$$\bigoplus_{l \in [N]} T_{u_{j_l}(y_l)} V = \text{dev}_{[N]}|_b(\mathcal{K}_b \mathcal{T} \cap T_b \tilde{\mathcal{S}}) + \bigoplus_{l \in [N]} T_{u_{j_l}(y_l)} \mu_l \quad \forall b \in \tilde{\mathcal{S}}(\mu) \equiv \tilde{\mathcal{S}} \cap \mathcal{M}_{\mathcal{T}}^{(0)}(\mu).$$

Note that all four definitions above are independent of the choice of metrics on V . Throughout this subsection, we assume that ν is λ^* -regular, $\mathcal{T}$ is semiregular, and μ is ν - and $\mathcal{T}$ -regular.

The space $\mathcal{M}_{\Sigma, t\nu, \lambda^*}$ consists of the maps $u: \Sigma \longrightarrow V$ such that $\bar{\partial}u|_z = t\nu(z, u(z))$ for all $z \in \Sigma$. Thus, the tangent space at u can be described as

$$T_u \mathcal{M}_{\Sigma, t\nu, \lambda^*} = \{\xi \in \Gamma(\Sigma; u^* TV) : D_{V,u} \xi - tL_{\nu,u} \xi = 0\},$$

where $L_{\nu,u} \xi$ is defined by

$$\{L_{\nu,u} \xi\}(z) = \nabla_{\xi(z)}^V \nu|_{(z, u(z))}.$$

The operator $D_{V,u} - tL_{\nu,u}$ is independent of the choice of the connection along $\mathcal{M}_{\Sigma, t\nu, \lambda^*}$ and by assumption has no cokernel if $t \in (0, 1)$. An orientation on $\mathcal{M}_{\Sigma, t\nu, \lambda^*}$ is determined by an orientation of the bundle $\Lambda_{\mathbb{R}}^{\text{top}} T\mathcal{M}_{\Sigma, t\nu, \lambda^*}$ over $\mathcal{M}_{\Sigma, t\nu, \lambda^*}$, which is the determinant line bundle of the elliptic operator $D_{V,u} - tL_{\nu,u}$. Since $L_{\nu,u}$ has order zero, the operator $D_{V,u} - tL_{\nu,u}$ is homotopic through elliptic operators to the operator $D_{V,u}$. Thus, $\Lambda_{\mathbb{R}}^{\text{top}} T\mathcal{M}_{\Sigma, t\nu, \lambda^*}$ is homotopic to

$$\det(D_{V,u}) = (\Lambda_{\mathbb{R}}^{\text{top}}(\ker D_{V,u})) \otimes (\Lambda_{\mathbb{R}}^{\text{top}}(\text{coker } D_{V,u}));$$

see [LM]. Since $D_{V,u}$ commutes with J , $\ker D_{V,u}$ and $\text{coker } D_{V,u}$ are both complex vector spaces and thus have natural orientations, which induce an orientation on the determinant line bundle of $D_{V,u}$ and via a homotopy of operators on the determinant bundle of $D_{V,u} - tL_{\nu,u}$. It follows that $\mathcal{M}_{\Sigma, t\nu, \lambda^*} \times \Sigma^N$ is naturally oriented. If μ is a ν -regular tuple of submanifold of V of total codimension

$$\text{codim } \mu = \dim \mathcal{M}_{\Sigma, t\nu, \lambda^*} \times \Sigma^N = \text{ind } D_{V,u} + 2|N| = 2\left(\langle c_1(V, J), \lambda^* \rangle + (\dim V)(1 - g(\Sigma)) + |N|\right),$$

the differential of the map

$$\text{ev}_{[N]}: \mathcal{M}_{\Sigma, t\nu, \lambda^*} \times \Sigma^N \longrightarrow \prod_{l \in [N]} V \quad (\Sigma, [N], \{\hat{0}\}; (\hat{0}, y), u_{\hat{0}}) \longrightarrow (u_{\hat{0}}(y_l))_{l \in [N]},$$

i.e. $\text{dev}_{[N]}$ as defined in (2) of Definition 3.24, induces an isomorphism between $T\mathcal{M}_{\Sigma, t\nu, \lambda^*} \oplus T\Sigma^N$ and the normal bundle of μ in V^N at each point of $\mathcal{M}_{\Sigma, t\nu, \lambda^*}(\mu)$. Here we identify the N -tuple μ with the submanifold

$$\prod_{l \in [N]} \mu_l \subset \prod_{l \in [N]} V \equiv V^N.$$

Since the normal bundle of μ is oriented, the evaluation map also induces an orientation on $T\mathcal{M}_{\Sigma,t\nu,\lambda^*} \oplus T\Sigma^N$ along $\mathcal{M}_{\Sigma,t\nu,\lambda^*}(\mu)$. Each element $b \in \mathcal{M}_{\Sigma,t\nu,\lambda^*}(\mu)$ is assigned a plus sign or is positively oriented if the two orientations agree, and a minus sign otherwise.

For any $v \in F^{(\emptyset)}\mathcal{T}$ such that q_v is defined, let $L_{\nu,v}: \Gamma(u_v) \longrightarrow \Gamma^{0,1}(u_v)$ be given by

$$\{L_{\nu,v}\xi\}(z) = \nabla_{\xi(z)}^{b_v} \nu|_{(z,f(z))}.$$

Denote by $\Gamma_{t,+}^{0,1}(v)$ the image of $\Gamma_+(v)$ under the map $D_v - tL_{\nu,v}$.

Lemma 3.25 *For any compact subset K of $\mathcal{M}_{\mathcal{T}}^{(0)}$, there exist $\delta, C > 0$ such that for all $v \in F^{(\emptyset)}\mathcal{T}_\delta|K$ and $t \in (0, \delta)$,*

- (1) $\|\xi\|_{v,p,1} \leq C\|D_v\xi - tL_{\nu,v}\xi\|_{v,p}$ for all $\xi \in \Gamma_+(v)$;
- (2) $L^p(v) = \Gamma_{t,+}^{0,1}(v) \oplus \Gamma_-^{0,1}(v)$;
- (3) if $D_{v,t}^-$ and $L_{\nu,v,t}^-$ are the $(-, -)$ -components of D_v and $L_{\nu,v}$ with respect to the decompositions $L_1^p(v) = \Gamma_+(v) \oplus \Gamma_-(v)$ and $L^p(v) = \Gamma_{t,+}^{0,1}(v) \oplus \Gamma_-^{0,1}(v)$, then

$$\pi_{v,-}: \ker \{D_v - tL_{\nu,v}: L_1^p(v) \longrightarrow L^p(v)\} \longrightarrow \ker \{D_{v,t}^- - tL_{\nu,v,t}^-: \Gamma_-(v) \longrightarrow \Gamma_-^{0,1}(v)\}$$

is an orientation-preserving isomorphism, provided one of the two operators is surjective.

Proof: (1) The first claim is immediate from (1) of Lemma 3.16 and (2) of Lemma 3.5. The second is obtained by the same argument as in the proof of (3) of Lemma 3.16.

(2) By construction, $\pi_{v,-}$ is an isomorphism of the two kernels of the lemma. In particular, $D_v - tL_{\nu,v}$ is surjective if and only if $D_{v,t}^- - tL_{\nu,v,t}^-$ is. Define

$$\begin{aligned} \Phi_\tau: L_1^p(v) \oplus \Gamma_-^{0,1}(v) &\longrightarrow L^p(v) \quad \text{and} \quad \Psi_\tau: \Gamma_-(v) \oplus \Gamma_-^{0,1}(v) \longrightarrow \Gamma_-^{0,1}(v) \quad \text{by} \\ \Phi_\tau(\xi, \eta) &= D_v\xi + \tau tL_{\nu,v}\xi + \eta \quad \text{and} \quad \Psi_\tau(\xi, \eta) = \tau(D_{v,t}^- + tL_{\nu,v,t}^-)\xi + \eta. \end{aligned}$$

The first map is surjective for all $\tau \in [0, 1]$ by (2) of the lemma, while the surjectivity of the second map is immediate from the definition. Furthermore, the maps

$$\phi_\tau: \ker \Phi_\tau \longrightarrow \Gamma_-(v), \quad \phi_\tau(\xi, \eta) = \pi_{v,-}\xi, \quad \text{and} \quad \psi_\tau: \ker \Psi_\tau \longrightarrow \Gamma_-(v), \quad \psi_\tau(\xi, \eta) = \xi$$

are isomorphisms such that

$$\psi_1^{-1}\phi_1(\xi, 0) = \pi_{v,-}\xi; \quad \text{if } \phi_1(\xi, \eta) = \psi_1(\xi', \eta'), \quad \eta = \eta'; \quad \psi_0^{-1}\phi_0J = J\psi_0^{-1}\phi_0.$$

It follows that $\pi_{v,-}$ is an orientation-preserving map between the two kernels of the lemma.

If K is a precompact open subset of $\mathcal{M}_{\mathcal{T}}$ and $\delta > 0$ is such that $\tilde{b}_{t\nu}(v)$ is defined for all $v \in F^{(\emptyset)}\mathcal{T}_\delta|K$ and $t \in (0, \delta)$, let $\mathcal{M}(K, \delta)$ and $\tilde{\mathcal{M}}_{t\nu}(K, \delta)$ denote the images of $F^{(\emptyset)}\mathcal{T}_\delta|K$ under the maps $\gamma_{\mathcal{T}}$ and $\tilde{\gamma}_{\mathcal{T},t\nu}$, respectively. Both maps are continuous and injective; see Subsection 4.2. The smooth structure of $F\mathcal{T}$ induces smooth structures on $\mathcal{M}(K, \delta)$ and $\tilde{\mathcal{M}}_{t\nu}(K, \delta)$, with the tangent bundles described by

$$\begin{aligned} T_{b(v)}\mathcal{M}(K, \delta) &= \left\{ \zeta'_\varpi = \frac{d}{d\tau} \zeta_\tau \varpi \Big|_{\tau=0} : \varpi \in T_v F\mathcal{T} \right\} \oplus \bigoplus_{l \in [N]} T_{y_l(v)} \Sigma; \\ T_{\tilde{b}_{t\nu}(v)}\tilde{\mathcal{M}}_{t\nu}(K, \delta) &= \left\{ \tilde{\zeta}'_\varpi = \frac{d}{d\tau} \tilde{\zeta}_\tau \varpi \Big|_{\tau=0} : \varpi \in T_v F\mathcal{T} \right\} \oplus \bigoplus_{l \in [N]} T_{y_l(v)} \Sigma, \end{aligned}$$

where $T_v F\mathcal{T}$ denotes $T_v F^\emptyset \mathcal{T}$; see Subsection 3.4. It is easy to see that $\varpi \longrightarrow \zeta'_\varpi$ is nearly complex linear and $\pi_{v,-}$ is almost the identity on the first component of $T_{b(v)} \mathcal{M}(K, \delta)$; both error terms are bounded by $C_K |v|$. Furthermore, by (1) of Lemma 3.6 and Corollary 4.7, $\varpi \longrightarrow \tilde{\zeta}'_\varpi$ also nearly computes with the complex structures and $\Pi_{b_v, \xi_{v, t\nu}} \pi_{v,-} \Pi_{b_v, \xi_{v, t\nu}}^{-1}$ is almost the identity on the first component of $T_{\tilde{b}_{t\nu}(v)} \tilde{\mathcal{M}}_{t\nu}(K, \delta)$; in the given case, the error terms are bounded by $C_K(t + |v|^{\frac{1}{p}})$. Thus, the orientations of $\mathcal{M}(K, \delta)$ and $\tilde{\mathcal{M}}_{t\nu}(K, \delta)$ induced by the natural orientation of $F\mathcal{T}$ agree with the orientations induced from the natural orientation on $\Gamma_-(v) \oplus \bigoplus_{l \in [N]} T_{y_l(v)} \Sigma$ via the maps $\pi_{v,-} \oplus id$ and $\pi_{v,-} \Pi_{b_v, \xi_{v, t\nu}}^{-1} \oplus id$, respectively.

By construction in Subsection 3.6,

$$\tilde{\psi}_{t\nu}: \tilde{\mathcal{M}}_{t\nu}(K, \delta) \longrightarrow \Gamma^{0,1}, \quad v \longrightarrow t\nu|_{\tilde{u}_{v, t\nu}} - \bar{\partial} \tilde{u}_{v, t\nu} \in \Gamma^{0,1}(\tilde{u}_{v, t\nu}),$$

determines a section of the bundle $\Pi \Gamma_-^{0,1}$ over $\tilde{\mathcal{M}}_{t\nu}(K, \delta)$, given by

$$\Pi \Gamma_-^{0,1}(\tilde{b}_{t\nu}(v)) = \Pi_{b_v, \xi_{v, t\nu}} \Gamma_-^{0,1}(v).$$

Note that the zero set of this section is precisely the space $(\mathcal{M}_{\Sigma, t\nu, \lambda} \times \Sigma^N) \cap \tilde{\mathcal{M}}_{t\nu}(K, \delta)$. A linearization of this section is given by

$$\begin{aligned} \nabla_{\tilde{\zeta}'_\varpi} (t\nu - \bar{\partial} \tilde{u}_{v, t\nu}) &\equiv \Pi_{b_v, \xi_{v, t\nu}} \pi_{v, t-}^{0,1} \nabla_{\pi_{v, -} \Pi_{b_v, \xi_{v, t\nu}}^{-1} \tilde{\zeta}'_\varpi} \Pi_{b_v, \xi_{v, t\nu}}^{-1} (t\nu - \bar{\partial} \tilde{u}_{v, t\nu}) \\ &= -\Pi_{b_v, \xi_{v, t\nu}} (D_{v, t}^{--} - tL_{\nu, v, t}^{--}) \pi_{v, -} \Pi_{b_v, \xi_{v, t\nu}}^{-1} \tilde{\zeta}'_\varpi, \end{aligned}$$

where $\pi_{v, t-}^{0,1}: L^p(v) = \Gamma_{+, t}^{0,1}(v) \oplus \Gamma_-^{0,1}(v) \longrightarrow \Gamma_-^{0,1}(v)$ is the projection map.

Corollary 3.26 *For any compact subset K of $\mathcal{M}_T^{(0)}$, there exists $\delta > 0$ such that for all $t \in (0, \delta)$, the orientation of $(\mathcal{M}_{\Sigma, t\nu, \lambda} \times \Sigma^N) \cap \tilde{\mathcal{M}}_{t\nu}(K, \delta)$ as the zero set of the section $\tilde{\psi}_{t\nu}$ agrees with its natural orientation.*

Proof: Suppose $\tilde{b}_{t\nu}(v) \in (\mathcal{M}_{\Sigma, t\nu, \lambda^*} \times \Sigma^N) \cap \tilde{\mathcal{M}}_{t\nu}(K, \delta)$. Since we can use any connection in $\tilde{u}_{v, t\nu}^* TV$ to define the natural orientation on $T_{\tilde{u}_{v, t\nu}} \mathcal{M}_{\Sigma, t\nu, \lambda^*}$, we can write

$$\{D_{\tilde{u}_{v, t\nu}} - tL_{\nu, \tilde{u}_{v, t\nu}}\} \xi = \Pi_{b_v, \xi_{v, t\nu}} \{D_v - tL_{\nu, v}\} \Pi_{b_v, \xi_{v, t\nu}}^{-1} \xi \quad \forall \xi \in \Gamma(\tilde{u}_{v, t}).$$

Thus, by Lemma 3.25, $\pi_{v, -} \circ \Pi_{b_v, \xi_{v, t\nu}}^{-1} \oplus id$ induces an orientation-preserving isomorphism between $T_{\tilde{b}_{t\nu}(v)} \mathcal{M}_{\Sigma, t\nu, \lambda^*} \oplus \bigoplus_{l \in [N]} T_{y_l(v)} \Sigma$ and $\ker(D_{v, t}^{--} - L_{\nu, v, t}^{--}) \oplus \bigoplus_{l \in [N]} T_{y_l(v)} \Sigma$ with their natural orientations.

By the preceding paragraph, the same is true for the zero set of $\tilde{\psi}_{t\nu}$.

If μ is an N -tuple of constraints as above, let

$$\begin{aligned} \mathcal{M}(K, \delta; \mu) &= \{b(v) \in \mathcal{M}(K, \delta): \text{ev}_{[N]}(b(v)) \in \mu\}, \\ \tilde{\mathcal{M}}_{t\nu}(K, \delta; \mu) &= \{\tilde{b}_{t\nu}(v) \in \tilde{\mathcal{M}}_{t\nu}(K, \delta): \text{ev}_{[N]}(\tilde{b}_{t\nu}(v)) \in \mu\}. \end{aligned}$$

Then $\mathcal{M}(K, \delta; \mu)$ and $\tilde{\mathcal{M}}_{t\nu}(K, \delta; \mu)$ are smooth manifolds. In fact, the smoothness of $\mathcal{M}(K, \delta; \mu)$ is immediate from the smoothness of $F\mathcal{T}|_{\mathcal{M}_T(\mu)}$, which is a consequence of $\mathcal{T}$ -regularity of μ , while

the smoothness of $\tilde{\mathcal{M}}_{t\nu}(K, \delta; \mu)$ follows from Lemma 3.28 below. Furthermore, since μ is ν -regular, the section $\tilde{\psi}_{t\nu}$ is transversal to zero in $\Pi_{-}^{0,1}$ over $\tilde{\mathcal{M}}_{t\nu}(K, \delta; \mu)$. By Corollary 3.26, the sign of $\tilde{b}_{t\nu}(v) \in \mathcal{M}_{\Sigma, t\nu, \lambda^*}(\mu)$ defined at the beginning of this subsection is its sign as an element of the zero set of the section $\tilde{\psi}_{t\nu}$ of $\Pi_{-}^{0,1}$ over $\tilde{\mathcal{M}}_{t\nu}(K, \delta; \mu)$.

If $b = (\Sigma, [N], I; x, (j, y), u) \in \mathcal{M}_{\mathcal{T}}^{(0)}(\mu)$, let

$$\mathcal{K}_b^\mu \mathcal{T} = \{(\xi, w_{\hat{I}+[N]}) \in \mathcal{K}_b \mathcal{T} : \xi_{j_l}(y_l) + du_{j_l}|_{y_l} w_l \in T_{u_{j_l}(y_l)} \mu_l \ \forall l \in [N]\}.$$

Denote by $\mathcal{N}_b^\mu \mathcal{T}$ the (L^2, b) -orthogonal complement of $\mathcal{K}_b^\mu \mathcal{T}$ in $\mathcal{K}_b \mathcal{T}$. Note that by (3) of Definition 3.24,

$$\bigoplus_{l \in [N]} T_{u_{j_l}(y_l)} V = \text{dev}_{[N]}|_b(\mathcal{N}_b^\mu \mathcal{T}) \oplus \bigoplus_{l \in [N]} T_{u_{j_l}(y_l)} \mu_l.$$

We denote by $\tilde{\mathcal{N}}^\mu \mathcal{T}$ the bundle over $\mathcal{M}_{\mathcal{T}}^{(0)}(\mu)$ with fibers $\mathcal{N}_b^\mu \mathcal{T}$ and by $\mathcal{N}^\mu \mathcal{T} \rightarrow \mathcal{M}_{\mathcal{T}}(\mu)$ its quotient by the natural $G_{\mathcal{T}}$ -action.

Suppose $\mathcal{S} \subset \mathcal{M}_{\mathcal{T}}$ is a smooth oriented submanifold such that μ is $\mathcal{S}$ -regular. Denote by $\tilde{\mathcal{S}} \subset \mathcal{M}_{\mathcal{T}}^{(0)}$ the preimage of $\mathcal{S}$ under the quotient projection map. Let $\mathcal{N}\mathcal{S} \rightarrow \mathcal{S}$ and $\mathcal{N}\tilde{\mathcal{S}} \rightarrow \tilde{\mathcal{S}}$ be the normal bundles. Choose a $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -equivariant orientation-preserving identification $\tilde{\phi}_{\mathcal{S}} : \mathcal{N}\tilde{\mathcal{S}} \rightarrow \mathcal{M}_{\mathcal{T}}^{(0)}$ of neighborhoods of $\tilde{\mathcal{S}}$ in $\mathcal{N}\tilde{\mathcal{S}}$ and $\mathcal{M}_{\mathcal{T}}^{(0)}$. Let $\tilde{\Phi}_{\mathcal{S}} : \pi_{\mathcal{N}\tilde{\mathcal{S}}}^* F^{(0)} \mathcal{T} \rightarrow F^{(0)} \mathcal{T}$ be a $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -equivariant vector-bundle isomorphism covering $\tilde{\phi}_{\mathcal{S}}$ such that $\tilde{\Phi}_{\mathcal{S}}$ is the identity on $\tilde{\mathcal{S}}$. Let

$$\phi_{\mathcal{S}} : \mathcal{N}\mathcal{S} \rightarrow \mathcal{M}_{\mathcal{T}} \quad \text{and} \quad \Phi_{\mathcal{S}} : \pi_{\mathcal{N}\mathcal{S}}^* F\mathcal{T} \rightarrow F\mathcal{T}$$

be the maps induced by $\tilde{\phi}_{\mathcal{S}}$ and $\tilde{\Phi}_{\mathcal{S}}$, respectively. Put

$$\mathcal{S}(\mu) = \mathcal{S} \cap \mathcal{M}_{\mathcal{T}}(\mu), \quad \tilde{\mathcal{S}}(\mu) = \tilde{\mathcal{S}} \cap \mathcal{M}_{\mathcal{T}}^{(0)}(\mu).$$

Since μ is $\mathcal{S}$ -regular, we can choose a $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -equivariant orientation-preserving identification $\tilde{\phi}_{\mathcal{S}}^\mu : \tilde{\mathcal{N}}^\mu \mathcal{T}_\delta | \tilde{\mathcal{S}}(\mu) \rightarrow \tilde{\mathcal{S}}$. Let

$$\tilde{\Phi}_{\mathcal{S}}^\mu : \pi_{\tilde{\mathcal{N}}^\mu \mathcal{T}}^* (\mathcal{N}\tilde{\mathcal{S}} \oplus F^{(0)} \mathcal{T}) \rightarrow \mathcal{N}\tilde{\mathcal{S}} \oplus F^{(0)} \mathcal{T}$$

be a $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -equivariant splitting-preserving vector-bundle isomorphism covering $\tilde{\phi}_{\mathcal{S}}^\mu$ such that $\tilde{\Phi}_{\mathcal{S}}^\mu$ is the identity on $\tilde{\mathcal{S}}(\mu)$. Denote by

$$\phi_{\mathcal{S}}^\mu : \mathcal{N}^\mu \mathcal{T}_\delta | \mathcal{S}(\mu) \rightarrow \tilde{\mathcal{S}} \quad \text{and} \quad \Phi_{\mathcal{S}}^\mu : \pi_{\mathcal{N}^\mu \mathcal{T}}^* (\mathcal{N}\mathcal{S} \oplus F\mathcal{T}) \rightarrow \mathcal{N}\mathcal{S} \oplus F\mathcal{T}$$

the maps induced by $\tilde{\phi}_{\mathcal{S}}^\mu$ and $\tilde{\Phi}_{\mathcal{S}}^\mu$, respectively.

Definition 3.27 *With notation as above and in Subsection 3.4, tuple $(\Phi_{\mathcal{S}}, \Phi_{\mathcal{S}}^\mu)$ is a regularization of $\mathcal{S}(\mu)$ if for all $b \in \tilde{\mathcal{S}}(\mu)$, $\vec{n} \in \mathcal{N}_b \tilde{\mathcal{S}}_{\delta(b)}$, and $\sigma \in \tilde{\mathcal{N}}_b^\mu \mathcal{T}_{\delta(b)}$, there exists $\varpi(\vec{n}, \sigma) \in \mathcal{K}_{\tilde{\phi}_{\mathcal{S}}(b, \vec{n})} \mathcal{T}$ such that*

$$\tilde{\Phi}_{\mathcal{S}} \tilde{\Phi}_{\mathcal{S}}^\mu(b, \sigma; \vec{n}, v_j) = \{\tilde{\Phi}_{\mathcal{S}}(b, \vec{n}; v)\}(\varpi(\vec{n}, \sigma)) \quad \forall v \in F_b^{(0)} \mathcal{T}.$$

Note that if μ is $\mathcal{S}$ -regular, $\mathcal{S}(\mu)$ admits a normalization. In fact, we can start with any choice of $\Phi_{\mathcal{S}}$ and $\Phi_{\mathcal{S}}^{\mu}|_{\pi_{\mathcal{N}^{\mu}\mathcal{T}}^*\mathcal{N}\mathcal{S}}$ as in the preceding paragraph, and then choose $\Phi_{\mathcal{S}}^{\mu}|_{\pi_{\mathcal{N}^{\mu}\mathcal{T}}^*F\mathcal{T}}$ so that the triple satisfies the requirements of the definition. In applications of Theorem 3.29 in [Z2], the exact choice of $\Phi_{\mathcal{S}}^{\mu}$ does not matter, but that of $\Phi_{\mathcal{S}}$ does play a role.

For the purposes of Theorem 3.29, we assume that $\Phi_{\mathcal{S}}$ and $\Phi_{\mathcal{S}}^{\mu}$ also encode the lifts of $\phi_{\mathcal{S}}$ and $\phi_{\mathcal{S}}^{\mu}$ to the bundles $\pi_{\mathcal{N}\mathcal{S}}^*\Gamma_{-}^{0,1} \rightarrow \mathcal{S}$ and $\pi_{\mathcal{N}^{\mu}\mathcal{T}}^*\Gamma_{-}^{0,1} \rightarrow \mathcal{S}(\mu)$, respectively. Put

$$\begin{aligned} F^{(0)}\mathcal{S} &= \mathcal{N}\tilde{\mathcal{S}} \oplus F^{(0)}\mathcal{T}, & F^{(0)}\mathcal{S} &= \{(b, \vec{n}, v_{\hat{I}}) \in F^{(0)}\mathcal{S} : v_{\hat{I}} \in F_b^{(0)}\mathcal{T}\}; \\ FS &= \mathcal{N}\mathcal{S} \oplus F\mathcal{T}, & F^{\emptyset}\mathcal{S} &= \{[b, \vec{n}, v_{\hat{I}}] \in F^{(0)}\mathcal{S} : [b, v_{\hat{I}}] \in F_b^{\emptyset}\mathcal{T}\}. \end{aligned}$$

Lemma 3.28 *For any $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -invariant precompact open subset K of $\tilde{\mathcal{S}}(\mu)$, there exist an open neighborhood U_K of K in $\mathcal{M}_{\mathcal{T}}^{(0)}$ and $\delta, C > 0$ with the following property. If $t \in (0, \delta)$, there exists a smooth $(G_{\mathcal{T}} \times \text{Aut}(\mathcal{T}))$ -equivariant section*

$$\tilde{\varphi}_{\mathcal{S}, t\nu}^{\mu} \in \Gamma(F^{(0)}\mathcal{S}_{\delta}|K; \pi_{F^{(0)}\mathcal{S}}^*\tilde{\mathcal{N}}^{\mu}\mathcal{T}),$$

such that $\|\tilde{\varphi}_{\mathcal{S}, t\nu}^{\mu}(v)\|_{b\nu, C^0} \leq C(t + |v|^{\frac{1}{p}})$ for all $v \in F^{(0)}\mathcal{S}_{\delta}|K$ and

$$F^{\emptyset}\mathcal{S}_{\delta}|K \rightarrow \tilde{\mathcal{M}}_{t\nu}(U_K, \delta; \mu), \quad [b, \vec{n}, v_{\hat{I}}] \rightarrow \tilde{\gamma}_{\mathcal{T}, t\nu}(\tilde{\Phi}_{\mathcal{S}}(\tilde{\Phi}_{\mathcal{S}}^{\mu}\tilde{\varphi}_{\mathcal{S}, t\nu}^{\mu}(b, \vec{n}, v_{\hat{I}}))),$$

is an orientation-preserving diffeomorphism.

Proof: Since μ is a regular value of $\text{ev}_{[N]}|\mathcal{S}$ and K is precompact, there exists $\delta > 0$ such that the map

$$\{(b, \vec{n}, v, \sigma) \in F^{(0)}\mathcal{S} \oplus \tilde{\mathcal{N}}^{\mu}\mathcal{T} | K : |\vec{n}|, \|\sigma\|_{b, C^0} < \delta\} \rightarrow F^{(0)}\mathcal{T}, \quad (b, \vec{n}, v, \sigma) \rightarrow \tilde{\Phi}_{\mathcal{S}}\tilde{\Phi}_{\mathcal{S}}^{\mu}(b, \sigma; \vec{n}, v),$$

is a $G_{\mathcal{T}}$ -equivariant orientation-preserving diffeomorphism onto its image. Thus, if $\delta > 0$ is sufficiently small, there exists $C > 0$ such that, with notation as in Definition 3.27,

$$C^{-1}\|\sigma - \sigma'\|_{b, C^0} \leq \|\varpi(\vec{n}, \sigma) - \varpi(\vec{n}, \sigma')\|_{b, C^0} \leq C\|\sigma - \sigma'\|_{b, C^0} \quad \forall b \in \tilde{\mathcal{S}}(\mu), \vec{n} \in \mathcal{N}_b\tilde{\mathcal{S}}_{\delta}, \sigma, \sigma' \in \tilde{\mathcal{N}}_b^{\mu}\mathcal{T}.$$

Then by Corollary 4.11 and definition of S'_{ϖ} in Subsection 3.4,

$$\begin{aligned} & \left| d_V(\tilde{\varphi}_{\mathcal{S}}\tilde{\Phi}_{\mathcal{S}}^{\mu}(b, \sigma; \vec{n}), \tilde{\gamma}_{\mathcal{T}, t\nu}(\tilde{\Phi}_{\mathcal{S}}\tilde{\Phi}_{\mathcal{S}}^{\mu}(b, \sigma; \vec{n}, v))) - d_V(\tilde{\varphi}_{\mathcal{S}}\tilde{\Phi}_{\mathcal{S}}^{\mu}(b, \sigma'; \vec{n}), \tilde{\gamma}_{\mathcal{T}, t\nu}(\tilde{\Phi}_{\mathcal{S}}\tilde{\Phi}_{\mathcal{S}}^{\mu}(b, \sigma'; \vec{n}, v))) \right| \\ & \leq C(t + |v|^{\frac{1}{p}})\|\sigma - \sigma'\|_{b, C^0} \quad \forall t \in (0, \delta), b \in \tilde{\mathcal{S}}(\mu), \vec{n} \in \mathcal{N}_b\tilde{\mathcal{S}}_{\delta}, \sigma, \sigma' \in \tilde{\mathcal{N}}_b^{\mu}\mathcal{T}, v \in F_b^{(0)}\mathcal{T}_{\delta}. \end{aligned}$$

On a neighborhood of $\text{ev}_{[N]}(b) \in \mu$, we can identify the normal bundle of μ in V^N g_V -isometrically with the trivial hermitian bundle of the same rank. Let π denote the projection onto the fiber. Since μ is $\mathcal{S}$ -regular,

$$\|\sigma - \sigma'\|_{b, C^0} \leq C|\pi \text{ev}_{[N]}(\tilde{\varphi}_{\mathcal{S}}\tilde{\Phi}_{\mathcal{S}}^{\mu}(b, \sigma; \vec{n})) - \pi \text{ev}_{[N]}(\tilde{\varphi}_{\mathcal{S}}\tilde{\Phi}_{\mathcal{S}}^{\mu}(b, \sigma'; \vec{n}))| \quad \forall b \in \tilde{\mathcal{S}}(\mu), \vec{n} \in \mathcal{N}_b\tilde{\mathcal{S}}_{\delta}, \sigma, \sigma' \in \tilde{\mathcal{N}}_b^{\mu}\mathcal{T}.$$

Thus, we can apply Corollary 3.23 to

$$h = \pi \circ \text{ev}_{[N]} \circ \tilde{\varphi}_{\mathcal{S}} \circ \tilde{\Phi}_{\mathcal{S}}^{\mu} \quad \text{and} \quad \tilde{h}_t = \pi \circ \text{ev}_{[N]} \circ \tilde{\gamma}_{\mathcal{T}, t\nu} \circ \tilde{\Phi}_{\mathcal{S}} \circ \tilde{\Phi}_{\mathcal{S}}^{\mu}.$$

We obtain $\delta, \epsilon > 0$ and for each $t \in (0, \delta)$ a section $\tilde{\varphi}_{\mathcal{S}, t\nu}^\mu$ with the claimed bound such that the map

$$\begin{aligned} F^{(0)}\mathcal{S}_\delta|K &\longrightarrow \left\{ (b, \vec{n}, v_I, \sigma) : \|\sigma\|_{b, C^0} < \epsilon, \text{ev}_{[N]}\tilde{\gamma}_{\mathcal{T}, t\nu}(\tilde{\Phi}_{\mathcal{S}}(\tilde{\Phi}_{\mathcal{S}}^\mu(b, \vec{n}; \sigma, v))) \in \mu \right\}, \\ (b, \vec{n}, v) &\longrightarrow (b, \vec{n}, v, \tilde{\varphi}_{\mathcal{S}, t\nu}^\mu(b, \vec{n}, v)) \end{aligned}$$

is an orientation-preserving diffeomorphism. Since

$$\begin{aligned} \{[b, \vec{n}, v, \sigma] : [b, \vec{n}, v] \in F^{(0)}\mathcal{S}_\delta, \|\sigma\|_{b, C^0} < \epsilon\} &\longrightarrow \tilde{\mathcal{M}}_{t\nu}(U_K, \delta), \\ [b, \vec{n}, \sigma, v] &\longrightarrow \tilde{\gamma}_{\mathcal{T}, t\nu}(\tilde{\Phi}_{\mathcal{S}}(\tilde{\Phi}_{\mathcal{S}}^\mu(b, \vec{n}; \sigma, v))), \end{aligned}$$

is orientation-preserving by the discussion above and our assumptions on $\tilde{\phi}_{\mathcal{S}}$, the claim follows. Above

$$U_K = \tilde{\phi}_{\mathcal{S}}\left(\tilde{\Phi}_{\mathcal{S}}^\mu(\{(b, \vec{n}, \sigma) \in \mathcal{N}\tilde{\mathcal{S}} \oplus \tilde{\mathcal{N}}^\mu\mathcal{T} | K : \|\vec{n}\|_{b, C^0} < \epsilon, \|\sigma\|_{b, C^0} < \epsilon\})\right).$$

Theorem 3.29 Suppose $\lambda^* \in H_2(V; \mathbb{Z})$, $\mathcal{T} = (\Sigma, [N], I; j, \lambda)$ is a (V, J) -semiregular bubble type, with $\sum_{i \in I} \lambda_i = \lambda^*$ and cokernel bundle $\Gamma_-^{0,1} \longrightarrow \mathcal{M}_{\mathcal{T}}$, and $(\tilde{\Gamma}_-, \Gamma_-^{0,1}, R)$ is an obstruction bundle setup. Let $\mathcal{S} \subset \mathcal{M}_{\mathcal{T}}$ be a smooth oriented submanifold,

$$\nu \in \Gamma^{0,1}(\Sigma \times V; \Lambda_{J,j}^{0,1} \pi_{\Sigma}^* T^* \Sigma \otimes \pi_V^* TV)$$

a λ^* -regular section, μ a ν -, $\mathcal{T}$ -, and $\mathcal{S}$ -regular N -tuple of submanifolds of V of total codimension

$$\text{codim } \mu = 2(\langle c_1(V, J), \lambda^* \rangle + (\dim V)(1 - g(S)) + |N|),$$

and $(\Phi_{\mathcal{S}}, \Phi_{\mathcal{S}}^\mu)$ is regularization of $\mathcal{S}(\mu)$. Then for every precompact open subset K of $\mathcal{S}(\mu)$, there exist a neighborhood U_K of K in $\bar{C}_{(\lambda^*, N)}^\infty(\Sigma; \mu)$ and $\delta, \epsilon, C > 0$ with the following property. For every $t \in (0, \epsilon)$, there exist a section

$$\varphi_{\mathcal{S}, t\nu}^\mu \in \Gamma(F^{(0)}\mathcal{S}_\delta|K; \pi_{F\mathcal{S}}^* \mathcal{N}^\mu \mathcal{T}), \quad \text{with} \quad \|\varphi_{\mathcal{S}, t\nu}^\mu(v)\|_{b_v, C^0} \leq C(t + |v|^{\frac{1}{p}}),$$

and a sign-preserving bijection between $\mathcal{M}_{\Sigma, t\nu, \lambda^*}(\mu) \cap U_K$ and the zero set of the section $\psi_{\mathcal{S}, t\nu}^\mu$ defined by

$$\begin{aligned} \psi_{\mathcal{S}, t\nu}^\mu &\in \Gamma(F^{(0)}\mathcal{S}_\delta|K; \pi_{F\mathcal{S}}^* \Gamma_-^{0,1}), \quad \Phi_{\mathcal{S}}^\mu(\varphi_{\mathcal{S}, t\nu}^\mu(v); \psi_{\mathcal{S}, t\nu}^\mu(v)) = \psi_{\mathcal{S}, t\nu}(\Phi_{\mathcal{S}}^\mu(\varphi_{\mathcal{S}, t\nu}^\mu(v))); \\ \psi_{\mathcal{S}, t\nu} &\in \Gamma(F^{(0)}\mathcal{S}_\delta|(\mathcal{S} \cap U_K); \pi_{F\mathcal{S}}^* \Gamma_-^{0,1}), \quad \Phi_{\mathcal{S}}(v; \psi_{\mathcal{S}, t\nu}(v)) = \psi_{\mathcal{T}, t\nu}(\Phi_{\mathcal{S}}(v)); \\ \psi_{\mathcal{T}, t\nu} &\in \Gamma(F^{(0)}\mathcal{T}_\delta|(\mathcal{M}_{\mathcal{T}} \cap U_K); \pi_{F\mathcal{T}}^* \Gamma_-^{0,1}), \quad R_v \psi_{\mathcal{T}, t\nu}(v) = \pi_{v, -}^{0,1}(t\nu_{v, t} - \bar{\partial}(u_{b_v} \circ q_v) - D_v \xi_{v, t\nu} - \tilde{\eta}_{v, t\nu}) \end{aligned}$$

for some $\nu_{v, t}, \tilde{\eta}_{v, t\nu} \in \Gamma^{0,1}(u_v)$ and $\xi_{v, t\nu} \in \tilde{\Gamma}_+(v)$, dependent smoothly on v , such that

$$\|\nu_{v, t} - \nu\|_{v, 2} \leq C(t + |v|^{\frac{1}{p}}), \quad \|\xi_{v, t\nu}\|_{v, p, 1} \leq C(t + |v|^{\frac{1}{p}}), \quad \|\tilde{\eta}_{v, t\nu}\|_{v, p} \leq C(t + |v|^{\frac{1}{p}})^2.$$

Furthermore, if $z \in \Sigma$ and $(B_{b_v}(u_v(z), C\delta), J, g_{V, b_v})$ is isometric to a ball in $\mathbb{C}^n$, then $\tilde{\eta}_{v, t\nu}(z) = 0$.

Remark: In specific applications, the main goal would be to express the number of zeros of $\psi_{\mathcal{S},tv}^\mu$ in terms of the cohomology ring of a closure of $\mathcal{S}_{\mathcal{T}}(\mu)$. One of the significant intermediate steps is to extract the leading-order terms from the section $\psi_{\mathcal{S},tv}^\mu$. If $\lambda_0 = 0$, the estimate on $\nu_{v,t}$ given above easily leads to a sufficiently good estimate on $\pi_{v,-}^{0,1}\nu_{v,t}$; see [I] and [Z2]. In such a case, one can also extract the first-order term from $\pi_{v,-}^{0,1}\bar{\partial}u_v$, which suffices for the computation in [I]. A power-series expansion for $\pi_{v,-}^{0,1}\bar{\partial}u_v$ is given in [Z2], where terms of up to third degree are used. With the choice of metrics in [Z2], the term $\pi_{v,-}^{0,1}\tilde{\eta}_{v,tv}$ vanishes. The remaining term is shown to be secondary for a good choice of the obstruction bundle setup.

Proof: Let $\delta, \epsilon > 0$ be as in Lemma 3.28 and its proof. We take $\varphi_{\mathcal{S},tv}^\mu$ to be the section descendent from the $G_{\mathcal{T}}$ -equivariant section $\tilde{\varphi}_{\mathcal{S},tv}^\mu$. Denote by U'_K the open set U_K of Lemma 3.28. By Corollary 4.22, there exists a neighborhood U_K of K in $\bar{C}_{(\lambda^*, N)}^\infty(\Sigma; \mu)$ such that $\mathcal{M}_{\Sigma, tv, \lambda^*}(\mu) \cap U_K$ is contained in $\tilde{\mathcal{M}}_{tv}(U'_K, \delta; \mu)$. The neighborhood U_K can always be chosen to contain all the zeros of the section $\tilde{\psi}_{tv}$ of the bundle $\Pi_-^{0,1}$ over $\mathcal{M}_{\Sigma, tv, \lambda^*}(\mu) \cap U_K$. By Corollary 3.26, $\mathcal{M}_{\Sigma, tv, \lambda^*}(\mu) \cap U_K$ is precisely the oriented zero set of the section $\tilde{\psi}_{tv}$. Since the map

$$F^\emptyset \mathcal{S}_\delta|K \longrightarrow \tilde{\mathcal{M}}_{tv}(U'_K, \delta; \mu), \quad v \longrightarrow \tilde{\gamma}_{\mathcal{T}, tv}(\Phi_{\mathcal{S}} \Phi_{\mathcal{S}}^\mu(\varphi_{\mathcal{S}, tv}^\mu(v))),$$

is an orientation-preserving diffeomorphism by Lemma 3.28, it induces a sign-preserving bijection between the zero set of $\tilde{\psi}_{tv}$ on $\tilde{\mathcal{M}}_{tv}(U'_K, \delta; \mu)$, and the zero set of the section

$$(\tilde{\gamma}_{\mathcal{T}, tv} \Phi_{\mathcal{S}} \Phi_{\mathcal{S}}^\mu \varphi_{\mathcal{S}, tv}^\mu)^* \tilde{\psi}_{tv} \in \Gamma(F^\emptyset \mathcal{S}_\delta|K; (\tilde{\gamma}_{\mathcal{T}, tv} \Phi_{\mathcal{S}} \Phi_{\mathcal{S}}^\mu \varphi_{\mathcal{S}, tv}^\mu)^* \Pi_-^{0,1}).$$

By equation (3.17), under the canonical identification

$$(\tilde{\gamma}_{\mathcal{T}, tv} \gamma_{\mathcal{T}}^{-1})^* \Pi_-^{0,1} = \Gamma_-^{0,1} \longrightarrow \mathcal{M}(U'_K, \delta),$$

the section $(\tilde{\gamma}_{\mathcal{T}, tv} \gamma_{\mathcal{T}}^{-1})^* \tilde{\psi}_{tv}$ corresponds to the section ψ_{tv} given by

$$\begin{aligned} \psi_{tv}(b(v)) &= tv|_{u_v} - \bar{\partial}u_v - \tilde{\pi}_{v,+}^{0,1}\eta_{v,tv} - N_{v,tv}P_v\eta_{v,tv} \\ &= \pi_{v,-}^{0,1}(tv|_{u_v} - \bar{\partial}u_v - \tilde{\pi}_{v,+}^{0,1}\eta_{v,tv} - N_{v,tv}P_v\eta_{v,tv}) \\ &= \pi_{v,-}^{0,1}(tv_{v,\xi_{v,tv}} - \bar{\partial}u_v - \tilde{\pi}_{v,+}^{0,1}\eta_{v,tv} - N_vP_v\eta_{v,tv}). \end{aligned} \tag{3.19}$$

The second equality above is automatic, since $\psi_{tv}(b(v)) \in \Gamma_-^{0,1}(v)$; the third follows from the definition of $N_{v,tv}$ in Subsection 3.6. The bounds on the terms $\nu_{v,\xi_{v,tv}}$, $\eta_{v,tv}$, $N_vP_v\eta_{v,tv}$ also follow from Subsection 3.6. By definition of $\Gamma_-^{0,1}$ in Subsection 3.4 and equation (3.19), under the canonical identification

$$\gamma_{\mathcal{T}}^* \Gamma_-^{0,1} = \pi_{F\mathcal{T}}^* \Gamma_-^{0,1} \longrightarrow F^\emptyset \mathcal{T}_\delta|U'_K,$$

the section $\gamma_{\mathcal{T}}^* \psi_{tv}$ corresponds to the section $\psi_{\mathcal{T}, tv}$, described in the statement of the theorem.

The next proposition describes a special case of the above theorem. It is obtained by fixing a metric g on Σ and going through the construction analogous to that in Subsection 3.6 and then modification for constraints as above. The sign statement below follows from the fact that the (L^2, g, g_V) -projection $\ker(D_{V,u_b} - \tau t L_{\nu, u_b}) \longrightarrow \ker D_{V, u_b}$ is an isomorphism for all $\tau \in [0, 1]$, t sufficiently small, and $b \in \mathcal{M}_{\Sigma, tv, \lambda^*}(\mu)$ sufficiently close to K .

Proposition 3.30 Suppose $\lambda^* \in H_2(V; \mathbb{Z})$, $\mathcal{T} = (\Sigma, [N], \{\hat{0}\}; \hat{0}, \lambda^*)$ is a (V, J) -regular bubble type,

$$\nu \in \Gamma^{0,1}(\Sigma \times V; \Lambda_{J,j}^{0,1} \pi_\Sigma^* T^* \Sigma \otimes \pi_V^* TV)$$

is any section, and μ is a $\mathcal{T}$ -regular N -tuple of submanifolds of V of total codimension

$$\text{codim } \mu = 2(\langle c_1(V, J), \lambda^* \rangle + (\dim V)(1 - g(S)) + |N|).$$

Then $\mathcal{M}_{\mathcal{T}}(\mu)$ is a discrete set and for every finite subset K of $\mathcal{S}(\mu)$, there exist a neighborhood U_K of K in $\bar{C}_{(\lambda^*, N)}^\infty(\Sigma; \mu)$, $\epsilon > 0$, and for each $t \in (0, \epsilon)$ a sign-preserving bijection between K and $\mathcal{M}_{\Sigma, t\nu, \lambda^*}(\mu) \cap U_K$.

3.9 Gluing Maps for Spaces $\bar{\mathcal{U}}_{\mathcal{T}}^{(0)}(\mu)$ and Orientations

We now return to spaces the case $\mathcal{T} = (S, M, I; j, \lambda)$ is a regular bubble type. Our primary interest is in the case $S = S^2$, so we assume that this is the case. However, most of the analysis in this subsection applies to any regular bubble type $\mathcal{T}$. Let μ be a generic $\tilde{M}$ -tuple of submanifolds in V , as defined below. If $I = \bigsqcup_{k \in K} I_k$ is the decomposition of I into rooted trees and $\{\mathcal{T}_k\}$ are the corresponding simple types derived from $\mathcal{T}$, the product gluing map,

$$(\tilde{\gamma}_{\mathcal{T}_k})_{k \in K} : \prod_{k \in K} F\mathcal{T}_{k, \delta_k} \longrightarrow \prod_{k \in K} \bar{\mathcal{M}}_{\langle \mathcal{T}_k \rangle},$$

may not map the total space of the bundle over $\mathcal{U}_{\mathcal{T}}^{(0)}(\mu)$ into $\bar{\mathcal{U}}_{\langle \mathcal{T} \rangle}^{(0)}(\mu)$. In this subsection, we remedy this deficiency of the product gluing map. We also show that all the spaces $\bar{\mathcal{U}}_{\langle \mathcal{T} \rangle}^{(0)}(\mu)$ and $\bar{\mathcal{U}}_{\langle \mathcal{T} \rangle}(\mu)$ are naturally oriented topological orbifolds and the gluing maps defined below preserve orientations.

Definition 3.31 If $\mathcal{T}$ is a (V, J) -regular bubble type, tuple μ of oriented submanifolds of V is $\mathcal{T}$ -regular if

$$\bigoplus_{l \in M} T_{u_{j_l}(y_l)} V = \text{Im } \text{dev}_{[N]}|_b + \bigoplus_{l \in M} T_{u_{j_l}(y_l)} \mu_l \quad \forall b = (S^2, M, I; x, (j, y), u) \in \mathcal{B}_{\mathcal{T}}(\mu);$$

where $\text{dev}_{[N]}|_b : \tilde{\mathcal{K}}_b \mathcal{T} \longrightarrow \bigoplus_{l \in M} T_{u_{j_l}(y_l)} V$, $\text{dev}_l|_b(\xi_I, w_{\hat{I}+[N]}) = \xi_{j_l}(y_l) + du_{j_l}|_{y_l} w_l$,

Let $\mathcal{T}$, $\mathcal{T}_k$, K , μ , and b be as above. Denote by $b_k = (S^2, M_k, I_k; x|_{\hat{I}_k}, (j, y)|_{M_k}, u|_{I_k})$ the corresponding $\mathcal{T}_k$ -bubble map; see Subsection 2.5. Let $\mathcal{N}_b^\mu \mathcal{T}$ be the (L^2, b) -orthogonal complement of

$$\tilde{\mathcal{K}}_b^\mu \mathcal{T} = \{(\xi, w_{\hat{I}+M}) \in \tilde{\mathcal{K}}_b \mathcal{T} : \xi_{j_l}(y_l) + du_{j_l}|_{y_l} w_l \in T_{u_{j_l}(y_l)} \mu_l \ \forall l \in M \cap \tilde{M}\} \subset \tilde{\mathcal{K}}_b \mathcal{T}$$

in $\bigoplus_{k \in K} \mathcal{K}_{b_k} \mathcal{T}_k$. Denote by $\tilde{\mathcal{N}}^\mu \mathcal{T} \longrightarrow \mathcal{B}_{\mathcal{T}}(\mu)$ and $\mathcal{N}^\mu \mathcal{T} \longrightarrow \mathcal{U}_{\mathcal{T}}^{(0)}(\mu)$ the corresponding vector bundles. Let

$$\mathcal{N}^\mu \mathcal{B}_{\mathcal{T}} = \tilde{\mathcal{N}}^\mu \mathcal{T} \oplus (\mathbb{C} \oplus \mathbb{R})^K \longrightarrow \mathcal{B}_{\mathcal{T}}(\mu), \quad \mathcal{N}^\mu \mathcal{U}_{\mathcal{T}}^{(0)} = \mathcal{N}^\mu \mathcal{T} \oplus (\mathbb{C} \oplus \mathbb{R})^K \longrightarrow \mathcal{U}_{\mathcal{T}}^{(0)}(\mu);$$

$$F^{(0)} \mathcal{T} = \bigoplus_{k \in K} F^{(0)} \mathcal{T}_k \longrightarrow \mathcal{B}_{\mathcal{T}}, \quad F\mathcal{T} = \bigoplus_{k \in K} F\mathcal{T}_k \longrightarrow \mathcal{U}_{\mathcal{T}}^{(0)}.$$

The last two vector bundles carry norms induced from the norms on $F^{(0)}\mathcal{T}_k$, while we define norms on the first two by

$$|(b, \sigma, (c, r))| = \|\sigma\|_{b, C^0} + |(c, r)|,$$

if $\sigma \in \mathcal{N}_b^\mu \mathcal{T} \subset \bigoplus_{k \in K} \tilde{\mathcal{K}}_{b_k} \mathcal{T}_k$ and $(c, r) \in (\mathbb{C} \oplus \mathbb{R})^K$. If δ is sufficiently small, define

$$\tilde{\phi}_T^\mu: \mathcal{N}^\mu \mathcal{B}_{T, \delta} \longrightarrow \prod_{k \in K} \mathcal{M}_{T_k}^{(0)} \quad \text{by} \quad \tilde{\phi}_T^\mu(\sigma, (c, r)) = ((c_k, r_k) \cdot H_{T_k}^{(0)}(\sigma_k))_{k \in K} \in \prod_{k \in K} \mathcal{M}_{T_k}^{(0)},$$

where $H_{T_k}^{(0)}$ is as at the end of Subsection 3.2 and $(c_k, r_k) \cdot$ denotes the action of a neighborhood of

$$0 \in \mathbb{C} \times \mathbb{R} = \mathbb{C} \times \mathbb{R} \times \{0\} \subset \mathbb{C} \times \mathbb{R} \times \mathbb{R}$$

described in Subsection 3.1. Since $\tilde{\phi}_T^\mu$ is $(\tilde{G}_T \times \text{Aut}(\mathcal{T}))$ -equivariant, it descends to a G_T^* -equivariant map

$$\phi_T^\mu: \mathcal{N}^\mu \mathcal{U}_{T, \delta}^{(0)} \longrightarrow \prod_{k \in K} \mathcal{M}_{T_k}.$$

Let $\Phi_T^\mu: \pi_{\mathcal{N}^\mu \mathcal{U}_T}^* F\mathcal{T} \longrightarrow F\mathcal{T}$ be a G_T^* -equivariant vector-bundle map covering the map ϕ_T^μ such that Φ_T^μ is the identity over $\mathcal{U}_T^{(0)}(\mu)$. Denote by $\tilde{\Phi}_T^\mu$ the lift of Φ_T^μ to $\mathcal{N}^\mu \mathcal{B}_{T, \delta}$. Let $\Phi_{T, k}^\mu$ and $\tilde{\Phi}_{T, k}^\mu$ be k th components of Φ_T^μ and $\tilde{\Phi}_T^\mu$, respectively.

Lemma 3.32 *With notation as above, there exist $(\tilde{G}_T \times \text{Aut}(\mathcal{T}))$ -invariant functions $\delta, C \in C^\infty(\mathcal{B}_T(\mu); \mathbb{R}^+)$ and a $(\tilde{G}_T \times \text{Aut}(\mathcal{T}))$ -equivariant section*

$$\tilde{\varphi}_T^\mu \in \Gamma(F^{(0)}\mathcal{T}_\delta; \pi_{F^{(0)}\mathcal{T}}^* \mathcal{N}^\mu \mathcal{B}_{T, \delta}),$$

such that $|\tilde{\varphi}_T^\mu(v)| \leq C(b_v)|v|^{\frac{1}{p}}$ and

$$F\mathcal{T}_\delta \longrightarrow \bar{\mathcal{U}}_{\langle T \rangle}^{(0)}(\mu), \quad v \longrightarrow (\tilde{\gamma}_{T_k}(\Phi_{T, k}^\mu \varphi_T^\mu(v)))_{k \in K},$$

is a homeomorphism onto an open neighborhood of $\mathcal{U}_T^{(0)}(\mu)$ in $\bar{\mathcal{U}}_{\langle T \rangle}^{(0)}(\mu)$. Furthermore, the restriction of this map to $F^0\mathcal{T}_\delta$ is an orientation-preserving diffeomorphism onto an open subset of $\mathcal{U}_{\langle T \rangle}^{(0)}(\mu)$.

Proof: Denote by $\mathcal{N}_T \mu$ the normal bundle of

$$X_T(\mu) \equiv \{x_K \in V^K: x_{k_1} = x_{k_2} \ \forall k_1, k_2 \in K\} \times \prod_{l \in M \cap \tilde{M}} \mu_l \subset V^K \times V^{M \cap \tilde{M}}.$$

Let $\tilde{\mathcal{N}}_T \mu = \mathcal{N}_T \mu \oplus (\mathbb{C} \oplus \mathbb{R})^K$. Since the $\tilde{G}_T$ -action does not change any evaluation maps and the constraints are in general position, the differential of the map

$$\begin{aligned} \Psi_{T, \tilde{M}}: \prod_{k \in K} \mathcal{M}_{T_k}^{(0)} &\longrightarrow V^K \times V^{M \cap \tilde{M}} \times (\mathbb{C} \times \mathbb{R})^K, \\ \Psi_{T, \tilde{M}} &= \left((\text{ev}_{T_k}(b_k))_{k \in K}; (\text{ev}_l(b))_{l \in M \cap \tilde{M}}; (\Psi_{T_k, \hat{0}_{I_k}}(b_k))_{k \in K} \right), \end{aligned}$$

where $\Psi_{\mathcal{T}_k, \hat{0}_{I_k}}(b_k) \in \mathbb{C} \times \mathbb{R}$ is as in Subsection 3.2, induces an isomorphism between $\mathcal{N}_b^\mu \mathcal{B}_{\mathcal{T}}$ and $\tilde{\mathcal{N}}_{\mathcal{T}} \mu$. This isomorphism is orientation-preserving by definition of orientations. Thus,

$$\tilde{\phi}_{\mathcal{T}}^\mu : \mathcal{N}^\mu \mathcal{B}_{\mathcal{T}, \delta} \longrightarrow \prod_{k \in K} \mathcal{M}_{\mathcal{T}_k}^{(0)}$$

is an orientation-preserving diffeomorphism onto an open neighborhood of $\mathcal{B}_{\mathcal{T}}(\mu)$ in $\prod_{k \in K} \mathcal{M}_{\mathcal{T}_k}^{(0)}$, provided $\delta \in C^\infty(\mathcal{B}_{\mathcal{T}}(\mu); \mathbb{R}^+)$ is sufficiently small. By the same argument as in Subsection 3.8, for any simple bubble type $\mathcal{T}'$, the map

$$\tilde{\gamma}_{\mathcal{T}'} : F^\emptyset \mathcal{T}'_\delta \longrightarrow \mathcal{M}_{\langle \mathcal{T}' \rangle} = \mathcal{H}_{\langle \mathcal{T}' \rangle}$$

is an orientation-preserving diffeomorphism onto an open subset of $\mathcal{M}_{\langle \mathcal{T}' \rangle}$ provided $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}'}; \mathbb{R}^+)$ is sufficiently small. Along with Corollary 4.23, this implies that the product map

$$\prod_{k \in K} \tilde{\gamma}_{\mathcal{T}_k} : \prod_{k \in K} F \mathcal{T}_{k, \delta} \longrightarrow \prod_{k \in K} \bar{\mathcal{M}}_{\langle \mathcal{T}_k \rangle}$$

is a homeomorphism onto an open neighborhood of $\prod_{k \in K} \mathcal{M}_{\mathcal{T}_k}$ in $\prod_{k \in K} \bar{\mathcal{M}}_{\langle \mathcal{T}_k \rangle}$ and its restriction to the preimage of $\prod_{k \in K} \mathcal{M}_{\langle \mathcal{T}_k \rangle}$ is an orientation-preserving diffeomorphism. The lemma now follows by applying an argument similar to the proof of Lemma 3.28 to the functions

$$h(v) = \Psi_{\mathcal{T}, \tilde{M}}(\tilde{\phi}_{\mathcal{T}}^\mu(\sigma, (c, r))), \quad \tilde{h}(v) = \tilde{h}_0(v) = \Psi_{\mathcal{T}, \tilde{M}}((\tilde{\gamma}_k \Phi_{\mathcal{T}, k}^\mu(v))_{k \in K}),$$

where we write $v = (\sigma, (c, r), v)$, with $(\sigma, (c, r)) \in \mathcal{N}^\mu \mathcal{B}_{\mathcal{T}}$ and $v \in F^{(0)} \mathcal{T}$. Since $\mathcal{B}_{\mathcal{T}}(\mu)$ is generally not precompact in $\prod_{k \in K} \mathcal{M}_{\mathcal{T}_k}^{(0)}$, we end up with $\delta, C \in C^\infty(\mathcal{B}_{\mathcal{T}}(\mu); \mathbb{R}^+)$, instead of $\delta, C \in \mathbb{R}^+$. Another difference is that $\tilde{h}$ is not necessarily smooth with respect to the standard smooth structure on $\mathcal{N}^\mu \mathcal{B}_{\mathcal{T}} \oplus F^{(0)} \mathcal{T}$. However, we can put a smooth structure on the total space such that the composite maps

$$\mathcal{N}^\mu \mathcal{B}_{\mathcal{T}, \delta} \oplus F^{(0)} \mathcal{T} \longrightarrow F^{(0)} \mathcal{T} \longrightarrow \mathbb{R}, \quad v \longrightarrow \tilde{\Phi}_{\mathcal{T}}^\mu(v), \quad v_h \longrightarrow |v_h|^{\frac{1}{3p}}, \quad h \in \hat{I}_k, \quad k \in K,$$

are smooth, whenever $\delta \in C^\infty(\mathcal{B}_{\mathcal{T}}(\mu); \mathbb{R}^+)$ is sufficiently small. Then by Corollary 4.5, $\tilde{h}$ is C^2 , which is sufficient for the arguments of Subsection 3.7. Finally, in the given case $\tilde{h}$ is defined on all of $(\mathcal{N}^\mu \mathcal{B}_{\mathcal{T}} \oplus F^{(0)} \mathcal{T})_\delta$ and thus the second condition on $\tilde{h}_t$ in Subsection 3.7 is redundant.

Suppose $\mathcal{T}$ is a bubble type and μ is an $\tilde{M}$ -tuple of constraints in general position. By Lemma 3.32, there exist $G_{\mathcal{T}}^*$ -invariant functions $\delta, C \in C^\infty(\mathcal{U}_{\mathcal{T}}^{(0)}(\mu); \mathbb{R}^+)$ and a $G_{\mathcal{T}}^*$ -equivariant section

$$\varphi_{\mathcal{T}}^\mu \in (F \mathcal{T}_\mu; \pi_{F \mathcal{T}}^* \mathcal{N}^\mu \mathcal{U}_{\mathcal{T}, \delta}^{(0)})$$

such that $|\varphi_{\mathcal{T}}^\mu(v)| \leq C(b_v)|v|^{\frac{1}{p}}$ and

$$\tilde{\gamma}_{\mathcal{T}}^\mu : F \mathcal{T}_\delta \longrightarrow \bar{\mathcal{U}}_{\langle \mathcal{T} \rangle}^{(0)}(\mu), \quad \tilde{\gamma}_{\mathcal{T}}^\mu(v) = (\tilde{\gamma}_{\mathcal{T}_k}(\Phi_{\mathcal{T}, k}^\mu \varphi_{\mathcal{T}}^\mu(v)))_{k \in K},$$

is a homeomorphism onto a neighborhood of $\mathcal{U}_T^{(0)}(\mu)$ in $\bar{\mathcal{U}}_T^{(0)}(\mu)$, which is an orientation-preserving diffeomorphism on a dense open subset of the domain. If T' is another regular bubble type such that $\langle T \rangle = \langle T' \rangle$ and μ is T' -regular, it follows that

$$\tilde{\gamma}_T^\mu{}^{-1} \tilde{\gamma}_{T'}^\mu : \tilde{\gamma}_{T'}^\mu{}^{-1}(\tilde{\gamma}_T^\mu(F\mathcal{T}_\delta)) \longrightarrow \tilde{\gamma}_T^\mu{}^{-1}(\tilde{\gamma}_{T'}^\mu(F\mathcal{T}'_{\delta'}))$$

is a homeomorphism provided $\delta' \in C^\infty(\mathcal{U}_{T'}^{(0)}(\mu); \mathbb{R}^+)$ is sufficiently small. Furthermore, by the above it is orientation-preserving on a dense open subset of its domain. It follows that $\tilde{\gamma}_T^{\mu^{-1}} \tilde{\gamma}_{T'}^\mu$ is an orientation-preserving homeomorphism everywhere. We thus obtain

Theorem 3.33 *Let $T^* = (S^2, M, I^*; j, \lambda^*)$ be a basic bubble type and μ an M -tuple of constraints such that μ is T -regular for every bubble type $T \leq T^*$.*

- (1) *The spaces $\bar{\mathcal{U}}_{T^*}^{(0)}(\mu)$ and $\bar{\mathcal{U}}_{T^*}(\mu)$ are oriented topological orbifolds.*
- (2) *Suppose $T < T^*$, $\phi_T^\mu : \mathcal{N}^\mu \mathcal{T}_\delta \longrightarrow \mathcal{U}_T^{(0)}$ is a G_{T^*} -equivariant identification of neighborhoods of $\mathcal{U}_T^{(0)}(\mu)$ in $\mathcal{N}^\mu \mathcal{T}$ and in $\mathcal{U}_T^{(0)}$, and $\Phi_T^\mu : \pi_{\mathcal{N}^\mu \mathcal{T}}^* F\mathcal{T} \longrightarrow F\mathcal{T}$ is a lift of ϕ_T^μ such that $\Phi_T^\mu|_{\mathcal{U}_T^{(0)}(\mu)} = 1$. Then there exist G_{T^*} -invariant functions $\delta, C \in C^\infty(\mathcal{U}_T^{(0)}(\mu); \mathbb{R}^+)$ and a G_{T^*} -equivariant continuous orientation-preserving identification*

$$\tilde{\gamma}_T^\mu : F\mathcal{T}_\delta \longrightarrow \bar{\mathcal{U}}_{T^*}^{(0)}(\mu),$$

of neighborhoods of $\mathcal{U}_T^{(0)}(\mu)$ in $F\mathcal{T}$ and in $\bar{\mathcal{U}}_{T^}^{(0)}(\mu)$, which is smooth on $F^\emptyset \mathcal{T}_\delta \longrightarrow \mathcal{U}_{T^*}(\mu)$. Furthermore, for every $v \in F\mathcal{T}_\delta$, there exists $\sigma(v) \in \mathcal{N}^\mu \mathcal{T}$ such that*

$$\|\sigma(v)\|_{b^*} \leq C(b^*)|v|^{\frac{1}{p}} \quad \text{and} \quad u_{\tilde{\gamma}_T^\mu(v)} = \exp_{V, u_{b'} \circ q_{\Phi_T^\mu(\sigma(v))}} \xi_v, \quad \text{where} \quad \Phi_T^\mu(\sigma(v)) \equiv [b', v'],$$

for some $\xi_v \in \Gamma(u_{b'} \circ q_{\Phi_T^\mu(\sigma(v))})$ with $\|\xi_v\|_{C^0} \leq C(b_v)|v|^{\frac{1}{p}}$.

Remark: The descriptive statement (2) of Theorem 3.33 is used in [Z2] for local excess-intersection type of computations on the spaces $\bar{\mathcal{U}}_{T^*}(\mu)$.

4 Technical Issues

4.1 Continuity of the Gluing Map

Let $\mathcal{T} = (S, M, I; j, \lambda)$ be a simple regular bubble type and H a nonempty subset of $\hat{I}$. Suppose $v_k \in F^{(\emptyset)} \mathcal{T}_\delta$ and the sequence v_k converges to $v^* \in F^{(H)} \mathcal{T}_\delta$. In this section, we show that $\tilde{\gamma}_T(v_k)$ converges to $\tilde{\gamma}_T(v^*)$ in the Gromov topology. Our main interest is the case $S = S^2$.

It is sufficient to assume that $\pi_h(v_k) = \pi_h(v^*)$ if $h \notin H$. In particular, $b \equiv b_{v^*} = b_{v_k}$. Denote by $\tilde{\mathcal{T}} = \mathcal{T}(H)$ the bubble type of $b(v^*)$. For each k , define

$$\tilde{v}_k = (\tilde{b}(v^*), (\tilde{v}_k)_H) \in F_{b(v^*)}^{(0)} \tilde{\mathcal{T}}$$

as follows. If $h \in H$, put

$$i_H h = \min \{i < h : \text{if } h' \in I \text{ \& } i < h' < h, h' \notin H\}.$$

Since I is a rooted $i_H h$ is well-defined. Let

$$\tilde{v}_{k,h} = \prod_{i_H h < h' \leq h} v_{k,h'}.$$

Since $v_k \longrightarrow v^*$, $\tilde{v}_{k,h} \longrightarrow 0$ for all $h \in H$. Furthermore, by construction $\Sigma_{v_k} = \Sigma_{\tilde{v}_k}$ and $q_{v_k} = q_{v^*} \circ q_{\tilde{v}_k}$. In particular, $u_{v_k} = u_{v^*} \circ q_{\tilde{v}_k}$.

For any $h \in H$ and $\delta > 0$, let

$$A_{h,\delta,k} = q_{v_k}^{-1} (\{(\iota_h, z) : r_{b,h}(z) \leq \delta\} \cup \{(h, z) : |q_S^{-1}(z)| \leq \delta\}) \subset \Sigma_{v_k},$$

$$A_{h,\delta}^* = q_{v^*}^{-1} (\{(\iota_h, z) : r_{b,h}(z) \leq \delta\} \cup \{(h, z) : |q_S^{-1}(z)| \leq \delta\}) \subset \Sigma_{v^*}, \quad \Sigma_\delta^* = \Sigma_{v^*} - \bigcup_{h \in H} A_{h,\delta}^*.$$

It is convenient to make the following definitions. If $\eta_k \in L^p(v_k)$, the sequence $\{\eta_k\}$ converges to $\eta^* \in L^p(v^*)$ if $q_{\tilde{v}_k}^{-1*} \eta_k$ converges to η^* in the L^p -norm on all precompact open subsets of $\Sigma_{v^*}^*$ and

$$\lim_{\delta \longrightarrow 0} \lim_{k \longrightarrow 0} \|\eta_k\|_{v_k, L^p(A_{h,\delta,k})} = 0 \quad \forall h \in H. \quad (4.1)$$

If $\xi_k \in L_1^p(v_k)$, the sequence $\{\xi_k\}$ converges to $\xi^* \in L_1^p(v^*)$ if $\xi_k \circ q_{\tilde{v}_k}^{-1}$ converges to ξ^* in the L_1^p -norm on all precompact open subsets of $\Sigma_{v^*}^*$ and

$$\lim_{\delta \longrightarrow 0} \lim_{k \longrightarrow 0} \|\xi_k\|_{v_k, L_1^p(A_{h,\delta,k})} = 0 \quad \forall h \in H. \quad (4.2)$$

In (4.1) and (4.2), we use the modified Sobolev norms.

Lemma 4.1 *There exist $C, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for any sequence $\{v_k \in F^{(\emptyset)} \mathcal{T}_\delta\}$ converging to v^* as above and $\xi \in \Gamma_+(v^*)$*

$$\|\pi_{v_k, -}(\xi \circ q_{\tilde{v}_k})\|_{v_k, p, 1} \leq C(b) |v_k - v^*| \|\xi\|_{v^*, p, 1}.$$

Proof: Note that $\Gamma_-(v_k) = \{\xi^- \circ q_{\tilde{v}_k} : \xi^- \in \Gamma(v^*)\}$. Thus, the difference $q_{\tilde{v}_k}^* \pi_{v^*, -} - \pi_{v_k, -} q_{\tilde{v}_k}^*$ arises entirely from the difference between the metrics $q_{\tilde{v}_k}^* g_{v^*}$ and g_{v_k} . By construction, the two metrics differ only on the annuli $A_{h, 2|v_k - v^*|^{\frac{1}{2}}, k}$ for $h \in H$. Thus, the claim follows from (2) of Lemma 3.5.

Lemma 4.2 *If η_k converges to η^* , then $P_{v_k} \eta_k$ converges to $P_{v^*} \eta^*$.*

Proof: (1) Let $\{\epsilon_k\}, \{\delta_k\} \subset \mathbb{R}^+$ be sequences converging to zero such that

$$\|\eta^*\|_{v^*, L^p(A_{h,\delta_k}^*)} \leq \epsilon_k; \quad \|P_{v^*} \eta^*\|_{v^*, L_1^p(A_{h,\delta_k}^*)} \leq \epsilon_k, \quad \lim_{k \longrightarrow \infty} \|\eta_k\|_{v_k, L^p(A_{h,\delta_{k^*}, k})} < \epsilon_k \quad \forall h \in H. \quad (4.3)$$

For every $k^* > 0$, choose N_{k^*} such that for all $k > N_{k^*}$

$$\|q_{\tilde{v}_k}^{-1*} \eta_k - \eta^*\|_{v^*, L^p(\Sigma_{\delta_{k^*}}^*)} \leq \epsilon_{k^*} \quad \text{and} \quad \|\eta_k\|_{v_k, L^p(A_{h,\delta_{k^*}, k})} \leq \epsilon_{k^*} \quad \forall h \in H. \quad (4.4)$$

It can be assumed that $2|v_k - v^*|^{\frac{1}{2}} \leq \delta_{k^*}$, ϵ_{k^*} whenever $k > N_{k^*}$. For any $k > N_{k^*}$, let $\tilde{\eta}_{k^*, k} \in L^p(v^*)$ be given by

$$\tilde{\eta}_{k^*, k} = \begin{cases} q_{\tilde{v}_k}^{-1*} \tilde{\eta}_k, & \text{on } \Sigma_{\delta_{k^*}}^*; \\ 0, & \text{outside of } \Sigma_{\delta_{k^*}}^*. \end{cases}$$

Then $\|\tilde{\eta}_{k^*,k}\|_{v^*,p} \leq \|\eta_k\|_{v_k,p}$. Let

$$\tilde{P}_{k^*,k}\eta_k = q_{\tilde{v}_k}^* P_{v^*} \tilde{\eta}_{k^*,k} \in L_1^p(v_k).$$

Then by Lemma 3.16 and the first assumptions of (4.3) and (4.4),

$$\begin{aligned} \left\| q_{\tilde{v}_k}^{-1*} \tilde{P}_{k^*,k}\eta_k - P_{v^*} \eta^* \right\|_{v^*, L_1^p(\Sigma_{\delta_{k^*}}^*)} &\leq \|P_{v^*} \tilde{\eta}_{k^*,k} - P_{v^*} \eta^*\|_{v^*,p,1} \\ &\leq C(b) \|\tilde{\eta}_{k^*,k} - \eta^*\|_{v^*,p} \leq 2C(b) \epsilon_{k^*}. \end{aligned} \quad (4.5)$$

Since $\|dq_{\tilde{v}_k}\|_{C^0} \leq C(b)$, by (4.3) and the first assumption of (4.4) for all $h \in H$,

$$\begin{aligned} \|\tilde{P}_{k^*,k}\eta_k\|_{v_k, L_1^p(A_{h,\delta_{k^*},k})} &\leq C(b) \|P_{v^*} \tilde{\eta}_{k^*,k}\|_{v^*, L_1^p(A_{h,\delta_{k^*}}^*)} \\ &\leq C(b) \left(\|P_{v^*} \eta^*\|_{v^*, L_1^p(A_{h,\delta_{k^*}}^*)} + \|P_{v^*} \tilde{\eta}_{k^*,k} - P_{v^*} \eta^*\|_{v^*,p,1} \right) \leq C'(b) \epsilon_{k^*}. \end{aligned} \quad (4.6)$$

(2) We now show that $\tilde{P}_{k^*,k}\eta_k$ is close to $P_{v_k}\eta_k$. By Lemmas 3.16 and 4.1,

$$\begin{aligned} \|\tilde{P}_{k^*,k}\eta_k - P_{v_k}\eta_k\|_{v_k,p,1} &\leq C(b) \left(\|D_{v_k} \tilde{P}_{k^*,k}\eta_k - \eta_k\|_{v_k,p} + \|\pi_{v_k, -\tilde{P}_{k^*,k}\eta_k}\|_{v_k,p,1} \right) \\ &\leq C(b) \left(\|D_{v_k} \tilde{P}_{k^*,k}\eta_k - \eta_k\|_{v_k,p} + |v_k - v^*| \|\eta_k\|_{v_k,p} \right). \end{aligned} \quad (4.7)$$

Since $D_{v^*} P_{v^*} \tilde{\eta}_{k^*,k} = \tilde{\eta}_{k^*,k}$ and $q_{\tilde{v}_k}$ is holomorphic outside of the annuli $A_{h,\delta_{k^*},k}$,

$$D_{v_k} \tilde{P}_{k^*,k}\eta_k = \eta_k \quad \text{on} \quad \Sigma_{v_k} - \bigcup_{h \in H} A_{h,\delta_{k^*},k}; \quad (4.8)$$

By equation (4.6),

$$\|D_{v_k} \tilde{P}_{k^*,k}\eta_k\|_{v_k, L^p(A_{h,\delta_{k^*},k})} \leq C(b) \|\tilde{P}_{k^*,k}\eta_k\|_{v_k, L_1^p(A_{h,\delta_{k^*},k})} \leq C'(b) \epsilon_{k^*}. \quad (4.9)$$

Thus, from equations (4.7)-(4.9) and the second assumption of (4.4), we conclude that for all $k > N_{k^*}$

$$\|\tilde{P}_{k^*,k}\eta_k - P_{v_k}\eta_k\|_{v_k,p,1} \leq C(b) \epsilon_{k^*} (1 + \|\eta^*\|_{v^*,p}). \quad (4.10)$$

Since $\|dq_{\tilde{v}_k}^{-1}\|_{C^0} \leq C(b)$ on $\Sigma_{\delta_{k^*}}^*$, by equations (4.5), (4.6), and (4.10),

$$\|q_{\tilde{v}_k}^{-1*} P_{v_k}\eta_k - P_{v^*} \eta^*\|_{v^*, L_1^p(\Sigma_{\delta_{k^*}}^*)} \leq C(b) \epsilon_{k^*} (1 + \|\eta^*\|_{v^*,p}); \quad (4.11)$$

$$\|P_{v_k}\eta_k\|_{v_k, L_1^p(A_{h,\delta_{k^*},k})} \leq C(b) \epsilon_{k^*} (1 + \|\eta^*\|_{v^*,p}) \quad \forall h \in H. \quad (4.12)$$

By equations (4.11) and (4.12), $P_{v_k}\eta_k$ converges to $P_{v^*} \eta^*$.

Lemma 4.3 *There exist $\tilde{C}, \delta \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R}^+)$ such that for all $v^* \in F^{(H)} \mathcal{T}_\delta$ and $h \in H$,*

$$\|P_{v^*} \bar{\partial} u_{v^*}\|_{g_{v^*}, C^1(A_{h,\delta(b_{v^*})}^*)} \leq \tilde{C}(b_{v^*}).$$

Proof: For each $h \in H$, this lemma is obtained by pasting $(P_{v^*} \bar{\partial} u_{v^*})|(A_{h, \delta(b_{v^*}^*)} \cap \Sigma_{v^*, \iota_h})$ and $(P_{v^*} \bar{\partial} u_{v^*})|(A_{h, \delta(b_{v^*}^*)} \cap \Sigma_{v^*, h})$ onto $\Sigma_{b_{v^*}^*, \iota_h}$ and $\Sigma_{b_{v^*}^*, h}$ via a cutoff function. We then use the usual elliptic estimates and Sobolev inequalities on $\Sigma_{b_{v^*}^*, \iota_h}$ and $\Sigma_{b_{v^*}^*, h}$ along with

$$\|P_{v^*} \bar{\partial} u_{v^*}\|_{v^*, p, 1} \leq C(b_{v^*}) |v^*|^{\frac{1}{p}}.$$

The bound obtained in this way is actually $C(b_{v^*}) |v^*|^{\frac{1}{p}}$.

Corollary 4.4 *There exist $C, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for any sequence $v_k \in F^{(0)} \mathcal{T}_\delta$ converging to $v^* \in F^{(H)} \mathcal{T}_\delta$ as above,*

$$\begin{aligned} \|q_{\tilde{v}_k}^{-1*} \eta_{v_k} - \eta_{v^*}\|_{v^*, L^p(\Sigma_{2|v_k - v^*|^{1/2}}^*)} &\leq C(b) |v_k - v^*|^{\frac{1}{p}}; \\ \|\eta_{v_k}\|_{v_k, L^p(A_{h, 2|v_k - v^*|^{1/2}, k})} &\leq C(b) |v_k - v^*|^{\frac{1}{p}} \quad \forall h \in H. \end{aligned}$$

Proof: Let $\delta_k = 2|v_k - v^*|^{\frac{1}{2}}$ and $\epsilon_k = (2\|\beta'\|_{C^0} + \tilde{C}(b)) |v_k - v^*|^{\frac{1}{p}}$, where $\tilde{C}$ is the function given by Lemma 4.3. Put

$$\begin{aligned} \eta^{(0)} &= -\bar{\partial} u_{v^*}, & \eta^{(m+1)} &= -\bar{\partial} u_{v^*} - N_{v^*} P_{v^*} \eta^{(m)} \quad m \geq 0; \\ \eta_k^{(0)} &= -\bar{\partial} u_{v_k}, & \eta_k^{(m+1)} &= -\bar{\partial} u_{v_k} - N_{v_k} P_{v_k} \eta_k^{(m)} \quad m \geq 0. \end{aligned}$$

By Lemma 4.3 and the explicit description of $\bar{\partial} q_{v_k}$ in Lemma 2.2, ϵ_k , δ_k , $\eta^{(0)}$, and $\eta_k^{(0)}$ satisfy (4.3) and (4.4). Suppose $\epsilon_k^{(m)}$ is such that $\epsilon_k^{(m)}$, δ_k , $\eta^{(m)}$, and $\eta_k^{(m)}$ satisfy (4.3) and (4.4). Since the map $q_{\tilde{v}_k}$ is holomorphic on $q_{\tilde{v}_k}^{-1}(\Sigma_{\delta_k}^*)$, by (4.11), (4.12), the estimates in the proof of Lemma 3.18 and the derivation of equation (3.11) in [Z1],

$$\begin{aligned} &\|q_{v_k}^{-1*} N_{v_k} P_{v_k} \eta_k^{(m)} - N_{v^*} P_{v^*} \eta^{(m)}\|_{v^*, L^p(\Sigma_{\delta_k}^*)} = \|N_{v^*} q_{v_k}^{-1*} P_{v_k} \eta_k^{(m)} - N_{v^*} P_{v^*} \eta^{(m)}\|_{v^*, L^p(\Sigma_{\delta_k}^*)} \\ &\leq C(b) \left(\|q_{v_k}^{-1*} P_{v_k} \eta_k^{(m)}\|_{v^*, L^p(\Sigma_{\delta_k}^*)} + \|P_{v_k} \eta_k^{(m)}\|_{v^*, L^p(\Sigma_{\delta_k}^*)} \right) \|q_{v_k}^{-1*} P_{v_k} \eta_k^{(m)} - P_{v^*} \eta^{(m)}\|_{v^*, L^p(\Sigma_{\delta_k}^*)} \\ &\leq C'(b) (\epsilon_{k^*}^{(m)} + |v_k|^{\frac{1}{p}}) \epsilon_{k^*}^{(m)}; \\ &\|N_{v_k} P_{v_k} \eta_k^{(m)}\|_{v_k, L^p(A_{h, \delta_k, k})} \leq C(b) |v_k|^{\frac{1}{p}} \|P_{v_k} \eta_k^{(m)}\|_{v_k, L_1^p(A_{h, \delta_k, k})} \leq C'(b) |v_k|^{\frac{1}{p}} \epsilon_{k^*}^{(m)}. \end{aligned}$$

Thus, we can take $\epsilon_k^{(m+1)} = \epsilon_k^{(m)} + C'(b) (\epsilon_{k^*}^{(m)} + |v_k|^{\frac{1}{p}}) \epsilon_{k^*}^{(m)}$. This sequence is bounded as long as $|v_k|^{\frac{1}{p}}$ is sufficiently small (depending only on b). Since η_{v^*} is the limit in the (v^*, p) -norm of the sequence $\eta^{(m)}$ and η_{v_k} is the limit in the (v_k, p) -norm of the sequence $\eta_k^{(m)}$, the claim follows.

Corollary 4.5 *If $\mathcal{T}$ is a simple regular bubble type, there exist $\delta, C \in C^\infty(\mathcal{M}_{\mathcal{T}}; \mathbb{R}^+)$ such that for any sequence $\{v_k \in F^\emptyset \mathcal{T}_\delta\}$ converging to $v^* \in F^H \mathcal{T}_\delta$, $\tilde{\gamma}(v_k)$ converges to $\tilde{\gamma}(v^*)$ with respect to the Gromov topology. Furthermore,*

$$\begin{aligned} d_V(ev(\tilde{\gamma}(v^*)), ev(\tilde{\gamma}(v_k))) &\leq C(b_{v^*}) |v_k - v^*|^{\frac{1}{p}} \quad \text{if } S = S^2; \\ d_V(ev(\tilde{\gamma}(v^*)), ev(\tilde{\gamma}(v_k))) &\leq C(b_{v^*}) |v_k - v^*|^{\frac{1}{p}} \quad \forall l \in M; \\ \left| \Psi_{\langle \mathcal{T}, \hat{0} \rangle}(\tilde{\gamma}(v_k)) - \Psi_{\mathcal{T}(v^*), \hat{0}}(\tilde{\gamma}(v^*)) \right| &\leq C(b_{v^*}) |v_k - v^*|^{\frac{1}{p}} \quad \text{if } S = S^2. \end{aligned}$$

Proof: It is sufficient to consider the case $\pi_h(v_k) = \pi_h(v^*)$ for all $h \notin H$ if $v^* \in F^{(H)}\mathcal{T}_\delta$. In such a case, $q_{\tilde{v}_k}$ maps the marked points of Σ_{v_k} to the marked points of Σ_{v^*} and $u_{v_k} = u_{v^*} \circ q_{v_k}$. By construction,

$$\tilde{u}_{v_k} = \exp_{b_{v^*}, u_{v_k}} P_{v_k} \eta_{v_k}, \quad \tilde{u}_{v_k} = \exp_{b_{v^*}, u_{v^*}} P_{v^*} \eta_{v^*}.$$

By Corollary 4.5 and the proof of Lemma 4.2,

$$\|q_{\tilde{v}_k}^{-1*} P_{v_k} \eta_{v_k} - P_{v^*} \eta_{v^*}\|_{v^*, L_1^p(\Sigma_{2|v_k - v^*|^{1/2}}^*)} \leq C(b_{v^*}) |v_k - v^*|^{\frac{1}{p}}; \quad (4.13)$$

$$\|P_{v_k} \eta_{v_k}\|_{v_k, L_1^p(A_{h, 2|v_k - v^*|^{1/2}, k})} \leq C(b_{v^*}) |v_k - v^*|^{\frac{1}{p}} \quad \forall h \in H. \quad (4.14)$$

Let $\zeta_k \in \Gamma(\Sigma^*; \tilde{u}_{v^*})$ be given by

$$\exp_{b_{v^*}, \tilde{u}_{v^*}} \zeta_k = \tilde{u}_{v_k} \circ q_{\tilde{v}_k}, \quad \|\zeta_k\|_{C^0} < \text{inj } g_{V, b_{v^*}}.$$

By equation (4.13) and the proof of (2) of Lemma 3.5,

$$\|\zeta_k\|_{C^0(\Sigma_{2|v_k - v^*|^{1/2}}^*)} \leq C(b_{v^*}) \|\zeta_k\|_{v^*, L_1^p(\Sigma_{2|v_k - v^*|^{1/2}}^*)} \leq C'(b_{v^*}) |v_k - v^*|^{\frac{1}{p}}. \quad (4.15)$$

On the other hand, by (4.14) and by the same argument as in (3) of the proof of Lemma 5.12, the variations of $P_{v^*} \eta_{v^*}$ on $A_{h, 2|v_k - v^*|^{1/2}}^*$ and $P_{v_k} \eta_{v_k}$ on $A_{h, 2|v_k - v^*|^{1/2}, k}^*$ are bounded $C(b_v) |v_k - v^*|^{\frac{1}{p}}$. This can be seen from equation (5.33); observe that an argument similar to the proof Lemma 4.3 shows that we can take δ to be any small number bigger than $2|v_k - v^*|^{\frac{1}{2}}$. Equation (4.15) and the small variation on the annuli imply that

$$\sup_{z \in \Sigma_{v_k}} d_V(\tilde{u}_{v^*}(q_{\tilde{v}_k}(z)), \tilde{u}_{v_k}(z)) \leq C(b_{v^*}) |v_k - v^*|^{\frac{1}{p}}.$$

It follows that $\tilde{\gamma}_{\mathcal{T}}(v_k)$ converges to $\tilde{\gamma}_{\mathcal{T}}(v^*)$ in the Gromov topology. The estimate on the evaluation maps is immediate from the above bound. The last estimate follows from equations (4.13) and (4.14), along with a Sobolev estimate on a neighborhood of $\infty \in \Sigma_{v^*, \hat{0}}$ which implies that the C^1 -norm of ζ_k there is bounded by $C(b_{v^*}) |v_k - v^*|^{\frac{1}{p}}$.

4.2 Injectivity of the Gluing Map

The goal of this subsection is to prove that the gluing maps of (3.15) and (3.16) are injective, as long as $\delta \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R}^+)$ is sufficiently small. We start by showing local injectivity on the subspaces $F^H \mathcal{T}_\delta$ of $F \mathcal{T}_\delta$, where H is a subset of $\hat{I}$.

If $\mathcal{T}$ is regular, we are only interested in the case $t=0$. If $\mathcal{T}$ is semiregular, we only consider the case $H = \emptyset$. We use the same notation as in Subsection 3.4. If $\|\varpi\|_v$ is sufficiently small, define $\tilde{\zeta}_{\varpi, t\nu} \in \Gamma(\tilde{u}_{v, t\nu})$ by

$$\exp_{b_v, \tilde{u}_{v, t\nu}} \tilde{\zeta}_{\varpi, t\nu} = u_{\varpi, t\nu}, \quad \|\tilde{\zeta}_{\varpi, t\nu}\|_{b_v, C^0} < \text{inj } g_{V, b}.$$

Lemma 4.6 *There exist $\delta, C \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R}^+)$ such that for all $v \in F^{(H)}\mathcal{T}_\delta$, where $H = \emptyset$ if $\mathcal{T}$ is semiregular, and $\varpi \in \tilde{T}_v F^H \mathcal{T}_{\delta(b_v)}$,*

- (1) $\|S_\varpi N_{\varpi, t\nu} R_\varpi \xi - N_{v, t\nu} \xi\|_{v, p} \leq C(b_v) \|\varpi\|_v \|\xi\|_{v, p, 1}^2$ for all $\xi \in \Gamma(u_v)$ with $\|\xi\|_{v, p, 1} \leq \delta(b_v)$ and $t \in [0, 1]$;
- (2) $\|S_\varpi \tilde{\pi}_{\varpi, \pm} R_\varpi \xi - \tilde{\pi}_{v, \pm} \xi\|_{v, p, 1} \leq C(b_v) \|\varpi\|_v \|\xi\|_{v, p, 1}$ for all $\xi \in \Gamma(u_v)$;
- (3) $\|S_\varpi P_\varpi R_\varpi \eta - P_v \eta\|_{v, p, 1} \leq C(b_v) \|\varpi\|_v \|\eta\|_{v, p}$ for all $\xi \in \Gamma^{0,1}(u_v)$.

Proof: Claim (1) follows from (2) of Lemma 3.6 and Riemannian geometry estimates such as in [Z1]. Claim (2) is a consequence of (5) of Lemma 3.6 and (2) of Definition 3.11. Finally, (3) is obtained from (1), (2), Definitions 3.11 and 3.13, and Lemmas 3.6 and 3.16 as follows. Writing $\Delta_{\varpi}P$ for $S_{\varpi}P_{\varpi}R_{\varpi} - P_v$, etc.,

$$\begin{aligned}\Delta_{\varpi}P &= P_v\pi_{v,+}^{0,1}D_v\tilde{\pi}_{v,+} + \Delta_{\varpi}P + \tilde{\pi}_{v,-}S_{\varpi}P_{\varpi}R_{\varpi} = P_v\pi_{v,+}^{0,1}D_v\Delta_{\varpi}P - (P_v\pi_{v,+}^{0,1}D_v - 1)\Delta_{\varpi}\tilde{\pi}_{v,+}S_{\varpi}P_{\varpi}R_{\varpi} \\ &= P_v\pi_{v,+}^{0,1}\Delta_{\varpi}\tilde{\pi}_{v,+}^{0,1} - \left(P_v\pi_{v,+}^{0,1}\Delta_{\varpi}D + (P_v\pi_{v,+}^{0,1}D_v - 1)\Delta_{\varpi}\tilde{\pi}_{v,+}\right)S_{\varpi}P_{\varpi}R_{\varpi} \\ &= -P_v\Delta_{\varpi}\pi_{v,+}^{0,1}S_{\varpi}\tilde{\pi}_{\varpi,+}^{0,1}R_{\varpi} - \left(P_v\pi_{v,+}^{0,1}\Delta_{\varpi}D + (P_v\pi_{v,+}^{0,1}D_v - 1)\Delta_{\varpi}\tilde{\pi}_{\varpi,+}S_{\varpi}P_{\varpi}R_{\varpi}\right).\end{aligned}$$

Corollary 4.7 *There exist $\delta, C \in C^{\infty}(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for all $t \in [0; \delta(b_v)]$, $v \in F^{(H)}\mathcal{T}_{\delta}$, where $H = \emptyset$ if $\mathcal{T}$ is semiregular, and $\varpi \in \tilde{T}_v F^H \mathcal{T}_{\delta(b_v)}$,*

$$C(b_v)^{-1}\|\varpi\|_v \leq \|\tilde{\zeta}_{\varpi,t\nu}\|_{v,p,1} + \sum_{h \in H} |w_h(\varpi)|_{g_v} + \sum_{l \in M} |w_l(\varpi)|_{g_v} \leq C(b_v)\|\varpi\|_v.$$

Furthermore, $\|S_{\varpi}\xi_{\varpi,t\nu} - \xi_{v,t\nu}\|_{v,p,1} \leq C(b_v)(t + |v|^{\frac{1}{p}})\|\varpi\|_v$.

Proof: The first claim of the lemma is immediate from the second and (1) of Lemma 3.6. On the other hand, by construction in Subsection 3.6,

$$\xi_{\varpi,t\nu} = tP_{\varpi}\nu - P_{\varpi}\bar{\partial}u_{\varpi} - P_{\varpi}N_{\varpi,t\nu}\xi_{\varpi,t\nu}.$$

Thus, if t and $|v|$ are sufficiently small (depending on b_v), the second claim follows from Lemmas 3.6, 4.6, Corollary 3.19, and equation (3.11).

Corollary 4.8 *If $\mathcal{T}$ is a simple bubble type and K is an open subset of $\mathcal{M}_{\mathcal{T}}$ with compact closure, there exists $\delta > 0$ such that for any $t \in [0, \delta]$, the map*

$$\tilde{\gamma}_{\mathcal{T},t\nu}: F^{\emptyset}\mathcal{T}_{\delta}|K \longrightarrow C_{(\lambda^*, L)}^{\infty}(S; V), \quad \tilde{\gamma}_{\mathcal{T},t\nu}(v) = \tilde{b}_{t\nu}(v),$$

is a differentiable embedding.

Proof: We first deduce from Corollary 4.7 that $\tilde{\gamma}_{\mathcal{T},t\nu}$ is injective if δ is sufficiently small. Suppose not, i.e. there exist sequences $v_k, v_k' \in F^{\emptyset}\mathcal{T}_{\delta}|K$ such that

$$v_k \longrightarrow b \in \bar{K}, \quad v_k' \longrightarrow b' \in \bar{K}, \quad \text{and} \quad \tilde{b}_{t\nu}(v_k) = \tilde{b}_{t\nu}(v_k').$$

It follows that $b = b'$, after possibly modifying the sequence $\{v_k'\}$ by actions of $G_{\mathcal{T}}$. If for some k , $v_k' = v_k(\varpi_k)$ with $\|\varpi_k\|_{v_k}$ sufficiently small, then by Corollary 4.7, $v_k' = v_k$. Otherwise, the difference between q_{v_k} and $q_{v_k'}$ is uniformly bounded below outside of the preimage of the zeroth component and the necks $A_{v_k, h}$. Thus, the bubble maps $b(v_k)$ and $b(v_k')$ are far apart unless b has an automorphism. In the latter case, v_k' can be replaced by an equivalent element of $F^{\emptyset}\mathcal{T}_{\delta}$. In the former case, $\tilde{u}_{v_k}$ and $\tilde{u}_{v_k'}$ cannot be the same because

$$\|P_{v_k}\eta_{v_k,t\nu}\|_{C^0} \leq C(t + |v_k|^{\frac{1}{p}}) \leq C'\delta^{\frac{1}{p}} \quad \text{and} \quad \|P_{v_k'}\eta_{v_k',t\nu}\|_{C^0} \leq C(t + |v_k'|^{\frac{1}{p}})\delta^{\frac{1}{p}}.$$

Thus $\tilde{\gamma}_{\mathcal{T},t\nu}$ is injective on $F^{\emptyset}\mathcal{T}_{\delta}|K$ provided δ is sufficiently small (depending on K). The smoothness of $\tilde{\gamma}_{\mathcal{T},t\nu}$ follows from the smooth dependence of solutions of equation (3.12) on the parameters. Finally, the differential of $\tilde{\gamma}_{\mathcal{T},t\nu}$ is nondegenerate by Corollary 4.7.

Corollary 4.9 *If $\mathcal{T}$ is regular, there exists $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}}; \mathbb{R}^+)$ such that for all m , the map*

$$\tilde{\gamma}_{\mathcal{T}} : \bigcup_{|H|=m} F^H \mathcal{T}_\delta \longrightarrow \bigcup_{|H|=m} \mathcal{M}_{\mathcal{T}(H)}, \quad \tilde{\gamma}_{\mathcal{T}}(v) = \tilde{b}(v),$$

is injective.

Proof: The same argument as in the proof of Corollary 4.8 shows that map

$$\tilde{\gamma}_{\mathcal{T}} : F^H \mathcal{T}_\delta \longrightarrow \mathcal{H}_{\mathcal{T}(H)}$$

is an embedding if δ is sufficiently small. It remains to see that $\tilde{\gamma}_{\mathcal{T}}^{(0)}(v) \neq g \tilde{\gamma}_{\mathcal{T}}^{(0)}(v')$ for any $g \in G_{\mathcal{T}(H)}$ whenever $[v] \neq [v']$. For each $v \in F^{(H)} \mathcal{T}_\delta$ and $i \in H$, we construct $(c_i(v), r_i(v)) \in \mathbb{C} \times \mathbb{R}$ such that

$$(c(v), r(v)) \cdot \tilde{\gamma}_{\mathcal{T}}^{(0)}(v) \in \mathcal{M}_{\mathcal{T}(H)}^{(0)}.$$

We define $c_i(v) \in \mathbb{C}$ and $r_i(v) \in \mathbb{R}$ by

$$\begin{aligned} \tilde{\Psi}((c_i(v), 0) \cdot \tilde{u}_{v,i}) + \sum_{\iota_h(v)=i} d_h(\mathcal{T}(H))(x_h(v) + c_i(v)) + \sum_{j_l(v)=i} (y_l(v) + c_i(v)) &= 0; \\ \Psi^{(3)}((c_i(v), r_i(v)) \cdot \tilde{u}_{v,i}) + \sum_{\iota_h(v)=i} d_h(\mathcal{T}(H))\beta((1 + r_i(v))|x_h(v) + c_i(v)|) \\ &+ \sum_{j_l(v)=i} \beta((1 + r_i(v))|y_l(v) + c_i(v)|) = \frac{1}{2}. \end{aligned}$$

Since the metric $g_{v,i}$ for $i > 0$ agrees with the standard metric on S^2 on a neighborhood of the south pole and $\Psi_{\mathcal{T},i}(b_v) = 0$, by Corollary 4.7 for any $\varpi \in T_v F^H \mathcal{T}$ with $\|\varpi\|_v$ sufficiently small,

$$|c_i(\varpi) - c_i(v)| \leq C(b_v)|v|^{\frac{1}{p}}\|\varpi\|_v, \quad |r_i(\varpi) - r_i(v)| \leq C(b_v)|v|^{\frac{1}{p}}\|\varpi\|_v. \quad (4.16)$$

Let $\bar{b}(v) = (c(v), r(v)) \cdot \tilde{b}(v)$. Write

$$\bar{b}(v) = (S, M, H \cup \{\hat{0}\}; \bar{x}(v), (j(v), \bar{y}), \bar{u}_v).$$

If $\|\varpi\|_v$ is sufficiently small, define $\bar{\zeta}_\varpi \in \Gamma(\bar{u}_v)$ by

$$\exp_{b_v, \bar{u}_v} \bar{\zeta}_\varpi = u_\varpi, \quad \|\bar{\zeta}_\varpi\|_{b_v, C^0} < \text{inj } g_{V, b_v}.$$

Similarly, for $h \in H$ and $l \in M$, define $\bar{w}_h(\varpi) \in T_{\bar{x}_h(v)} \Sigma_{v, \iota_h(v)}$ and $\bar{w}_l(\varpi) \in T_{\bar{y}_l(v)} \Sigma_{v, j_l(v)}$ by

$$\begin{aligned} \exp_{g_v, \bar{x}_h(v)} \bar{w}_h(\varpi) &= \bar{x}_h(\varpi), \quad |\bar{w}_h(\varpi)| \equiv |\bar{w}_h(\varpi)|_{g_v} < \text{inj}(g_v, x_h(v)); \\ \exp_{g_v, \bar{y}_l(v)} \bar{w}_l(\varpi) &= \bar{y}_l(\varpi), \quad |\bar{w}_l(\varpi)| \equiv |\bar{w}_l(\varpi)|_{g_v} < \text{inj}(g_v, y_l(v)). \end{aligned}$$

Then by equation (4.16) and Corollary 4.7,

$$C''(b_v)^{-1} \|\varpi\|_v \leq \|\bar{\zeta}_\varpi\|_{v, p, 1} + \sum_{h \in H} |\bar{w}_h(\varpi)| + \sum_{l \in M} |\bar{w}_l(\varpi)| \leq C(b_v) \|\varpi\|_v. \quad (4.17)$$

It follows that the map

$$F^H \mathcal{T}_\delta \longrightarrow \mathcal{M}_{\mathcal{T}(H)}^{(0)}, \quad v \longrightarrow \bar{b}(v),$$

is a local embedding. By the same argument as in the proof of Lemma 4.8, we can conclude that this map is injective as long as $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ is sufficiently small. Since this map is $G_{\mathcal{T}(H)}$ -equivariant by construction, it follows that the induced map on the quotient, i.e. the map of Corollary 4.9, is injective.

Corollary 4.10 *If $S = S^2$, there exists $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}}; \mathbb{R}^+)$ such that the map*

$$\tilde{\gamma}_{\mathcal{T}}: F\mathcal{T}_\delta|_{\mathcal{M}_{\mathcal{T}}} \longrightarrow \bar{\mathcal{M}}_{\langle \mathcal{T} \rangle}$$

is injective. Furthermore, the restriction

$$\tilde{\gamma}_{\mathcal{T}}: F^\emptyset \mathcal{T}_\delta|_{\mathcal{M}_{\mathcal{T}}} \longrightarrow \mathcal{M}_{\langle \mathcal{T} \rangle}$$

is a differentiable embedding.

In order to adjust the gluing procedure in the presence of constraints, below we state the analogue of Corollary 4.7 for $\varpi \in \mathcal{K}_{b_v} \mathcal{T} \subset T_v F^\emptyset \mathcal{T}$. It is obtained in the same way as Corollary 4.7, except the analogue of Lemma 4.6 would make use of Lemma 3.8, instead of Lemma 3.6, and of (2b), instead of (2a), of Definitions 3.11 and 3.13. We also use (3) of Lemma 3.5.

Corollary 4.11 *There exist $\delta, C \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ such that for all $t \in [0; \delta(b_v)]$, $v \in F^{(\emptyset)} \mathcal{T}_\delta$, and $\varpi \in \mathcal{K}_{b_v} \mathcal{T}_{\delta(b_v)}$,*

$$C(b_v)^{-1} \|\varpi\| \leq \|\tilde{\zeta}_{\varpi, tv}\|_{v, p, 1} + \sum_{l \in M} |w_l(\varpi)|_{g_v} \leq C(b_v) \|\varpi\|.$$

Furthermore, $\|S'_\varpi \xi_{\varpi, tv} - \xi_{v, tv}\|_{v, C^0} \leq C(b_v)(t + |v|^{\frac{1}{p}}) \|\varpi\|$.

4.3 The Basic Gluing Map and the Space of Balanced Maps

Our next goal is to show that the gluing map of Subsection 3.6 is surjective in the appropriate sense. More precisely, if $\mathcal{T}$ is a regular bubble type, we show that the image of $\tilde{\gamma}_{\mathcal{T}}$ contains a neighborhood of $\mathcal{M}_{\mathcal{T}}$ in $\bar{\mathcal{M}}_{\langle \mathcal{T} \rangle}$. If $\mathcal{T}$ is a semiregular, we show that all elements in $\mathcal{M}_{\Sigma, tv, \lambda^*}$ that are close to any given compact subset of $\mathcal{M}_{\mathcal{T}}$ are in the image of the gluing map $\tilde{\gamma}_{\mathcal{T}, tv}$ if t is sufficiently small. The major difficulty in doing this is the following. If $v \in F\mathcal{T}$, a small change in the singular points of b_v may lead to a very large change in the map u_v . This is precisely the reason we used the norm $\|\varpi\|_v$ on $T_v F^H \mathcal{T}$ instead of just $\|\varpi\|$ in Subsection 3.4. In order to deal with this issue, we need Corollary 4.13, which is proved in this subsection. We continue to assume that $\mathcal{T}$ is a simple bubble type.

Recall that $\mathcal{H}_{\mathcal{T}}$ is the set of tuples $b = (S, M, I; x, (j, y), u)$ such that $u_{\iota_h}(x_h) = u_h(\infty)$ for all $h \in \hat{I}$ and $\bar{\partial}u_i = 0$ for all $i \in I$. Furthermore, $\mathcal{M}_{\mathcal{T}}^{(0)}$ is the subset of $\mathcal{H}_{\mathcal{T}}$ consisting of the tuples b such that $\Psi_{\mathcal{T}, h}(b) = 0$ for all $h \in \hat{I}$. It is convenient to make the following definitions. If H is a subset of $\hat{I}$ and $\epsilon \geq 0$, let

$$\begin{aligned} \mathcal{M}_{\mathcal{T}, \epsilon}^{(H)} = \{ & b = (S, M, I; x, (j, y), u) : \bar{\partial}u_i = 0 \ \forall i \in I; \ d_V(u_{\iota_h}(x_h), u_h(\infty)) \leq \epsilon \ \forall h \in \hat{I}; \\ & |\Psi_{\mathcal{T}, h}(b)| \leq \epsilon \ \forall h \in \hat{I} - H \}. \end{aligned}$$

Lemma 4.12 *There exist $\delta, C \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R}^+)$ with the following property. Suppose $b^* \in \mathcal{M}_T^{(0)}$, $\epsilon < \delta(b^*)$, $b \in \mathcal{M}_{T,\epsilon}^{(H)}$ is such that $d(b^*, b) \leq \delta(b^*)$, and $v = (b, v_{\hat{I}}) \in F_b^{(H)} \mathcal{T}_{\delta(b^*)}$. Then there exist*

$$\tilde{b} \in \mathcal{M}_{T,\epsilon^2}^{(H)} \quad \text{and} \quad \tilde{v} = (\tilde{b}, \tilde{v}_{\hat{I}}) \in F_{\tilde{b}}^{(H)} \mathcal{T}$$

such that

- (1) $d(b, \tilde{b}) \leq C(b^*)\epsilon$ and $|\tilde{v}_h - v_h|_b \leq C(b^*)\epsilon|v_h|_b$ for all $h \in \hat{I}$;
- (2) if $q_v(z) \in \Sigma_{T,i}$, $|r_{b,h}(q_v(z))| \geq 2|v_h|^{\frac{1}{2}}$ for all $h \in \hat{I} - H$ such that $\iota_h = i$ and $|q_S^{-1}(q_v(z))| \geq 2|v_i|^{\frac{1}{2}}$ if $i \in \hat{I} - H$, then $d_b(q_v(z), q_{\tilde{v}}(z)) \leq \delta(b^*)\epsilon$.

Proof: (1) Let $b = (S, M, I; x, (j, y), u)$. If δ is sufficiently small, by Proposition 3.3, we can choose $\xi_i \in \Gamma(u_i)$ such that $\|\xi_i\|_{g_{b,i}, C^1} \leq C(b^*)\epsilon$ and

$$b' \equiv (S, M, I; x, (j, y), u') \in \mathcal{H}_T,$$

where $u'_i = \exp_{u_i} \xi_i$. The C^1 -bound on ξ_i and the assumption $b \in \mathcal{M}_{T,\epsilon}^{(H)}$ imply that $|\Psi_{T,h}(b')| \leq C'(b^*)\epsilon$ for all $h \in \hat{I} - H$.

(2) We now define $\tilde{b}' \equiv (S, M, I; \tilde{x}', (j, \tilde{y}'), \tilde{u}')$ and $\tilde{v}' = (\tilde{b}', \tilde{v}'_{\hat{I}})$ as follows. Suppose $i^* \in I$ and for all $i \in \hat{I}$ with $i > i^*$, $h \in \hat{I}$ with $\iota_h = i$, and $l \in M$ with $j_l = i$, we have constructed

- (i) $(c_i, r_i) \in \mathbb{C} \times \mathbb{R}$ such that $|(c_i, r_i)| \leq C(b^*)\epsilon$;
- (ii) $\tilde{x}'_h, \tilde{y}'_l \in \Sigma_{T,i}$ such that $|r_{b,h}(\tilde{x}'_h)| \leq C(b^*)\epsilon$ and $|\phi_{y_l} \tilde{y}'_l| \leq C(b^*)\epsilon$;
- (iii) $\tilde{v}'_h \in \mathbb{C}$ such that $|\tilde{v}'_h - v_h| \leq C(b^*)\epsilon|v_h|$;
- (iv) if $x_i \in S^2$, $\bar{x}_i \in S^2$, such that $|r_{b,i}(\bar{x}_i)| \leq C(b^*)\epsilon|v_i|$;
- (v) if $x_i \in S^2$, $\bar{v}_i \in \mathbb{C}$ such that $|\bar{v}_i - v_i| \leq C(b^*)\epsilon|v_i|$,

such that

- (I1) if $i \notin H$, $\Psi_{T,i}(\tilde{b}') = 0$ where $\tilde{u}'_i = (c_i, r_i) \cdot u'_i$;
- (I2) if $\Sigma_{T,\iota_i} = S^2$, $z \in \Sigma_{T,\iota_i}$, and $|\phi_{\bar{x}_h} q_{i,(x_i, v_i)}(z)| \leq \frac{2}{3}|\bar{v}_h|^{\frac{1}{2}}$ for some $h \in \hat{I} - H$, then

$$\left| \phi_{\tilde{x}'_h} q_{i,(\bar{x}_i, \bar{v}_i)}(z) \right| \leq |\tilde{v}'_h|^{\frac{1}{2}} \quad \text{and} \quad q_{h,(\tilde{x}'_h, \tilde{v}'_h)}(q_{i,(\bar{x}_i, \bar{v}_i)}(z)) = q_{h,(\bar{x}_h, \bar{v}_h)}(q_{i,(x_i, v_i)}(z)),$$

where $q_{i,(x_i, v_i)}$, etc., are the maps defined in Section 2.2.

Note that while we have not defined $\tilde{b}'$ completely yet, (I1) is still a well-defined statement. The function $\Psi_{T,i}$ depends only the i th bubble component of $\tilde{b}'$, which has already been constructed by the induction assumptions.

If $i^* \in H$, we take $c_{i^*} = 0$ and $r_{i^*} = 0$. If $i^* \in \hat{I} - H$, let $(c_{i^*}, r_{i^*}) \in \mathbb{C} \times \mathbb{R}$ be given by

$$\begin{aligned} \tilde{\Psi}((c_{i^*}, 0)u'_{i^*}) + \sum_{\iota_h = i^*} d_h(\mathcal{T})(\bar{x}_h + c_{i^*}) + \sum_{j_l = i^*} (y_l + c_{i^*}) &= 0; \\ \Psi^{(3)}((c_{i^*}, r_{i^*})u'_{i^*}) + \sum_{\iota_h = i^*} d_h(\mathcal{T})\beta((1 + r_{i^*})|\bar{x}_h + c_{i^*}|) + \sum_{j_l = i^*} \beta((1 + r_{i^*})|y_l + c_{i^*}|) &= \frac{1}{2}. \end{aligned}$$

If ϵ is sufficiently small, by the proof of Lemma 3.3 such $(c_{i^*}, r_{i^*}) \in \mathbb{C} \times \mathbb{R}$ exists and satisfies $|c_{i^*}|, |r_{i^*}| \leq C(b^*)\epsilon$. For all $h \in \hat{I}$ with $\iota_h = i^*$ and $l \in M$ with $j_l = i^*$, put

$$\begin{aligned} \tilde{x}'_h &= (1 + r_{i^*})(\bar{x}_h + c_{i^*}), \quad \tilde{v}'_h = (1 + r_{i^*})\bar{v}_h, \quad \tilde{y}'_l = (1 + r_{i^*})(y_l + c_{i^*}); \\ \bar{x}_{i^*} &= x_{i^*} - c_{i^*}v_{i^*}, \quad \bar{v}_{i^*} = (1 + r_{i^*})^{-1}v_{i^*} \quad \text{if} \quad x_{i^*} \in S^2. \end{aligned}$$

Continuing in this way, for all $i \in \hat{I}$, $h \in \hat{I}$ with $\iota_h = i$, and $l \in M$ with $j_l = i$, we obtain elements (i)-(v) satisfying (I1),(I2). Let $\tilde{u}'_{\hat{0}} = u'_{\hat{0}}$. If $l \in M$ and $j_l = \hat{0}$, take $\tilde{y}'_l = y_l$.

(3) If $S = S^2$, let $(\tilde{x}'_h, \tilde{v}'_h) = (\bar{x}_h, \bar{v}_h)$ if $\iota_h = 0$, $\tilde{b} = \tilde{b}'$, and $\tilde{v} = \tilde{v}'$. By the inductive construction, $\tilde{b}$ and $\tilde{v}$ satisfy the requirements of the lemma. In fact, $\tilde{b} \in \mathcal{M}_{\mathcal{T},0}^{(H)}$. If $S = \Sigma$, we could extend the above construction to the principal component Σ as we did for $S = S^2$ if $q_{\tilde{v}'}$ were defined using the metric $g_{b,\hat{0}}$ on Σ , which may differ slightly from $g_{\tilde{b},\hat{0}}$. This problem is fixed in below.

(4) If $l \in M$ and $j_l = \hat{0}$, we take $\tilde{y}_l = y_l$ as before. For all $h \in \hat{I}$ with $\iota_h = \hat{0}$, let $\tilde{x}_h \in \Sigma$, $\tilde{v}_h \in T_{\tilde{x}_h}\Sigma$, and $\Theta_h : B_{2|v_h|_b}^{-\frac{1}{2}}(0; \mathbb{C}) \longrightarrow \mathbb{C}$ be such that

$$(\Sigma_{\hat{0}}1) \quad d_b(x_h, \tilde{x}_h) \leq C(b^*)\epsilon|v_h|, \quad ||\tilde{v}_h|_{\tilde{b}} - |v_h|_b| \leq C(b^*)\epsilon|v_h|_b;$$

$$(\Sigma_{\hat{0}}2) \quad \text{for all } z \in B_b(x_h, 2|v_h|_b^{\frac{1}{2}}),$$

$$\frac{\phi_{\tilde{b},h}z}{\tilde{v}_h} = (1 + r_h) \left\{ c_h + \frac{\phi_{b,h}z}{v_h} + \Theta_h \left(\frac{\phi_{b,h}z}{v_h} \right) \right\};$$

$$(\Sigma_{\hat{0}}3) \quad \Theta_h \text{ is holomorphic, } \Theta_h(0) = 0, \Theta'_h(0) = 0, \text{ and } \|\Theta''_h\|_{C^0} \leq C(b^*)|v|^2\epsilon.$$

Note that even though we have not defined $\tilde{b}$ completely yet, $(\Sigma_{\hat{0}}1)$ and $(\Sigma_{\hat{0}}2)$ are still well-defined statements, since the metric $g_{\tilde{b},\hat{0}}$ on Σ depends only on the singular points $\{\tilde{x}_h : \iota_h = \hat{0}\}$ on Σ . Existence of such $\tilde{x}_h$, $\tilde{v}_h$, and Θ_h follows from Corollary 5.5, provided δ is sufficiently small.

If $\iota_i = \hat{0}$ and $j_l = i$, let $(i, \tilde{y}_l) = q_{\tilde{v},i}q_{v,i}^{-1}(i, y_l)$. The map $q_{\tilde{v},i}$ is well-defined even though $\tilde{v}$ has not been defined completely yet. By $(\Sigma_{\hat{0}}2)$,

$$\tilde{y}_l = q_{\tilde{v},i}q_{v,i}^{-1}(y_l) = \frac{\phi_{\tilde{b},\tilde{x}_i}\phi_{b,x_i}^{-1}(y_lv_i)}{\tilde{v}_i} = (1 + r_i) \{c_i + y_l + \Theta_i(y_l)\}. \quad (4.18)$$

Since $\tilde{y}'_l = (1 + r_i)(y_l + c_i)$, $|\tilde{y}_l - \tilde{y}'_l| \leq C(b^*)|v|^2\epsilon$ by $(\Sigma_{\hat{0}}3)$.

Suppose $h \in \hat{I}$, $\iota_h \in \hat{I}$, and for every $i \in \hat{I}$ with $i < h$ and $j \in M$ with $j_l = i$, we have defined

$$\tilde{x}_i \in \Sigma_{\mathcal{T},\iota_i}, \quad \tilde{y}_l \in \Sigma_{\mathcal{T},i}, \quad \tilde{v}_i \in \begin{cases} T_{\tilde{x}_i}\Sigma, & \text{if } \iota_i = 0; \\ \mathbb{C}, & \text{if } \iota_i > 0; \end{cases} \quad \tilde{c}_i \in \mathbb{C}, \quad \Theta_i : B_{2|v_i|_b}^{-\frac{1}{2}}(0; \mathbb{C}) \longrightarrow \mathbb{C}$$

such that

$$(\Sigma1) \quad |\phi_{\tilde{b}',i}\tilde{x}_i|_{\tilde{b}'} \leq C(b^*)|v|^2\epsilon \text{ if } \iota_i > 0 \text{ and } |\phi_{\tilde{b}',\tilde{y}'_l}\tilde{y}_l|_{\tilde{b}'} \leq C(b^*)|v|^2\epsilon;$$

$$(\Sigma2) \quad ||\tilde{v}_i|_{\tilde{b}} - |v_i|_b| \leq C(b^*)\epsilon|v_i|_b;$$

$$(\Sigma3) \quad |\tilde{c}_i - c_i| \leq C(b^*)|v|^2\epsilon;$$

$$(\Sigma4) \quad \text{for all } z \in \Sigma \text{ such that } r_{b,i}q_{v,\iota_i}(z) \leq 2|v_i|_b^{\frac{1}{2}},$$

$$\frac{\phi_{\tilde{b},i}q_{\tilde{v},\iota_i}z}{\tilde{v}_i} = (1 + r_i) \left\{ \tilde{c}_i + \frac{\phi_{b,i}q_{v,\iota_i}z}{v_i} + \Theta_i \left(\frac{\phi_{b,i}q_{v,\iota_i}z}{v_i} \right) \right\}$$

$$(\Sigma5) \quad \Theta_i \text{ is holomorphic, } \Theta_i(0) = 0, \Theta'_i(0) = 0, \text{ and } \|\Theta''_i\|_{C^0} \leq C(b^*)|v|^2\epsilon.$$

If $h \in H$, we take $\tilde{x}_h = \tilde{x}'_h$, $\tilde{v}_h = \tilde{v}'_h = 0$, $\tilde{y}_l = \tilde{y}'_l$ if $j_l = h$, $\tilde{c}_h = c_h = 0$, and $\Phi_h(z) = 0$. If $h \notin H$, let

$$(\iota_h, \tilde{x}_h) = q_{\tilde{v},\iota_h}q_{v,\iota_h}^{-1}(h, \bar{x}_h).$$

By an argument similar to (4.18), from $(\Sigma 4)$ we obtain

$$\tilde{x}_h = (1 + r_{\iota_h}) \{ \tilde{c}_{\iota_h} + \bar{x}_h + \Theta_{\iota_h}(\bar{x}_h) \}. \quad (4.19)$$

Since $\tilde{x}'_h = (1 + r_{\iota_h})(\bar{x}_h + c_{\iota_h})$, (4.19), $(\Sigma 3)$, and $(\Sigma 5)$ imply the first part of $(\Sigma 1)$ with $i = h$. Furthermore, by assumption $(\Sigma 4)$,

$$\begin{aligned} \phi_{\tilde{b},h} q_{\tilde{v},\iota_h}(z) &= q_{\tilde{v},\iota_h}(z) - \tilde{x}_h = \frac{\phi_{\tilde{b},\iota_h} q_{\tilde{v},\iota_h}(z)}{\tilde{v}_{\iota_h}} - \tilde{x}_h \\ &= (1 + r_{\iota_h}) \left\{ \left(\frac{\phi_{b,\iota_h} q_{v,\iota_h}(z)}{v_{\iota_h}} - \bar{x}_h \right) + \left(\Theta_{\iota_h} \left(\frac{\phi_{b,\iota_h} q_{v,\iota_h}(z)}{v_{\iota_h}} \right) - \Theta_{\iota_h}(\bar{x}_h) \right) \right\}. \end{aligned} \quad (4.20)$$

Since Θ_{ι_h} is holomorphic, and

$$\frac{\phi_{b,\iota_h} q_{v,\iota_h}(z)}{v_{\iota_h}} - \bar{x}_h = \phi_{b,h} q_{v,\iota_h}(z) + c_h v_h,$$

we can rewrite (4.20) as

$$\phi_{\tilde{b},h} q_{\tilde{v},\iota_h}(z) = (1 + r_{\iota_h}) (1 + a_h) v_h \left\{ \tilde{c}_h + \frac{\phi_{b,h} q_{v,\iota_h}(z)}{v_h} + \Theta_h \left(\frac{\phi_{b,h} q_{v,\iota_h}(z)}{v_h} \right) \right\}, \quad (4.21)$$

where the complex numbers $a_h, \tilde{c}_h \in \mathbb{C}$ and the holomorphic function $\Theta_h: B_{2|v_h|^{-\frac{1}{2}}}(0, \mathbb{C}) \rightarrow \mathbb{C}$ are given by

$$a_h = \frac{d}{dz} \Theta_{\iota_h}(z) \Big|_{z=x_h}, \quad (1 + a_h) \tilde{c}_h = c_h + \frac{\Theta_{\iota_h}(x_h) - \Theta_{\iota_h}(x_h - c_h v_h)}{v_h}, \quad (4.22)$$

$$\Theta_h(z) = \frac{\Theta_{\iota_h}(v_h z + x_h) - v_h z \Theta'_{\iota_h}(x_h) - \Theta_{\iota_h}(x_h)}{(1 + a_h) v_h}. \quad (4.23)$$

By (4.23), $\Theta_h(0) = 0$ and $\Theta'_h(0) = 0$. By our assumptions on Θ_{ι_h} and (4.22), (4.23),

$$|a_h| \leq C(b^*) |v|^2 \epsilon |x_{\iota_h}| \leq C'(b^*) |v|^2 \epsilon; \quad (4.24)$$

$$|\tilde{c}_h - c_h| \leq C(b^*) (\epsilon |a_h| + |v_h|^{-1} |v|^2 \epsilon |c_h v_h|) \leq C'(b^*) |v|^2 \epsilon, \quad (4.25)$$

$$\|\Theta''_h\|_{C^0} \leq C(b^*) |v_h|^{-1} |v|^2 \epsilon |v_h|^2 \leq C'(b^*) |v|^2 \epsilon. \quad (4.26)$$

We now take

$$\tilde{v}_h = (1 + a_h) \tilde{v}'_h = (1 + a_h) (1 + r_{\iota_h}) (1 + r_h)^{-1} v_h.$$

It follows from (4.24)-(4.26) that the induction hypotheses $(\Sigma 2)$ -($\Sigma 5$) with $i = h$ are satisfied. If $j_l = h$, let $(h, \tilde{y}_l) = q_{\tilde{v},h}^{-1}(h, y_l)$. By the same argument as in the case $\iota_h = \hat{0}$ above, $(\Sigma 3)$ -($\Sigma 5$) of the $i = \iota_h$ case imply that the second part of $(\Sigma 1)$ with $i = h$ is satisfied. Continuing in this way, we obtain tuples

$$\tilde{b} = (\Sigma, M, I; \tilde{x}, (j, \tilde{y}), \tilde{u}'), \quad \tilde{c} = c_{\tilde{f}}, \quad \tilde{v} = (\tilde{b}, \tilde{v}_{\tilde{f}}),$$

satisfying $(\Sigma 1)$ -($\Sigma 5$). Since $\tilde{b}' \in \mathcal{M}_{T,0}^{(H)}$, by $(\Sigma 1)$ $\tilde{b} \in \mathcal{M}_{T,\epsilon}^{(H)}$ if δ is sufficiently small. Finally, $(\Sigma 1)$ -($\Sigma 5$) along with (I1) and (I2) show that $\tilde{b}$ and $\tilde{v}$ satisfy the two requirements of the lemma.

Corollary 4.13 *If $\mathcal{T}$ is a simple bubble type, there exist $\delta, C \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R}^+)$ with the following property. Suppose $b^* \in \mathcal{M}_{\mathcal{T}}^{(0)}$, $\epsilon < \delta(b^*)$, $b \in \mathcal{M}_{\mathcal{T}, \epsilon}^{(H)}$ is such that $d(b^*, b) \leq \delta(b^*)$, and $v = (b, v_{\hat{I}}) \in F_b^{(H)} \mathcal{T}_{\delta(b^*)}$. Then there exist $\tilde{b} \in \mathcal{M}_{\mathcal{T}, 0}^{(H)}$ and $\tilde{v} = (\tilde{b}, \tilde{v}_{\hat{I}}) \in F_{\tilde{b}}^{(H)} \mathcal{T}$ such that*

- (1) $d(b, \tilde{b}) \leq C(b^*)\epsilon$ and $|\tilde{v}_h - v_h| \leq C(b^*)\epsilon|v_h|$ for all $h \in \hat{I}$;
- (2) if $q_v(z) \in \Sigma_{\mathcal{T}, i}$, $|r_{b, h}(q_v(z))| \geq 3|v_h|^{\frac{1}{2}}$ for all $h \in \hat{I} - H$ such that $\iota_h = i$ and $|q_S^{-1}(q_v(z))| \geq 3|v_i|^{\frac{1}{2}}$ if $i \in \hat{I} - H$, then $d_b(q_v(z), q_{\tilde{v}}(z)) \leq \epsilon$.

Proof: If $S = S^2$, the tuples $\tilde{b}$ and $\tilde{v}$ constructed in the first half of the proof of Lemma 4.12 satisfy the requirements of the corollary. In fact, $d_b(q_v(z), q_{\tilde{v}}(z)) = 0$ if z is as in (2) above. If $S = \Sigma$, let

$$\tilde{\epsilon} = \epsilon^2 \prod_{h \in [I] - H} |v_h|_b^2 > 0.$$

If $C(b^*)\delta(b^*)$ is sufficiently small, by repeated applications of Lemma 4.12, we can replace the tuples b and v by $b' \in \mathcal{M}_{\mathcal{T}, \tilde{\epsilon}}^{(H)}$ and $v' = (b', v'_{\hat{I}}) \in F^{(H)} \mathcal{T}$ such that

- (1) $d(b, b') \leq C'(b^*)\epsilon$ and $|v'_h - v_h| \leq C'(b^*)\epsilon|v_h|_b$ for all $h \in \hat{I}$;
- (2) if $q_v(z) \in \Sigma_{\mathcal{T}, i}$, $|r_{b, h}(q_v(z))| \geq \frac{5}{2}|v_h|^{\frac{1}{2}}$ for all $h \in \hat{I} - H$ such that $\iota_h = i$ and $|q_S^{-1}(q_v(z))| \geq \frac{5}{2}|v_i|^{\frac{1}{2}}$ if $i \in \hat{I} - H$, then $d_b(q_v(z), q_{v'}(z)) \leq 2\delta(b)\epsilon$.

Applying the construction of the first half of the proof of Lemma 4.12 to the tuples b' and v' , we obtain tuples $\tilde{b} \in \mathcal{M}_{\mathcal{T}}^{(0)}$ and $\tilde{v} = (\tilde{b}, \tilde{v}_{\hat{I}}) \in F_{\tilde{b}}^{(H)} \mathcal{T}$ such that

$$d(b', \tilde{b}) \leq C(b^*)\tilde{\epsilon} \quad \text{and} \quad |\tilde{v}_h - v'_h| \leq C(b^*)\tilde{\epsilon}|v'_h|_{b'} \quad \forall h \in \hat{I}.$$

Then if z is as in the requirement (2) of the corollary,

$$d_{b'}(q_{v'}(z), q_{\tilde{v}}(z)) \leq C(b^*)\tilde{\epsilon} \left(\prod_{h \in \hat{I} - H} |v'_h|_{b'} \right)^{-1} \leq \epsilon^2$$

if δ is sufficiently small. Thus, the tuples $\tilde{b}$ and $\tilde{v}$ satisfy both requirements of the corollary.

4.4 Gromov Convergence and the L^p -norm of the Differential

Let $b_k = (S, M, I; x, (j, y_k), u_k)$ be a sequence of smooth maps converging to

$$b^* = (S, M, I^*; x^*, (j^*, y^*), u^*) \in \mathcal{M}_{\mathcal{T}^*}^{(0)}$$

with respect to the Gromov topology such that $\bar{\partial}u_{k, \hat{0}} = t_k \nu$ with $t_k \rightarrow 0$ and $\bar{\partial}u_{k, h} = 0$ if $h \in \hat{I}$. We assume that $\mathcal{T}^*$ is a simple bubble type. In the next subsection, it is proved that b_k lies in the image of the gluing map $\tilde{\gamma}_{\mathcal{T}, t_k \nu}$ for some k . In this subsection, we show the differentials of $du_{k, i}$ satisfy a certain condition which holds for all bubble maps in the image of $\tilde{\gamma}_{\mathcal{T}, t_k \nu}$.

By definition of convergence, for all k sufficiently we can choose

(a) curves $\mathcal{C}_k = (S, M, I^*; x'_k, (j^*, y^*))$ with $\lim_{k \rightarrow \infty} x'_{k, h} \rightarrow x_h^*$ for all $h \in \hat{I}^*$, and

(b) vectors $(v_k)_{\hat{I}^*} \in F_{\mathcal{C}_k}^{(0)}$ with $16|v_k|_{g_b} \leq r_{\mathcal{C}_k} g_b$,

such that $\lim_{k \rightarrow \infty} |v_k| = 0$, $\mathcal{C}(v_k) = (S, M, I; x_k, (j_k, y(v_k)))$, and

$$\lim_{k \rightarrow \infty} \sup_{z \in \Sigma_{\mathcal{C}(v_k)}} d_V(u_{b^*}(q_{v_k}(z)), u_{b_k}(z)) = 0, \quad \lim_{k \rightarrow \infty} q_{v_k}(j_{k, l}, y_{k, l}) = (j_l^*, y_l^*) \quad \forall l \in M,$$

where $v_k = (\mathcal{C}_k, (v_k)_{\hat{f}^*})$ and g_b denotes the standard metric on $\Sigma_{\mathcal{C}_k}$ if $S = S^2$. Let

$$\phi_{k,h} = \begin{cases} \phi_{x'_{k,h}}, & \text{if } x'_{k,h} \in S^2; \\ \phi_{g_{b,\hat{0}},x'_{k,h}}, & \text{if } x'_{k,h} \in \Sigma; \end{cases} \quad r_{k,h} = \begin{cases} r_{x'_{k,h}}, & \text{if } x'_{k,h} \in S^2; \\ r_{g_{b,\hat{0}},x'_{k,h}}, & \text{if } x'_{k,h} \in \Sigma. \end{cases}$$

Let g_{v_k} be the metric on $\Sigma_{b_k} = \Sigma_{v_k}$ defined as in Section 3.3, using the metric $g_{b,\hat{0}}$ on Σ if $S = \Sigma$.

For any element in the image $\tilde{\gamma}_{T,t\nu}$ that lies near b^* , the modified (L^p, g_{v_k}) -norm of $d\tilde{u}_v$ is bounded above by a constant dependent only on b^* . Furthermore, as $v \rightarrow 0$ and the size of the necks is reduced, the modified (L^p, g_v) -norm of $d\tilde{u}_v$ on such necks tends to zero. The modified (L^p, g_v) -norm is bounded above by the usual (L^{2p}, g_v) -norm times some constant dependent only on b^* . In this subsection, we show that the (L^{2p}, g_{v_k}) -norm of du_{b_k} is uniformly bounded and tends to zero on the “necks.” Instead of using our usual cutoff function β , we will use the family of cutoff functions provided by the following lemma. The proof can be found in [MS, p166]. The statement below is somewhat sharper than in [MS], but the proof in [MS] suffices.

Lemma 4.14 *For every $\epsilon > 0$, there exists a smooth function $\tilde{\beta}_\epsilon: \mathbb{R} \rightarrow [0, 1]$ such that*

$$\int_{\mathbb{C}} |\tilde{\beta}'_\epsilon(r)|^2 r dr d\theta \leq 8\epsilon, \quad \text{and} \quad \tilde{\beta}_\epsilon(r) = \begin{cases} 1, & \text{if } r \geq 1; \\ 0, & \text{if } r \leq e^{-1/\epsilon}. \end{cases}$$

Similarly to the above, given $r > 0$, we denote by $\tilde{\beta}_{\epsilon,r}$ the cutoff function defined by $\tilde{\beta}_{\epsilon,r}(t) = \tilde{\beta}_\epsilon(r^{-\frac{1}{2}}t)$.

We now define nearly holomorphic maps $f_{k,i} \in C^\infty(\Sigma_{\mathcal{C}_k,i}; V)$. In order to simplify computations, we fix a finite family of J -invariant metrics on V such that for some fixed $\varepsilon > 0$ and for every $q \in V$ there exists a metric $g_{V,q}$ in this family such that $(B_{g_{V,q}}(q, \varepsilon), J, g_{V,q})$ is isomorphic to a ball in $\mathbb{C}^n$. Since V is compact and the family of metrics is finite, all estimates below that depend on a particular metric $g_{V,q}$ will involve bounds dependent only on V . We denote by $\exp_q$ the exponential map of (the Levi-Civita connection of) the metric $g_{V,q}$ and the $g_{V,q}$ -geodesic about q or radius ϵ by $B_q(\epsilon)$. If $\delta > 0$ and $h \in I^* - I$, let

$$\begin{aligned} B_{h,k}^+(\delta) &= \{(\iota_h^*, z) \in \Sigma_{\mathcal{C}_k, \iota_h^*} : r_{k,h}(\iota_h^*, z) \leq \delta\}, \\ B_{h,k}^-(\delta) &= \{(h, z) \in \Sigma_{\mathcal{C}_k, h} : |q_S^{-1}(z)| \leq \delta\}. \end{aligned} \tag{4.27}$$

Choose a sequence $\epsilon_k \in \mathbb{R}^+$ converging to zero. Let $r_k = \left(2 \sum_{i \in I^*} \|du_i^*\|_{b^*, C^2}\right)^{-1} \epsilon_k$. By taking a subsequence if necessary, it can be assumed that

$$|t_k| \leq \epsilon_k, \quad d_V(u_{b^*}(q_{v_k}(z)), u_{b_k}(z)) \leq \epsilon_k, \quad r_{b^*,h}(\iota_h^*, x'_{k,h}) \leq r_k, \quad e^{\frac{2p}{\epsilon_k}} |v_{k,h}|_{b^*}^{\frac{1}{2}} \leq r_k. \tag{4.28}$$

Let $q_h = u_h^*(\infty)$ and

$$\tilde{A}_{h,k}^\pm = B_{h,k}^\pm \left(|v_{k,h}|_{b^*}^{\frac{1}{2}} \right) - B_{h,k}^\pm \left(e^{-\frac{1}{\epsilon_k}} |v_{k,h}|_{b^*}^{\frac{1}{2}} \right).$$

By (4.28), $u_{b_k} \left(q_{v_k}^{-1} \left(B_{h,k}^\pm \left(e^{\frac{1}{\epsilon_k}} |v_{k,h}|_{b^*}^{\frac{1}{2}} \right) \right) \right) \subset B_{q_h}(C(b^*)\epsilon_k)$. Thus, we can define $\xi_{k,h}^\pm \in C^\infty(\tilde{A}_{h,k}^\pm; T_{q_h} V)$ by

$$\begin{aligned} \exp_{q_h, q_h} \xi_{k,h}^+(z) &= u_{b_k} \left(q_{v_k, \iota_h^*}^{-1}(\iota_h^*, z) \right), & |\xi_{k,h}^+(z)|_{g_{V, q_h}} &< \varepsilon; \\ \exp_{q_h, q_h} \xi_{k,h}^-(z) &= u_{b_k} \left(q_{v_k, \iota_h^*}^{-1}(\iota_h^*, \phi_{k,h}^{-1}(z v_{k,h})) \right), & |\xi_{k,h}^-(z)|_{g_{V, q_h}} &< \varepsilon, \end{aligned} \tag{4.29}$$

provided k is sufficiently large (depending on b^*). Let $\bar{\xi}_{k,h}^\pm \in T_{q_h} V$ be given by

$$\bar{\xi}_{k,h}^\pm = \frac{1}{\text{Area}(\tilde{A}_{k,h}^\pm)} \int_{\tilde{A}_{k,h}^\pm} \xi_{k,h}^\pm, \quad (4.30)$$

where the area and the integral are computed using the metric g_{b^*,ι_h^*} on Σ_{b^*,ι_h^*} and $g_{b^*,h}$ on $\Sigma_{b^*,h}$. Define $f_{k,i} \in C^\infty(\Sigma_{b^*,i}; V)$ by

$$f_{k,i}(z) = \begin{cases} \exp_{q_h, q_h} \left\{ \bar{\xi}_{k,h}^+ + \tilde{\beta}_{\epsilon_k, |v_{k,h}|_{b^*}}(r_{k,h}(z)) (\xi_{k,h}^+(z) - \bar{\xi}_{k,h}^+) \right\}, & \text{if } r_{k,h}(z) \leq |v_{k,h}|_{b^*}^{\frac{1}{2}}; \\ \exp_{q_i, q_i} \left\{ \bar{\xi}_{k,i}^- + \tilde{\beta}_{\epsilon_k, |v_{k,i}|_{b^*}}(|q_S^{-1}(z)|) (\xi_{k,i}^-(z) - \bar{\xi}_{k,i}^-) \right\}, & \text{if } i \in I^* - I \text{ \& } |q_S^{-1}(z)| \leq |v_{k,i}|_{b^*}^{\frac{1}{2}}; \\ u_{b_k}(q_{v_k}^{-1}(i, z)), & \text{otherwise.} \end{cases}$$

Let $\zeta'_{k,i} \in \Gamma(u_i^*)$ be given by

$$\exp_{b^*, u_i^*} \zeta'_{k,i} = f_{k,i}, \quad \|\zeta'_{k,i}\|_{b^*, C^0} < \text{inj } g_{V, b^*}.$$

Lemma 4.15 *There exists $C > 0$ such that for all k sufficiently large and $i \in I^*$,*

$$\|\zeta'_{k,i}\|_{b^*, C^0} \leq C\epsilon_k, \quad \|\bar{\partial} f_{k,i}\|_{g_{b^*,i}, 2p} \leq C\epsilon_k^{\frac{1}{2p}} (\|df_{k,i}\|_{g_{b^*,i}, 2p} + 1).$$

Proof: The first statement is clear from (4.28) and the construction of $f_{k,i}$ above. Suppose $z \in \Sigma_{b^*,i}$. If $z \notin B_{h,k}^+(|v_{k,h}|_{b^*}^{\frac{1}{2}})$ for all $h \in I^* - I$ and $z \notin B_{i,k}^-(|v_{k,i}|_{b^*}^{\frac{1}{2}})$ if $i \in I^* - I$, then

$$|\bar{\partial} f_{k,i}|_{g_{b^*,i}, z} \leq C_\nu t_k \leq C_\nu \epsilon_k. \quad (4.31)$$

Suppose $z \in \tilde{A}_{h,k}^+$ with $h \in I^* - I$. Since the metric $g_{\mathbb{P}^n, q_h}$ is flat near q_h ,

$$\bar{\partial} f_{k,i}|_z = d \exp_{q_h, q_h} \bar{\partial} \left\{ \bar{\xi}_{k,h}^+ + \tilde{\beta}_{\epsilon_k, |v_{k,h}|_{b^*}}(r_{k,h}(\cdot)) (\xi_{k,h}^+ - \bar{\xi}_{k,h}^+) \right\}_z. \quad (4.32)$$

It follows from (4.32) that

$$|\bar{\partial} f_{k,i}|_{g_{b^*,i}, z} \leq C \left(|v_{k,h}|_{b^*}^{-\frac{1}{2}} |d\tilde{\beta}_{\epsilon_k}|_{|v_{k,h}|_{b^*}^{-\frac{1}{2}} r_{k,h}(z)} |\xi_{k,h}^+ - \bar{\xi}_{k,h}^+|_z + |\bar{\partial} \xi_{k,h}^+|_z \right) \quad (4.33)$$

By Poincare Lemma, see Lemma ?? in [Z1] applied with $r = |v_{h,k}|_{b^*}^{-\frac{1}{2}}$ and $2p$ instead of p ,

$$\begin{aligned} \left\| |v_{k,h}|_{b^*}^{-\frac{1}{2}} |d\tilde{\beta}_{\epsilon_k}|_{|v_{k,h}|_{b^*}^{-\frac{1}{2}} r_{k,h}(\cdot)} |\xi_{k,h}^+ - \bar{\xi}_{k,h}^+| \right\|_{g_{b^*,i}, L^{2p}(\tilde{A}_{h,k}^+)} &\leq C |v_{k,h}|_{b^*}^{-\frac{p-1}{2p}} \|d\tilde{\beta}_{\epsilon_k}\|_2^{\frac{1}{p}} \|\xi_{k,h}^+ - \bar{\xi}_{k,h}^+\|_{b^*, C^0} \\ &\leq C' \epsilon_k^{\frac{1}{2p}} \|d\xi_{k,h}^+\|_{g_{b^*,i}, L^{2p}(\tilde{A}_{h,k}^+)} \leq C' \epsilon_k^{\frac{1}{2p}} \|df_{k,h}\|_{g_{b^*,i}, 2p}. \end{aligned}$$

The last two equations give

$$\|\bar{\partial} f_{k,i}\|_{g_{b^*,i}, L^{2p}(\tilde{A}_{h,k}^+)} \leq C(\epsilon_k^{\frac{1}{2p}} \|df_{k,h}^+\|_{g_{b^*,i}, 2p} + \epsilon_k). \quad (4.34)$$

The same estimate applies to $\|\bar{\partial}f_{k,i}\|_{g_{b^*,i},L^{2p}(\bar{A}_{i,k}^-)}$ if $i \in I^* - I$. Here the exponent of $\frac{2p}{\epsilon_k}$ in (4.28) is crucial:

$$\begin{aligned} \|\bar{\partial}\xi_{k,i}^-\|_{g_{b^*,i},L^{2p}(\bar{A}_{i,k}^-)}^{2p} &\leq \int_{|v_{k,i}|_{b^*}^{-\frac{1}{2}} \leq r \leq e^{\frac{1}{\epsilon_k}} |v_i|_{b^*}^{-\frac{1}{2}}} t_k^{2p} |\nu \circ dq_{v_k, \iota_i^*}^{-1}|_{g_{b^*, \iota_i^*}}^{2p} |v_{k,i}|_{b^*}^{2p} (1+r^2)^{2p-2} r dr d\theta \\ &\leq C t_k^{2p} |v_{k,i}|_{b^*}^{2p} \left(|v_{k,i}|_{b^*}^{-\frac{1}{2}} e^{\frac{1}{\epsilon_k}} \right)^{4p-2} \leq C t_k^{2p}. \end{aligned} \quad (4.35)$$

Since $f_{k,i}$ is constant on $B_{h,k}^+(e^{-\frac{1}{\epsilon_k}} |v_{k,h}|_{b^*}^{\frac{1}{2}})$ for $h \in I^* - I$ with $\iota_h^* = i$ and on $B_{i,k}^-(e^{-\frac{1}{\epsilon_k}} |v_{k,i}|_{b^*}^{\frac{1}{2}})$ if $i \in I^* - I$, the second claim is proved.

Corollary 4.16 *There exists $C > 0$ such that for all k sufficiently large,*

$$\|df_{k,i}\|_{g_{b^*,i},2p} \leq C \quad \text{and} \quad \|\zeta'_{k,i}\|_{g_{b^*,i},2p,1} \leq C \epsilon_k^{\frac{1}{2p}}.$$

Proof: By the quadratic expansion of $\bar{\partial}_{u_i^*} \zeta'_{k,i}$ as in Subsection 3.6,

$$D_{b^*,u_i^*} \zeta'_{k,i} + N_{\bar{\partial},u_i^*} \zeta'_{k,i} = \bar{\partial}_{u_i^*} \zeta'_{k,i}, \quad (4.36)$$

where

$$\|\bar{\partial}_{u_i^*} \zeta'_{k,i}\|_{g_{b^*,i},2p} \leq C \epsilon_k^{\frac{1}{2p}} (\|df_{k,i}\|_{g_{b^*,i},2p} + 1) \quad (4.37)$$

by Lemma 4.15 and

$$\|N_{\bar{\partial},u_i^*} \zeta'_{k,i}\|_{g_{b^*,i},2p} \leq C \|\zeta'_{k,i}\|_{C^0} \|\zeta'_{k,i}\|_{g_{b^*,i},2p,1} \leq C \epsilon_k \|\zeta'_{k,i}\|_{g_{b^*,i},2p,1}, \quad (4.38)$$

by Proposition ?? in [Z1] and Lemma 4.15. Thus, by standard elliptic estimates for u_{b^*} and (4.36)-(4.38),

$$\begin{aligned} \|\zeta'_{k,i}\|_{g_{b^*,i},2p,1} &\leq C (\|D_{b^*,u_i^*} \zeta'_{k,i}\|_{g_{b^*,i},2p} + \|\zeta'_{k,i}\|_{g_{b^*,i},2p}) \\ &\leq C' \epsilon_k^{\frac{1}{2p}} (\|\zeta'_{k,i}\|_{g_{b^*,i},2p,1} + \|df_{k,i}\|_{g_{b^*,i},2p} + 1). \end{aligned} \quad (4.39)$$

On the other hand, since $f_{k,i} = \exp_{b^*,u_i^*} \zeta'_{k,i}$,

$$\|df_{k,i}\|_{g_{b^*,i},2p} \leq C (\|du_i^*\|_{g_{b^*,i},2p} + \|\zeta'_{k,i}\|_{g_{b^*,i},2p,1}). \quad (4.40)$$

If ϵ_k is sufficiently small, the claim follows from equations (4.39) and (4.40).

Corollary 4.17 *There exists $C > 0$ such that for all k sufficiently large, $h \in \hat{I}^*$, and $\delta > 0$,*

$$\|du_{b_k}\|_{g_{v_k},L^{2p}(q_{v_k}^{-1}(B_{h,k}^\pm(\delta)))} \leq C (\epsilon_k^{\frac{1}{2p}} + \delta^{\frac{1}{p}}).$$

Proof: If $h \in \hat{I}$, the statement is immediate from Corollary 4.16; so we assume $h \in I^* - I$. The metric g_v on $q_v^{-1}(B_{h,k}^+(\delta))$ differs by a bounded factor from the metric $q_{v_k, \iota_h^*}^* g_{b^*, i}$. Thus,

$$\begin{aligned} \|du_{b_k}\|_{g_{v_k}, L^{2p}(q_v^{-1}(B_{h,k}^+(\delta)))} &\leq C \|d(f_k \circ q_{v_k, \iota_h^*}^{-1})\|_{g_{b^*, \iota_h^*}^*, L^{2p}(B_{h,k}^+(\delta) - B_{h,k}^+(|v_k, h|^{\frac{1}{2}}))} \\ &= C \|df_{k, \iota_h^*}\|_{g_{b^*, \iota_h^*}^*, L^{2p}(B_{h,k}^+(\delta) - B_{h,k}^+(|v_k, h|^{\frac{1}{2}}))} \\ &\leq C \|df_{k, \iota_h^*}\|_{g_{b^*, \iota_h^*}^*, L^{2p}(B_{h,k}^+(\delta))}. \end{aligned} \quad (4.41)$$

Since $f_{k, \iota_h^*} = \exp_{b^*, u_{\iota_h^*}} \zeta'_{k, \iota_h^*}$, by Corollary 4.16,

$$\|df_{k, \iota_h^*}\|_{g_{b^*, \iota_h^*}^*, L^{2p}(B_{h,k}^+(\delta))} \leq C \left(\|du_{\iota_h^*}\|_{g_{b^*, \iota_h^*}^*, L^{2p}(B_{h,k}^+(\delta))} + \|\zeta'_{k, \iota_h^*}\|_{g_{b^*, \iota_h^*}^*, 2p, 1} \right) \leq C' (\delta^{\frac{1}{p}} + \epsilon^{\frac{1}{2p}}). \quad (4.42)$$

The claim for $B_{h,k}^+(\delta)$ follows from (4.41) and (4.42). The metric g_v on $q_v^{-1}(B_{h,k}^-(\delta))$ differs by a bounded factor from the metric which is the pullback of the metric $g_{b^*, h}$ by the map

$$z \longrightarrow q_N \left(\frac{\phi_{k,h} q_{v_k, \iota_h^*}^*(z)}{v_{k,h}} \right).$$

Thus, similarly to the above,

$$\|du_{b_k}\|_{g_{v_k}, L^{2p}(q_v^{-1}(B_{h,k}^-(\delta)))} \leq C \|df_{k,h}\|_{g_{b^*, h}, L^{2p}(B_{h,k}^-(\delta))}; \quad (4.43)$$

$$\|df_{k,h}\|_{g_{b^*, h}, L^{2p}(B_{h,k}^-(\delta))} \leq C (\delta^{\frac{1}{p}} + \epsilon^{\frac{1}{2p}}). \quad (4.44)$$

The claim for $B_{h,k}^-(\delta)$ follows from (4.43) and (4.44).

4.5 Surjectivity of the Gluing Map

We continue with the notation of Subsection 4.4. In this subsection, for k sufficiently, we use Corollary 4.13 to construct

$$\tilde{v}_k = (\tilde{b}_k, (\tilde{v}_k)_{\hat{I}^*}) \in F^{(0)} \mathcal{T}_\delta$$

and $\tilde{\zeta}_k \in \Gamma(u_{\tilde{v}_k})$ such that $\tilde{b}_k$ is very close to b in $\mathcal{M}_{\mathcal{T}}^{(0)}$, $\|\tilde{\zeta}_k\|_{\tilde{v}_k, p, 1}$ is small, and $u_{b_k} = \exp_{\tilde{v}_k} \tilde{\zeta}_k$. We then look at the elements of $F^{(0)} \mathcal{T}_\delta$ near $\tilde{v}_k$ to find $\tilde{v}'_k$ and $\tilde{\zeta}_k \in \tilde{\Gamma}_+(\tilde{v}'_k)$ such that $u_{b_k} = \exp_{\tilde{v}'_k} \tilde{\zeta}'_k$. If $\mathcal{T}$ is semiregular, we consider only the case $I = \hat{0}$; if $\mathcal{T}$ is regular, we assume $t = 0$.

Let $H = \hat{I} \subset \hat{I}^*$. If $\delta > 0$ and $i \in I^*$, put

$$\Sigma_{i, \delta} = \{(i, z) \in \Sigma_{b^*, i} : r_{b^*, h}(i, z) \leq \delta \ \forall h \in \hat{I} - H \text{ s.t. } \iota_h = i, \ |q_S^{-1}(z)| \geq \delta \text{ if } i \in \hat{I} - H\}.$$

In addition to (4.28), we can assume that our sequence satisfies

$$\|\zeta_{k,i}\|_{g_{b^*, i}, C^2(\Sigma_{i, r_k})} \leq \epsilon_k. \quad (4.45)$$

Let $b'_k = (S, M, I^*; x'_k, (j^*, y^*), u^*)$. By the second assumption in (4.28),

$$d(b^*, b'_k) \leq C \epsilon_k \implies b'_k \in \mathcal{M}_{I^*, C \epsilon_k}^{(H)},$$

since $b^* \in \mathcal{M}_{\mathcal{T}^*}^{(0)}$, where $C > 0$ depends only on b^* . By the last assumption of (4.28), $|v_k|_{b'_k} \leq C\epsilon_k$. Thus, if $\epsilon_k > 0$ is sufficiently small, by Corollary 4.13, there exist

$$\tilde{b}_k \in \mathcal{M}_{\mathcal{T}^*}^{(H)} \quad \text{and} \quad \tilde{v}_k = (\tilde{b}_k, (\tilde{v}_k)_{\hat{I}^*}) \in F^{(H)}\mathcal{T}^*$$

such that

- (1) $d(b, \tilde{b}) \leq C'\epsilon_k$ and $|\tilde{v}_{k,h} - v_{k,h}| \leq C'\epsilon_k |v_{k,h}|_b$ for all $h \in \hat{I}^*$;
(2) if $q_{v_k}(z) \in \Sigma_{\mathcal{T}^*, i}$, $|r_{b^*, h}(q_{v_k}(z))| \geq 3|v_{k,h}|^{\frac{1}{2}}$ for all $h \in \hat{I}^* - H$ such that $\iota_h^* = i$ and $|q_S^{-1}(q_{v_k}(z))| \geq 3|v_{k,i}|^{\frac{1}{2}}$ if $i \in I^* - H$, then $d_b(q_{v_k}(z), q_{\tilde{v}_k}(z)) \leq \epsilon_k$.

It then follows from the second and third assumptions of (4.28) that there exist $\tilde{\zeta}_k \in \Gamma(u_{\tilde{v}_k})$, $\tilde{w}_{k,h} \in T_{x_h(\tilde{v}_k)}\Sigma_{\tilde{v}_k, \iota_h}$ for $h \in H$, and $\tilde{w}_{k,l} \in T_{y_l(\tilde{v}_k)}\Sigma_{\tilde{v}_k, j_l}$ for $l \in M$ such that

$$\begin{aligned} \exp_{\tilde{v}_k} \tilde{\zeta}_k &= u_{b_k}, \quad \exp_{g_{\tilde{v}_k, x_{k,h}}(\tilde{v}_k)} \tilde{w}_{k,h} = x_{k,h}, \quad \exp_{g_{\tilde{v}_k, y_{k,l}}(\tilde{v}_k)} \tilde{w}_{k,l} = y_{k,l}; \\ \|\tilde{\zeta}_k\|_{b^*, C^0}, |\tilde{w}_{k,h}|_{g_{\tilde{v}_k, x_{k,h}}(\tilde{v}_k)} |\tilde{w}_{k,l}|_{g_{\tilde{v}_k, y_{k,l}}(\tilde{v}_k)} &\leq C'\epsilon_k. \end{aligned}$$

Lemma 4.18 *There exists $C > 0$ such that for all k ,*

$$\|\tilde{\zeta}_k\|_{\tilde{v}_k, p, 1} \leq C\epsilon_k^{\frac{1}{2p}}.$$

Proof: By (4.45), (1), and (2), $\|\tilde{\zeta}_k\|_{g_v, C^1} \leq C\epsilon_k$ outside of the necks

$$\tilde{A}_{k,h} = q_{v_k}^{-1} \left(B_{k,h}^+(r_k) \cup B_{k,h}^-(r_k) \right).$$

On the other hand, $\|du_{\tilde{v}_k}\|_{\tilde{v}_k, C^0} \leq C$ by Lemma 3.5 and

$$\|du_{\tilde{v}_k}\|_{\tilde{v}_k, L^p(\tilde{A}_{k,h})} \leq C(\epsilon_k^{\frac{1}{2p}} + r_k^{\frac{1}{p}}) \leq C'\epsilon_k^{\frac{1}{2p}}$$

by Corollary 4.17. The three estimates imply the claim.

Suppose $\hat{I} = \emptyset$ and thus $H = \emptyset$. If k is sufficiently large and $\varpi \in T_{\tilde{v}_k} F^{\emptyset} \mathcal{T}^*$ is such that $2\|\varpi\|_{\tilde{v}_k} < \delta(b^*)$, where δ is as in Lemmas 3.6 and 4.6, let

$$\tilde{b}_k(\varpi) = \tilde{b}_{t\nu}(\tilde{v}_k(\varpi)) = (S, M, \{\hat{0}\};, (\hat{0}, \bar{y}(\varpi)), \tilde{u}_{\varpi, t\nu})$$

be the tuple defined as in Subsections 3.4 and 3.6. Let $\tilde{\zeta}_k(\varpi) \in \Gamma(\tilde{u}_{\varpi, t\nu})$ and $\tilde{w}_{k,l}(\varpi) \in T_{y_l(\varpi)}\Sigma_{\varpi, j_l}$ for $l \in M$ be given by

$$\exp_{\tilde{v}_k(\varpi)} \tilde{\zeta}_k(\varpi) = u_{b_k}, \quad \exp_{g_{\tilde{v}_k(\varpi), y_{k,l}}(\varpi)} \tilde{w}_{k,l}(\varpi) = y_{k,l}; \quad \|\tilde{\zeta}_k(\varpi)\|_{b^*, C^0}, |\tilde{w}_{k,l}(\varpi)|_{g_{\tilde{v}_k(\varpi), y_{k,l}}(\varpi)} \leq 2C'\epsilon_k.$$

We need to find ϖ such that $\tilde{\pi}_{\varpi, -}\tilde{\zeta}_k(\varpi) = 0$ and $y_l(\varpi) = y_{k,l}$, or equivalently

$$S_{\varpi} \tilde{\pi}_{\varpi, -} \tilde{\zeta}_k(\varpi) = 0 \quad \text{and} \quad S_{\varpi} \tilde{w}_{k,l}(\varpi) = 0, \tag{4.46}$$

where $S_{\varpi} \tilde{w}_{k,l}(\varpi)$ denotes the parallel transform of $\tilde{w}_{k,l}(\varpi)$ back to $y_l(\tilde{v}_k)$ along the g_v -geodesic

$$s \longrightarrow \exp_{y_l(\tilde{v}_k)} s w_l(\varpi).$$

Lemma 4.19 *There exists $C > 0$ such that for all k sufficiently large and $\varpi, \varpi' \in T_{\tilde{v}_k} F^{(\emptyset)} \mathcal{T}^*$ with $2\|\varpi\|_{\tilde{v}_k} < \delta(b^*)$,*

$$\begin{aligned} S_{\varpi} \tilde{\pi}_{\varpi, -\tilde{\zeta}_k}(\varpi) &= \tilde{\pi}_{v, -\tilde{\zeta}_k} + \tilde{N}^{(0)}(\tilde{\zeta}_k, \varpi) - \tilde{\pi}_{v, -\zeta_{\varpi}} + N^{(0)}(\varpi), \\ S_{\varpi} \tilde{w}_{k, h}(\varpi) &= \tilde{w}_{k, h} + \tilde{N}^{(h)}(\tilde{w}_{k, h}, \varpi) - w_h(\varpi) + N^{(h)}(\varpi) \quad \forall h \in M, \end{aligned}$$

where ζ_{ϖ} is as in Section 3.4 and $\tilde{N}^{(l)}$ and $N^{(l)}$ satisfy

$$\begin{aligned} \|\tilde{N}^{(0)}(\tilde{\zeta}_k, \varpi) - \tilde{N}^{(0)}(\tilde{\zeta}_k, \varpi')\|_{\tilde{v}_k, 2} &\leq C \|\tilde{\zeta}_k\|_{\tilde{v}_k, 2} \|\varpi - \varpi'\|_{\tilde{v}_k}; \\ |\tilde{N}^{(l)}(\varpi, \tilde{w}_{k, l}) - \tilde{N}^{(l)}(\varpi', \tilde{w}_{k, l})|_{g_{\tilde{v}_k, y_l}(\tilde{v}_k)} &\leq C \|\tilde{w}_{k, l}\|_{g_{v_k, y_l}(\tilde{v}_k)} \|\varpi - \varpi'\|_{\tilde{v}_k} \quad \forall l \in M; \\ \|N^{(0)}(\varpi) - N^{(0)}(\varpi')\|_{\tilde{v}_k, 2} &\leq C (\|\varpi\|_{\tilde{v}_k} + \|\varpi'\|_{\tilde{v}_k}) \|\varpi - \varpi'\|_{\tilde{v}_k}; \\ |N^{(l)}(\varpi) - N^{(l)}(\varpi')|_{g_{\tilde{v}_k, y_l}(\tilde{v}_k)} &\leq C (\|\varpi\|_{\tilde{v}_k} + \|\varpi'\|_{\tilde{v}_k}) \|\varpi - \varpi'\|_{\tilde{v}_k} \quad \forall l \in M. \end{aligned} \tag{4.47}$$

Proof: This lemma follows from a pointwise Riemannian geometry estimate on $S_{\varpi} \tilde{\zeta}_k(\varpi) - (\tilde{\zeta}_k - \zeta_{\varpi})$ and the fact that all statements in Lemmas 3.6 and 4.6 can be written in a form similar to (4.47), e.g. for all $\xi \in \Gamma(\tilde{v}_k)$

$$\|S_{\varpi} \tilde{\pi}_{\varpi, -R_{\varpi} \xi} - S_{\varpi'} \tilde{\pi}_{\varpi', -R_{\varpi'} \xi}\|_{\tilde{v}_k, 2} \leq C \|\varpi - \varpi'\|_{\tilde{v}_k, 2} \|\xi\|_{\tilde{v}_k, 2}.$$

The latter fact can be seen from the two lemmas and the definitions of R_{ϖ} and S_{ϖ} in Subsection 3.4.

Lemma 4.20 *There exist $C, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}^*}^{(0)}, \mathbb{R}^+)$ such that for all $v \in F^{(H)} \mathcal{T}_\delta^*$ and $\varpi \in T_v F^{(H)} \mathcal{T}_\delta^*$ with $\|\varpi\|_{\tilde{v}_k} \leq \delta(b)$,*

$$\|\zeta_{\varpi}\|_{v, 2} \leq C(b) \|\tilde{\pi}_{v, -\zeta_{\varpi}}\|_{v, 2}.$$

Proof: It can be seen directly from the definitions that

$$\|\zeta_{\varpi}\|_{v, 2} \leq (1 + C(b_v)|v|) \|\pi_{v, -\zeta_{\varpi}}\|_{v, 2}.$$

The claim then follows from the proof of (2b) of Lemma 3.12.

Corollary 4.21 *There exist a neighborhood U of b^* in $\mathcal{M}_{\mathcal{T}^*}^{(0)}$ and $\delta, \epsilon > 0$ such that for all $v \in F^{(\emptyset)} \mathcal{T}_\delta^*|U$, $\xi \in \tilde{\Gamma}_-(v)$ with $\|\xi\|_{v, 2} < \delta$, $w_h \in T_{x_h(v)} \Sigma_{v, \iota_h}$ with $|w_h|_{g_{v, x_h}(v)} < \delta$ and $w_l \in T_{y_l(v)} \Sigma_{v, j_l}$ for $l \in M$ with $|w_l|_{g_{v, y_l}(v)} < \delta$, the system of equations*

$$\tilde{\pi}_{v, -\zeta_{\varpi}} - N^{(0)}(\varpi) = \xi, \quad w_h(\varpi) - N^{(l)}(\varpi) = w_h \quad \forall h \in H, M$$

has a (unique) solution $\varpi \in T_v F^{(\emptyset)} \mathcal{T}$ with $\|\varpi\|_v < \epsilon$.

Proof: By Lemmas 3.6 and 4.20,

$$C^{-1} \|\varpi\|_v \leq \|\tilde{\pi}_{v, -\xi}\|_{v, 2} + \sum_{h \in H} |w_h(\varpi)|_{g_{v, x_h}(v)} + \sum_{l \in M} |w_l(\varpi)|_{g_{v, y_l}(v)} \leq C \|\varpi\|_v.$$

whenever b_v lies near b^* . Thus, the claim follows from (4.47) by the usual contraction-principle argument.

Corollary 4.22 *Let $\mathcal{T}^* = (S, M, \{\hat{0}\}; j^*, \lambda^*)$ be a simple bubble type. If $\mathcal{T}^*$ is regular, the map*

$$\tilde{\gamma}_{\mathcal{T}^*}: F\mathcal{T}_\delta^* \longrightarrow \bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle}$$

contains a neighborhood of $\mathcal{M}_{\mathcal{T}^}$ in $\bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle}$. If $\mathcal{T}^*$ is semiregular, $H = \emptyset$, and k is sufficiently large, there exists $\tilde{v}_k \in F^{(\emptyset)}\mathcal{T}_\delta^*$ such that $b_k = \tilde{\gamma}_{\mathcal{T}^*, t_k \nu}(\tilde{v}_k)$.*

Proof: The second statement is immediate from Lemmas 4.18 and 4.19 and Corollary 4.21. If $\mathcal{T}^*$ is regular, what we have shown is that the image of $\tilde{\gamma}_{\mathcal{T}^*}$ contains a neighborhood of $\mathcal{M}_{\mathcal{T}^*}$ in $\bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle} \cup \mathcal{M}_{\mathcal{T}^*}$. Furthermore, there exists a sequence of neighborhoods $U_1 \supset U_2 \supset \dots$ of b^* in $\bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle}$ such that $\bigcap_k U_k = \{[b^*]\}$. If $[b_k] \in \mathcal{M}_{\mathcal{T}}$ is a sequence of bubble maps converging to $[b^*] \in \mathcal{M}_{\mathcal{T}^*}$, it can be assumed that $[b_k] \in U_k$. By the above statement applied to $\mathcal{T}$, we can choose sequences

$$\{[b_{kr}]\} \subset \mathcal{M}_{\langle \mathcal{T}^* \rangle} = \mathcal{M}_{\langle \mathcal{T} \rangle}$$

such that for each fixed k the sequence $\{[b_{kr}]\}$ converges to $[b_k]$. Since U_k is an open neighborhood of $[b_k]$, it can be assumed that $[b_{kr}] \in U_k$ for all r . By the above, the image of $\tilde{\gamma}_{\mathcal{T}^*}|F\mathcal{T}_{\frac{1}{2}\delta}^*$ contains $U_k \cap \mathcal{M}_{\langle \mathcal{T}^* \rangle}$ if k is sufficiently large. Thus, for all r there exists $v_{kr} \in F\mathcal{T}_{\frac{1}{2}\delta}^*$ such that $\tilde{\gamma}_{\mathcal{T}^*}(v_{kr}) = [b_{kr}]$. Let $\tilde{v}_k \in F\mathcal{T}_\delta^*$ be the limit of the sequence v_{kr} with k fixed. Then, by continuity of the map $\tilde{\gamma}_{\mathcal{T}^*}$, see Corollary 4.5,

$$\tilde{\gamma}_{\mathcal{T}^*}(\tilde{v}_k) = \lim_{r \rightarrow \infty} \tilde{\gamma}_{\mathcal{T}^*}(\tilde{v}_{kr}) = \lim_{r \rightarrow \infty} [b_{kr}] = [b_k].$$

Thus, the image of $\tilde{\gamma}_{\mathcal{T}^*}$ contains a neighborhood of $\mathcal{M}_{\mathcal{T}^*}$ in $\bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle}$.

Corollary 4.23 *If $\mathcal{T}^* = (S, M, I^*; j^*, \lambda^*)$ is a simple regular bubble type, the map*

$$\tilde{\gamma}_{\mathcal{T}^*}: F\mathcal{T}_\delta^* \longrightarrow \bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle}$$

is a homeomorphism onto an open neighborhood of $\mathcal{M}_{\mathcal{T}^}$ in $\bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle}$ provided $\delta \in C^\infty(\mathcal{M}_{\mathcal{T}^*}; \mathbb{R}^+)$ is sufficiently small.*

Proof: By Corollaries 4.5, 4.10, 4.22, the map $\tilde{\gamma}_{\mathcal{T}^*}: F\mathcal{T}_\delta^* \longrightarrow \bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle}$ is a continuous bijection onto a neighborhood of $\mathcal{M}_{\mathcal{T}^*}$ in $\bar{\mathcal{M}}_{\langle \mathcal{T}^* \rangle}$. In addition, the proof of Corollary 4.22 shows that $\tilde{\gamma}_{\mathcal{T}^*}$ is an open map.

5 Appendix

5.1 Properties of Smooth Families of Metrics on Σ

Let m be a positive integer and

$$\aleph = \{x = x_{[m]} : x_h \in \Sigma, x_h \neq x_l \text{ if } h \neq l\}.$$

Suppose $\{g_x : x \in \aleph\}$ is a smooth family of metrics on Σ such that for any $x = x_{[m]} \in \aleph$ the metric g_x is flat on a neighborhood of x_h in Σ for all $h \in [m]$. If $x = x_{[m]} \in \aleph$ and $v \in T_y \Sigma$, let

$$T_x \aleph = \bigoplus_{h \in [m]} T_{x_h} \Sigma, \quad |v|_x = |v|_{g_x, y}.$$

If $w = w_{[m]} \in T_x \mathbb{N}$, let $|w|$ denote $\sum_{h \in [m]} |w_h|_x$. Define $x(w) \in \Sigma^m$ by

$$x(w) = (x_1(w), \dots, x_m(w)) = (\exp_{g_x, x_1} w_1, \dots, \exp_{g_x, x_m} w_m).$$

We denote by $\phi_{x,y}$ the map $\phi_{g_x, y}$ and by $B_x(y, \delta)$ the set $B_{g_x}(y, \delta)$ described in Subsection 1.3. If $\delta: \mathbb{N} \rightarrow \mathbb{R}$, let

$$T\mathbb{N}_\delta = \{(x, w): x \in \mathbb{N}; w \in T_x \mathbb{N}, |w|_x < \delta(x)\}.$$

Lemma 5.1 *There exist $\delta \in C^\infty(\mathbb{N}; \mathbb{R}^+)$ and a smooth families of holomorphic maps*

$$\{\tilde{p}_{h,(x,w)}: \{z \in B_x(x_h, \delta(x))\} \rightarrow \Sigma \mid (x, w) \in T\mathbb{N}_\delta\},$$

such that each map $\tilde{p}_{h,(x,w)}$ is a $(g_x, g_{x(w)})$ -isometry,

$$\begin{aligned} d\phi_{x,x_h}|_{x_h(w)} \phi_{x(w), x_h(w)} \tilde{p}_{h,(x,w)}(z) &= \phi_{x,x_h}(z), \\ \text{and} \quad d_{g_x}(z, \tilde{p}_{h,(x,w)}(z)) &\leq 2|w|_x \quad \forall z \in B_x(x_h, \delta(x)). \end{aligned} \quad (5.1)$$

In particular, both sides of (5.1) are defined.

Proof: We choose δ such that if $w \in T_x \mathbb{N}$ and $|w| \leq 4\delta(x)$, then $x(w) \in \mathbb{N}$ and the metric $g_{x(w)}$ is flat on $B_x(x_h, 2\delta(x))$. This choice of δ insures that both sides of (5.1) are defined. Equation (5.1) is equivalent to

$$\phi_{x(w), x_h(w)} \tilde{p}_{h,(x,w)}(z) = d \exp_{g_x, x_h} |_{w_h} \phi_{x,x_h}(z) = \phi_{x,x_h}(z) + d \exp_{g_x, x_h} |_{w_h} w_h, \quad (5.2)$$

since the metric g_x is flat on $B_x(x_h, 2\delta(x))$. This equation defines the required map $\tilde{p}_{h,(x,w)}$. Since the metrics g_x and $g_{x(w)}$ are flat on $B_x(x_h, 2\delta(x))$, the maps $\phi_{x,x_h(w)} z$ and $\phi_{x(w), x_h(w)}$ are holomorphic, and thus $\tilde{p}_{h,(x,w)}$ is holomorphic. Taking the differential of (5.2), we obtain

$$d\phi_{x(w), x_h(w)} |_{\tilde{p}_{h,(x,w)}(z)} \circ d\tilde{p}_{h,(x,w)} |_z = d\phi_{x,x_h(w)} |_z. \quad (5.3)$$

Since $\phi_{x(w), x_h(w)}$ and $\phi_{x,x_h(w)}$ are $(g_{x(w)}, g_{x(w)})$ - and (g_x, g_x) -isometries, respectively, on $B_x(x_h, 2\delta(x))$, it follows that $\tilde{p}_{h,(x,w)}$ is a $(g_x, g_{x(w)})$ -isometry on $B_x(x_h, 2\delta(x))$. By (5.2),

$$d_{g_x}(z, \tilde{p}_{h,(x,w)}(z)) \leq |w_h|_x + |(\phi_{x(w), x_h(w)} - \phi_{x,x_h(w)}) \tilde{p}_{h,(x,w)}(z)|_x \leq |w_h|_x + C(x)|w|\delta(x), \quad (5.4)$$

since the family of metrics is smooth. If $C(x)\delta(x) < 1$, the remaining claim of the lemma follows from (5.4).

Lemma 5.2 *There exist $\delta, C_k \in C^\infty(\mathbb{N}; \mathbb{R}^+)$, where k is a positive integer, $\alpha_h \in C^\infty(T\mathbb{N}_\delta; \mathbb{C})$, and smooth families of maps*

$$\{\Theta_{w,h}: \{v \in T_{x_h} \mathbb{N} : |v|_x < \delta(x)\} \rightarrow T_{x_h} \Sigma \mid (x, w) \in T\mathbb{N}_\delta\}$$

such that each map $\Theta_{w,h}$ is holomorphic, $\Theta_{w,h}(0) = 0$, $\Theta'_{w,h}(0) = 0$, $\|\Theta_{w,h}^{(k)}\|_{C^0} \leq C_k(x)|w|$, $|\alpha_h(w)| \leq C_0(x)|w|$, and

$$d\phi_{x,x_h}|_{x_h(w)} d\phi_{x(w), x_h(w)} |_{x_h} (\phi_{x(w), x_h} z) = (1 + \alpha_h(w)) \phi_{x,x_h} z + \Theta_{w,h}(\phi_{x,x_h} z). \quad (5.5)$$

for all $z \in B_x(x_h, \delta(x))$. In particular, both sides of (5.5) are defined.

Proof: We choose δ such that if $w \in T_x \mathfrak{N}$ and $|w| \leq 4\delta(x)$, then $x(w) \in \mathfrak{N}$ and the metric $g_{x(w)}$ is flat on $B_x(x_h, 4\delta(x))$. This choice of δ insures that both side of (5.5) are defined. If w and z are as in the statement of the lemma, by the flatness of the metric $g_{x(w)}$ near x_h , $\mathbb{C}$ -linearity of the differential of the exponential map near zero, and the smoothness of the family of the metrics

$$d\phi_{x,x_h}|_{x_h(w)} d\phi_{x(w),x_h(w)}|_{x_h}(\phi_{x(w),x_h} z) = (1 + a_h(w))(\phi_{x(w),x_h} z), \quad (5.6)$$

for some $a_h \in C^\infty(T\mathfrak{N}_\delta; \mathbb{C})$ such that $a_h(0) = 0$. Note that if $g_{x(w)} = g_x$, $a_h(w) = 0$, since the metric g_x is flat on $B_x(x_h, |w|)$. The map

$$\{v \in T_{x_h} \Sigma : |v|_x < 2\delta(x)\} \longrightarrow T_{x_h} \Sigma, \quad v \longrightarrow \phi_{x(w),x_h} \phi_{x,x_h}^{-1} v - v$$

is holomorphic since $\phi_{x(w),x_h}$ and ϕ_{x,x_h} are, and vanishes at 0. Thus,

$$\phi_{x(w),x_h} \phi_{x,x_h}^{-1} v = (1 + b_h(w))v + \Theta_{w,h}(v), \quad (5.7)$$

for some $b_h(w) \in \mathbb{C}$ and holomorphic function $\Theta_{w,h}$ such that $\Theta_{w,h}(0), \Theta'_{w,h}(0) = 0$. Equation (5.5) follows from (5.6) and (5.7). Smoothness of b_h and $\Theta_{w,h}$ in w follows from the smoothness of the family of the metrics. The bounds on α_h and the derivatives of $\Theta_{w,h}$ follow from their smoothness and compactness of the fibers of

$$\{w \in T_x \mathfrak{N} : |w| \leq \delta(x)\} \longrightarrow \mathfrak{N}.$$

Lemma 5.3 *There exist $\delta, C \in C^\infty(\mathfrak{N}; \mathbb{R}^+)$ and smooth families of maps*

$$N_h : \{(x, w) : x \in \mathfrak{N}; (x, w) \in T\mathfrak{N}_\delta\} \longrightarrow T\mathfrak{N}$$

such that $|N_h(w, w_h)|_x \leq C(x)|w||w_h|$ and

$$d\phi_{x,x_h}|_{x_h(w)}(\phi_{x(w),x_h(w)} x_h) = -w_h + N_h(w, w_h). \quad (5.8)$$

In particular, the left-hand side of (5.8) is defined.

Proof: We take δ as in Lemma 5.2. Then, the left-hand side of (5.8) is defined and

$$\begin{aligned} & d\phi_{x,x_h}|_{x_h(w)}(\phi_{x(w),x_h(w)} x_h) \\ &= d\phi_{x,x_h}|_{x_h(w)}(\phi_{x,x_h(w)} x_h) + d\phi_{x,x_h}|_{x_h(w)} \{(\phi_{x(w),x_h(w)} x_h) - (\phi_{x,x_h(w)} x_h)\} \\ &= -w_h + N_h(w, w_h), \end{aligned} \quad (5.9)$$

where $N(\cdot, \cdot)$ is some smooth function of both variables, that vanishes if either input is zero. Equation (5.8) is thus proved, while the bound on N_h is obtained from its smoothness and compactness of the fibers as in the proof of Lemma 5.2.

Lemma 5.4 *There exists $\delta \in C^\infty(\mathfrak{N}; \mathbb{R}^+)$ such that for any $x \in \mathfrak{N}$, $v \in T_x \mathfrak{N}$ with $|v| < \delta(x)$, and $c = c_{[m]} \in \mathbb{C}^m$ with $|c||v| < \delta(x)$, there exists $w \in T\mathfrak{N}$ with $|w_h|_x < 2|c_h||v_h|_x$ such that for $z \in B_x(x_h, 4\delta(x)^{\frac{1}{2}})$*

$$d\phi_{x,x_h}|_{x_h(w)}(\phi_{x(w),x_h(w)} z) = (1 + \alpha_h(w))(c_h v_h + \phi_{x,x_h} z) + \Theta_{w,h}(\phi_{x,x_h} z), \quad (5.10)$$

where $\alpha_h(w)$ and $\Theta_{w,h}$ are as in Lemma 5.2. In particular, both sides of (5.10) are defined.

Proof: We start by choosing δ so that $8\delta^{\frac{1}{2}}$ is smaller than the function δ of Lemmas 5.2 and 5.3. By flatness of the metric $g_{x(w)}$ on $B(x_h, 8\delta(x)^{\frac{1}{2}})$ for $w \in T_x \mathfrak{N}$ with $|w| < \delta(x)$

$$\phi_{x(w), x_h(w)} z = d\phi_{x(w), x_h(w)}|_{x_h} \phi_{x(w), x_h} z + \phi_{x(w), x_h(w)} x_h \quad (5.11)$$

for any $z \in B(x_h, 4\delta(x)^{\frac{1}{2}})$. Taking $d\phi_{x, x_h}|_{x_h(w)}$ of both sides of (5.11) and applying Lemmas 5.2 and 5.3, we obtain

$$d\phi_{x, x_h}|_{x_h(w)}(\phi_{x(w), x_h(w)} z) = (1 + \alpha_h(w)) \phi_{x, x_h} z + \Theta_{w, h}(\phi_{x, x_h} z) - w_h + N_h(w, w_h). \quad (5.12)$$

Thus, we need to solve the equations

$$-w_h + N_h(w, w_h) = (1 + \alpha_h(w)) c_h v_h. \quad (5.13)$$

Let $\Psi_h(w) = N_h(w, w_h) - (1 + \alpha_h(w)) c_h v_h$. If $|w| \leq 2|c_h||v_h|$, then by Lemmas 5.2 and 5.3,

$$\begin{aligned} |\Psi(w)| &\leq C(x)|c||v|(2|c||v| + 1) \leq 2|c||v|, \\ |\Psi(w) - \Psi(w')| &\leq C(x)|c||v||w - w'| \leq \frac{1}{2}|w - w'|, \end{aligned} \quad (5.14)$$

provided $4C(x)\delta(x) < 1$. In such a case, Ψ is a contracting operator, and thus (5.13) has a unique solution $w \in T_x \mathfrak{N}$ with $|w| < 2|c||v|$. The estimate $|w_h| < 2|c_h||v_h|$ follows directly from (5.13) if $\delta(x)$ is sufficiently small.

Corollary 5.5 *There exist $\delta, C_k \in C^\infty(\mathfrak{N}; \mathbb{R}^+)$, where k is a positive integer, such that for any $x \in \mathfrak{N}$, $v \in T_x \mathfrak{N}$ with $|v| < \delta(x)$, $c = c_{[m]} \in \mathbb{C}^m$ with $|c| < \delta(x)$, and $r = r_{[m]} \in \mathbb{R}^m$ with $|r| < \frac{1}{2}$, there exists $\tilde{x} \in \mathfrak{N}$ and $\tilde{v} \in T_{\tilde{x}} \mathfrak{N}$ such that*

- (1) $\tilde{x}_h \in B_x(x_h, 2|c_h||v_h|)$, $|\frac{g_{\tilde{x}}}{g_x} - 1| \leq C_1(x)|c||v|$, $|\tilde{v}_h|_{\tilde{x}} - |v_h|_x| \leq C_1(x)(|c||v| + |r_h|)|v_h|$;
- (2) for any $z \in B_x(x_h, 4\delta(x)^{1/2})$,

$$\frac{\phi_{\tilde{x}, \tilde{x}_h} z}{\tilde{v}_h} = (1 + r_h) \left\{ c_h + \frac{\phi_{x, x_h} z}{v_h} + \Theta_{v, c, r, h} \left(\frac{\phi_{x, x_h} z}{v_h} \right) \right\}, \quad (5.15)$$

where $\Theta_{v, c, r, h}$ is a holomorphic function, varying smoothly with the parameters, such that

$$\Theta_{v, c, r, h}(0) = 0, \quad \Theta'_{v, c, r, h}(0) = 0, \quad \|\Theta_{v, c, r, h}^{(k)}\|_{C^0} \leq C_k(x)|c||v||v_h|^{k-1}.$$

Proof: Let δ be as in Lemma 5.4. Given v and c as in the statement of the lemma, let $w \in T_x \mathfrak{N}$ be the element provided by Lemma 5.4. Take

$$\tilde{x}_h = x_h(w), \quad \tilde{v}_h = (1 + r_h)^{-1} (1 + a_h(w)) d\phi_{x, x_h}^{-1}|_{w_h} v_h.$$

The estimates in (1) are immediate from Lemma 5.4, provided δ is sufficiently small. The inequalities in (2) arise from the smooth dependence of w on x , v , and c in Lemma 5.4, and the fact that w is zero if either $v = 0$ or $c = 0$.

5.2 Sobolev Inequalities for the Metrics g_v

In this subsection, we prove (3) of Lemma 3.5. The reason this estimate holds is that (Σ_v, g_v) can be written as a union of the surfaces $(\Sigma_{b_v, i}, g_v)$ with small disks missing and annuli $(\tilde{A}_{v, h}^\pm, g_v)$ that are uniformly equivalent to annuli in $\mathbb{R}^2$ with the smaller radius less than half of the larger.

Suppose $\mathcal{T} = (S, M, I; j, \lambda)$ is a bubble type and

$$v = (b, v_{\hat{I}}) = ((S, M, I; x, (j, y), u), v_{\hat{I}}) \in F^{(0)}\mathcal{T}_\delta.$$

For any $h \in \hat{I}$ and $i \in I$, put

$$\begin{aligned} \tilde{A}_{v, h}^- &= q_{v, i_h}^{-1} \left(\left\{ (\iota_h, z) \in \Sigma_{b_v, \iota_h} : (2\delta_{\mathcal{T}}(b_v))^{-1} |v_h|_b \leq r_{b, h}(z) \leq |v_h|_b^{\frac{1}{2}} \right\} \right); \\ \tilde{A}_{v, h}^+ &= q_{v, i_h}^{-1} \left(\left\{ (\iota_h, z) \in \Sigma_{b_v, \iota_h} : |v_h|_b^{\frac{1}{2}} \leq r_{b, h}(z) \leq 2\delta_{\mathcal{T}}(b_v) \right\} \right); \\ S_{v, i} &= q_{v, i}^{-1} \left(\left\{ (i, z) \in S_{b_v, i} : r_{b, h}(z) \geq \delta_{\mathcal{T}}(b_v) \text{ if } \iota_h = i; |q_S^{-1}(z)| \geq \delta_{\mathcal{T}}(b_v) \text{ if } i > 0 \right\} \right). \end{aligned}$$

Let $\tilde{A}_{v, h}$ denote $\tilde{A}_{v, h}^- \cup \tilde{A}_{v, h}^+$.

Lemma 5.6 *For any $p > 2$, there exists $C_p \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R})$ such that for any $v \in F^{(0)}\mathcal{T}_\delta$ and $h \in \hat{I}$,*

$$\xi \in \tilde{\Gamma}_c(\tilde{A}_{v, h}; u_v^* TV) \implies \|\xi\|_{C^0} \leq C_p(b_v) \|\xi\|_{g_v, p, 1}.$$

Proof: By construction of the metric g_v , $g_v|_{\tilde{A}_{v, h}}$ is the pullback of the metric g_{v, ι_h} on $q_{v, \iota_h}(\tilde{A}_{v, h})$ by the map q_{v, ι_h} . Furthermore, the metric g_{v, ι_h} on $q_v(\tilde{A}_{v, h}^\pm)$ differs from the standard metric on the annulus $B_{2\delta_{\mathcal{T}}(b_v), |v_h|_b^{\frac{1}{2}}} \subset \mathbb{R}^2$ by factors bounded by $C(b_v)$. Since $\|du_v\|_{g_v, p} \leq C_p(b_v)$ by (1) of Lemma 3.5, the claim follows from Proposition ?? in [Z1].

Proposition 5.7 *For any $p > 2$, there exists $C_p \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R})$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$,*

$$\|\xi\|_{C^0} \leq C_p(b_v) \|\xi\|_{g_v, p, 1} \quad \text{for all } \xi \in \Gamma(u_v).$$

Proof: (1) Note that $g_v|_{S_{v, i}}$ is the pull-back of the metric $g_{b_v, i}$ on $q_{v, i}(S_{v, i})$ by the map $q_{v, i}$. Thus, by Proposition ?? in [Z1], if $\xi \in \Gamma_c(S_{v, i}; u_v^* TV)$,

$$\|\xi\|_{C^0} = \|\xi \circ q_{v, i}\|_{C^0} \leq C_p(\|du_v \circ q_{v, i}\|_{g_{b_v, i}, p}) \|\xi \circ q_{v, i}\|_{g_{b_v, i}, p, 1} = C_p(b_v) \|\xi\|_{g_v, p, 1},$$

since ξ vanishes outside of $S_{v, i}$.

(2) We now define a partition of unity subordinate to $\{S_{v, i}, \tilde{A}_{v, h} : i \in I, h \in \hat{I}\}$. Put

$$\begin{aligned} \eta_{v, h}^+(z) &= \begin{cases} 1 - \beta_{\delta_{\mathcal{T}}^2(b_v)}(r_{b_v, h}(q_{v, \iota_h}(z))), & \text{if } q_{v, \iota_h}(z) \in \Sigma_{b_v, \iota_h}; \\ 1, & \text{otherwise;} \end{cases} \\ \eta_{v, h}^-(z) &= \begin{cases} 1 - \beta_{\delta_{\mathcal{T}}^2(b_v)}(|q_S^{-1}q_v(z)|), & \text{if } q_{v, h}(z) \in \Sigma_{b_v, h}; \\ 1, & \text{otherwise;} \end{cases} \quad \tilde{\eta}_v(z) = 1 - \prod_{h \in \hat{I}} \eta_{v, h}^-(z) \eta_{v, h}^+(z). \end{aligned}$$

Note that $d\eta_{v,h}^\pm$ is supported in $\tilde{A}_{v,h}^\pm$. It follows from the definition of g_v that

$$\|d\eta_{v,h}^\pm\|_{g_v, C^1} = \|d(\eta_{v,h}^\pm \circ q_{v,i_h}^{-1})\|_{g_{v,i_h}, C^1} \leq C(b_v).$$

Thus, if $\xi \in \Gamma(u_v)$ by (1) and Lemma 5.6,

$$\begin{aligned} \|\xi\|_{C^0} &\leq \sum_{i \in I} \|\tilde{\eta}\xi\|_{C^0(S_{v,i})} + \sum_{h \in \hat{I}} \|\eta_{v,h}^- \eta_{v,h}^+ \xi\|_{C^0} \leq C_p(b_v) \left(\|\tilde{\eta}\xi\|_{g_v, p, 1} + \sum_{h \in \hat{I}} \|\eta_{v,h}^- \eta_{v,h}^+ \xi\|_{g_v, p, 1} \right) \\ &\leq C_p(b_v) \left(|I| \|\xi\|_{g_v, p, 1} + 2 \sum_{h \in \hat{I}} \|\eta_{v,h}^- \eta_{v,h}^+\|_{g_v, C^1} \|\xi\|_{g_v, p} \right) \leq C'_p(b_v) \|\xi\|_{g_v, p, 1}. \end{aligned}$$

5.3 Elliptic Estimates for the Metrics g_v

This subsection contains the proof of (4) of Lemma 3.5, the main elliptic estimate for the operators D_v and the modified Sobolev norms. This estimate does *not* hold for the standard Sobolev norms. The argument is essentially the same as in [LT], but we do include all of the details, based on [Z1], and state a sharper estimate.

Let $\mathcal{T}$, v , $\tilde{A}_{v,h} = \tilde{A}_{v,h}^- \cup \tilde{A}_{v,h}^+$, and $S_{v,i}$ be as in Subsection 5.2. If $\iota_h = 0$, the metric $g_{b_v, \hat{0}}$ is flat on $B_{b_v, h}(\delta_{\mathcal{T}}(b_v)^{\frac{1}{2}})$. Thus, for any $h \in \hat{I}$, we can choose conformal polar coordinates (r, θ) on $\tilde{A}_{v,h}$ such that $r(z) = r_{b,v}(q_{v,\iota_h}(z))$. Since $g_v|_{\tilde{A}_{v,h}}$ is the pullback of the metric g_{v,ι_h} on $q_{v,\iota_h}(\tilde{A}_{v,h})$ by the map q_{v,ι_h} ,

$$g_v = \left\{ (1 - \beta_{|v_h|}(2r)) \frac{2C(b_v)}{|v_h| + |v_h|^{-1}r^2} + \beta_{|v_h|}(r) \right\} (dr^2 + r^2 d\theta^2) \quad \text{on } \tilde{A}_{v,h}. \quad (5.16)$$

Similarly, since $\rho_v = \rho_{v,\iota_h} \circ q_{v,\iota_h}$

$$\rho_v = r^2 + \frac{|v_h|^2}{r^2} \quad \text{on } \tilde{A}_{v,h}. \quad (5.17)$$

Lemma 5.8 *For all $p > 1$, there exists $C_p \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R})$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$ and $h \in \hat{I}$,*

$$\xi \in \Gamma_c(\tilde{A}_{v,h}; u_v^* TV) \implies \left(\int_{\tilde{A}_{v,h}} \rho_v^{-\frac{p-2}{p}} |\nabla^{b_v} \xi|^2 \right)^{\frac{1}{2}} \leq C_p(b_v) (\|D_v \xi\|_{v,p} + \|\xi\|_{v,p}).$$

Proof: (1) Let ϵ_1 and ϵ_2 denote $(2\delta_{\mathcal{T}}(b_v))^{-1} |v_h|$ and $2\delta_{\mathcal{T}}(b_v)$, respectively. Note that the integral on the left-hand side in the statement of the lemma is conformally invariant. With respect to the metric $dr^2 + r^2 d\theta^2$,

$$|D_v \xi|_{(r,\theta)} = \left| \frac{D}{dr} \xi + Jr^{-1} \frac{D}{d\theta} \xi \right|_{(r,\theta)},$$

where $\frac{D}{dr}$ and $\frac{D}{d\theta}$ denote covariant differentiation with respect to the connection ∇^{b_v} and the norms are taken with respect to the metric $g_{V,b}$ on V . Thus,

$$a_h^2 \equiv \int_{\tilde{A}_{v,h}} \rho_v^{-\frac{p-2}{p}} |\nabla^{b_v} \xi|^2 \leq \|D_v \xi\|_{v,p}^2 - 2 \int_0^{2\pi} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \left\langle \frac{D}{dr} \xi, J \frac{D}{d\theta} \xi \right\rangle dr d\theta. \quad (5.18)$$

Since $\nabla^{b_v} J = 0$, using integration by parts twice, we obtain

$$\begin{aligned}
& \int_0^{2\pi} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \left\langle \frac{D}{dr} \xi, J \frac{D}{d\theta} \xi \right\rangle dr d\theta = - \int_0^{2\pi} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \left\langle \frac{D}{d\theta} \frac{D}{dr} \xi, J \xi \right\rangle dr d\theta \\
& = - \int_0^{2\pi} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \left(\left\langle \frac{D}{dr} \frac{D}{d\theta} \xi, J \xi \right\rangle - \langle \mathcal{R}(u_r, u_\theta) \xi, J \xi \rangle \right) dr d\theta \\
& = \int_0^{2\pi} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \left(\left\langle \frac{D}{d\theta} \xi, J \frac{D}{dr} \xi \right\rangle - \frac{(p-2)}{p} \frac{\rho'_v(r)}{\rho_v(r)} \left\langle \frac{D}{d\theta} \xi, J \xi \right\rangle + \langle \mathcal{R}_{b_v}(u_r, u_\theta) \xi, J \xi \rangle \right) dr d\theta,
\end{aligned} \tag{5.19}$$

where u_r and u_θ denote $\frac{d}{dr} u_v$ and $\frac{d}{d\theta} u_v$, respectively, and $\mathcal{R}_{b_v}$ is the curvature tensor of the connection ∇^{b_v} . Since $\langle \frac{D}{d\theta} \xi, J \frac{D}{dr} \xi \rangle = - \langle \frac{D}{dr} \xi, J \frac{D}{d\theta} \xi \rangle$, by (5.19) and (1) of Lemma 3.5,

$$\begin{aligned}
& \left| \int_0^{2\pi} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \left\langle \frac{D}{dr} \xi, J \frac{D}{d\theta} \xi \right\rangle dr d\theta \right| \\
& \leq \frac{|p-2|}{2p} \left| \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \frac{\rho'_v(r)}{\rho_v(r)} \int_0^{2\pi} \left\langle \frac{D}{d\theta} \xi, J \xi \right\rangle d\theta dr \right| + C(b_v) \|\xi\|_{v,p}^2.
\end{aligned} \tag{5.20}$$

(2) By Poincare Lemma, see Proposition ?? in [Z1], for every circle with r fixed,

$$\begin{aligned}
\left| \int_0^{2\pi} \left\langle \frac{D}{d\theta} \xi, J \xi \right\rangle d\theta \right| & \leq \int_0^{2\pi} \left| \frac{D}{d\theta} \xi \right|^2 d\theta + C(g_{b_v}) \left\{ \left(\int_0^{2\pi} |u_\theta|^2 d\theta \right) \left(\int_0^{2\pi} |\xi|^2 d\theta \right) \right. \\
& \quad \left. + \left(\int_0^{2\pi} |u_\theta| d\theta \right) \left(\int_0^{2\pi} |\xi|^2 d\theta \right)^{\frac{1}{2}} \left(\int_0^{2\pi} \left| \frac{D}{d\theta} \xi \right|^2 d\theta \right)^{\frac{1}{2}} \right\}.
\end{aligned} \tag{5.21}$$

Since $\left| \frac{\rho'_v(r)}{\rho_v(r)} \right| \leq 2r^{-1}$ on $\tilde{A}_{v,h}$, by Holder's inequality and the first part of Lemma 3.5,

$$\begin{aligned}
& \frac{1}{2} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \left| \frac{r \rho'_v(r)}{\rho_v(r)} \right| \left(\int_0^{2\pi} r^{-1} |u_\theta| d\theta \right) \left(\int_0^{2\pi} |\xi|^2 d\theta \right)^{\frac{1}{2}} \left(\int_0^{2\pi} r^{-2} \left| \frac{D}{d\theta} \xi \right|^2 d\theta \right)^{\frac{1}{2}} r dr \\
& \leq C \|\xi\|_{v,p} \left(\int_{\tilde{A}_{v,h}} \rho_v^{-\frac{p-2}{p}} r^{-2} \left| \frac{D}{d\theta} \xi \right|^2 \right)^{\frac{1}{2}}.
\end{aligned} \tag{5.22}$$

Similarly,

$$\frac{1}{2} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \left| \frac{r \rho'_v(r)}{\rho_v(r)} \right| \left(\int_0^{2\pi} r^{-2} |u_\theta|^2 d\theta \right) \left(\int_0^{2\pi} |\xi|^2 d\theta \right) r dr d\theta \leq C(b_v) \|\xi\|_{v,p}^2. \tag{5.23}$$

Combining equations (5.21)-(5.23), we obtain

$$\begin{aligned}
& \frac{1}{2} \left| \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} \frac{\rho'_v(r)}{\rho_v(r)} \int_0^{2\pi} \left\langle \frac{D}{d\theta} \xi, J \xi \right\rangle d\theta dr \right| \\
& \leq \int_0^{2\pi} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} r^{-2} \left| \frac{D}{d\theta} \xi \right|^2 r dr d\theta + C(b_v) (\|\xi\|_{v,p}^2 + \|\xi\|_p a_h).
\end{aligned} \tag{5.24}$$

Note that

$$\begin{aligned}
& \int_0^{2\pi} \int_{\epsilon_1}^{\epsilon_2} \rho_v^{-\frac{p-2}{p}} r^{-2} \left| \frac{D}{d\theta} \xi \right|^2 r dr d\theta \\
& = \frac{1}{2} \int_{\tilde{A}_{v,h}} \rho_v^{-\frac{p-2}{p}} \left(r^{-2} \left| \frac{D}{d\theta} \xi \right|^2 + \left| \left(\frac{D}{dr} \xi + J r^{-1} \frac{D}{d\theta} \xi \right) - \frac{D}{dr} \xi \right|^2 \right) \leq \frac{1+\epsilon}{2} a_h^2 + C_\epsilon \|D_v \xi\|_{v,p}^2
\end{aligned} \tag{5.25}$$

for any $\epsilon > 0$. Combining equations (5.18), (5.20), (5.24) and (5.25), we obtain

$$a_h^2 \leq \frac{|p-2|}{p}(1+\epsilon)a_h^2 + (C(b_v) + C_\epsilon)(\|D_v \xi\|_{v,p}^2 + \|\xi\|_{v,p}^2 + \|\xi\|_{v,p} a_h).$$

Since $\frac{|p-2|}{p} < 1$, the claim follows by choosing ϵ sufficiently small.

Lemma 5.9 *For all $p \geq 1$, there exists $C_p \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R})$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$ and $h \in \hat{I}$,*

$$\xi \in \Gamma_c(\tilde{A}_{v,h}|u_v^*TV) \implies \|\nabla^{b_v} \xi\|_{g_v,p} \leq C_p(b_v) \left(\|D_v \xi\|_{v,p} + \|\xi\|_{v,p} + \left(\int_{\tilde{A}_{v,h}} \rho_v^{-\frac{p-2}{2p}} |\nabla \xi|^2 \right)^{\frac{1}{2}} \right).$$

Proof: Choose a sequence

$$\delta_0 > \dots > \delta_{N+1} > 0 \quad \text{such that} \quad \delta_0 = \epsilon_2, \quad \delta_{N+1} = \epsilon_1, \quad \frac{1}{3} \leq \frac{\delta_{l+1}}{\delta_l} \leq \frac{1}{2}.$$

For each $l = 1, \dots, N-1$, let g_l denote the metric

$$g_l = \left(\delta_l^2 + \frac{|v_h|^2}{\delta_l^2} \right)^{-1} g_v \quad \text{on} \quad \tilde{A}_l = \{(r, \theta) \in \tilde{A}_{v,h} : \delta_{l+2} \leq r \leq \delta_{l-1}\}.$$

Let $\rho_l = \delta_l^2 + |v_h|^2 \delta_l^{-2}$ and denote by A_l the annulus $\{(r, \theta) \in \tilde{A}_{v,h} : \delta_{l+1} \leq r \leq \delta_l\}$. The pullback of the metric g_l on $\tilde{A}_l$ to the annulus $(\frac{\delta_{l+2}}{\delta_l}, \frac{\delta_{l-1}}{\delta_l}) \times S^1 \subset \mathbb{R}^2$ by the map $(r, \theta) \longrightarrow (\delta_l r, \theta)$ differs from the Euclidean metric by a conformal factor bounded by $C(b_v)$, since

$$\frac{1}{100} \leq \left\{ (1 - \beta_{|v_h|}(\delta_l r)) \frac{2}{|v_h| + |v_h|^{-1} \delta_l^2 r^2} + \beta_{|v_h|}(\delta_l r) \right\} \left(\delta_l^2 + \frac{|v_h|^2}{\delta_l^2} \right)^{-\frac{1}{2}} \delta_l \leq 200,$$

whenever $r \in (\frac{1}{9}, 3)$ and $\delta_l \in (|v_h|, 1)$. Thus, by Proposition ?? in [Z1],

$$\|\nabla^{b_v} \xi\|_{g_l, L^p(A_l)} \leq C \left(\|D_v \xi\|_{g_l, L^p(\tilde{A}_l)} + \|\nabla^{b_v} \xi\|_{g_l, L^p(\tilde{A}_l)} + \|\xi du\|_{g_l, L^p(\tilde{A}_l)} \right), \quad (5.26)$$

or equivalently

$$\|\nabla^{b_v} \xi\|_{g_v, L^p(A_l)} \leq C \left(\|D_v \xi\|_{g_v, L^p(\tilde{A}_l)} + \|\rho_l^{-\frac{p-2}{2p}} \nabla^{b_v} \xi\|_{L^2(\tilde{A}_l)} + \|\xi du\|_{g_v, L^p(\tilde{A}_l)} \right). \quad (5.27)$$

Since $\frac{\rho_v(r)}{\rho_l} \in [\frac{1}{81}, 81]$ when $r \in [\delta_{l+2}, \delta_{l-1}]$, (5.27) is equivalent to

$$\left(\int_{A_l} |\nabla^{b_v} \xi|^p \right)^{\frac{1}{p}} \leq C_p(b_v) \left(\left(\int_{\tilde{A}_l} |D_v \xi|^p \right)^{\frac{1}{p}} + \left(\int_{\tilde{A}_l} \rho_v^{-\frac{p-2}{p}} |\nabla^{b_v} \xi|^2 \right)^{\frac{1}{2}} + \|\xi du_v\|_{g_v, L^p(\tilde{A}_l)} \right).$$

The claim follows by summing up the last inequality over all l and using (1) of Lemma 3.5.

Remark: The above proof does not quite apply to the two outermost annuli A_1 and A_N . However, since ξ is compactly supported in $\tilde{A}_{v,h}$, the proof of Proposition ?? in [Z1] can be applied to A_1 with $A_1 \cup A_2$ replacing $\tilde{A}_1$ to (5.26), and similarly for A_N . Alternatively, for the purposes of proving Proposition 5.11 below, it is sufficient to prove Lemma 5.9 and Corollary 5.10 for ξ that vanish on A_1 and A_N .

Corollary 5.10 For all $p > 1$, there exists $C_p \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R})$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$ and $h \in \hat{I}$

$$\xi \in \Gamma_c(\tilde{A}_{v,h}; u^*TV) \implies \|\xi\|_{v,p,1} \leq C_p(b_v)(\|D_v\xi\|_{v,p} + \|\xi\|_{v,p}).$$

Proof: This corollary follows immediately from Lemmas 5.10 and 5.10.

Proposition 5.11 For all $p > 1$, there exists $C_p \in C^\infty(\mathcal{M}_T^{(0)}; \mathbb{R})$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$,

$$\|\xi\|_{v,p,1} \leq C_p(b_v)(\|D_v\xi\|_{v,p} + \|\xi\|_{v,p}) \quad \text{for all } \xi \in \Gamma(u_v).$$

Proof: This proposition follows from Corollary 5.10 and Proposition ?? in [Z1] by taking a partition of unity as in the proof of Proposition 5.7.

5.4 Fiber-Uniform Inverse for the Operator D_v

Lemma 5.12 Let $\{v_k\}$ be a sequence in $F^{(0)}\mathcal{T}_\delta$ that converges to $b^* \in \mathcal{M}_T^{(0)}$. Suppose $\xi_k \in \Gamma(u_{v_k})$ is such that $\|\xi_k\|_{v_k,p,1} \leq 1$ for all k , while $\|D_{v_k}\xi_k\|_{v_k,p} \longrightarrow 0$ as $k \longrightarrow \infty$ for some $p > 2$. Then a subsequence of $\{\xi_k\}$ C^0 -converges $\xi^* \in \Gamma_-(b^*)$. Furthermore, $\|\xi_k\|_{v_k,C^0}$ converges to $\|\xi^*\|_{b^*,C^0}$.

Proof: (1) Write $b^* = (S, M, I; x^*, (j, y^*), u_I)$ and

$$v_k = (b_k, v_k) = ((S, M, I; x_k, (j, y_k), u_k), (v_k)_{\hat{I}}).$$

For each $i \in I$ and $\delta > 0$, put

$$S_{i,\delta}^* = \{z \in \Sigma_{b,i} : r_{b^*,h}(z) \geq \delta \ \forall h \in \hat{I}; |q_S^{-1}(z)| \geq \delta \text{ if } i > 0\}.$$

For $i \in I$ and all k sufficiently large (depending on δ), define $\zeta_{k,i}, \xi'_{k,i} \in \Gamma(u_i^*|S_{i,\delta}^*)$ by

$$\exp_{b^*, u_i^*(z)} \zeta_{k,i}(z) = u_{v_k}(q_{v_k}^{-1}(z)), \quad \|\zeta_{k,i}\|_{b^*,C^0} < \text{inj } g_{V,b^*}; \quad \Pi_{b^*, \zeta_{k,i}(z)} \xi'_{k,i}(z) = \xi_k(q_{v_k}^{-1}(z)).$$

Since $\|\nabla^{b^*} \zeta_{k,i}\|_{b^*,C^0} \leq C$ for k sufficiently large, (1) of Lemma 3.5 and by Corollary ?? in [Z1],

$$\begin{aligned} \|\xi'_{k,i}\|_{b^*,p,1} &\leq (1 + \epsilon_k)\|\xi_k\|_{v_k,p,1} + \epsilon_k\|\xi_k\|_{v_k,C^0}, \\ \|D_{b,u_i}\xi'_{k,i}\|_{b^*,p} &\leq (1 + \epsilon_k)\|D_{v_k}\xi_k\|_{v_k,p} + \epsilon_k\|\xi_k\|_{v_k,C^0}, \end{aligned} \tag{5.28}$$

where $\epsilon_k \longrightarrow 0$ as $k \longrightarrow \infty$. Since $\|\xi_k\|_{v_k,p,1} \leq 1$, (2) of Lemma 3.5 applied to (5.28),

$$\|\xi'_{k,i}\|_{b^*,p,1} \leq (1 + \tilde{\epsilon}_k)\|\xi_k\|_{v_k,p,1} + \tilde{\epsilon}_k, \quad \|D_{b,u_i}\xi'_{k,i}\|_{b^*,p} \leq (1 + \tilde{\epsilon}_k)\|D_{v_k}\xi_k\|_{v_k,p} + \tilde{\epsilon}_k, \tag{5.29}$$

where $\tilde{\epsilon}_k \longrightarrow 0$ as $k \longrightarrow \infty$. Sobolev's embedding theorem then implies that $\xi'_{k,i}$ converges to a vector field $\xi_i^* \in \Gamma(u_i|\Sigma_{b^*,i}^*)$ in the C^0 -norm on the compact subsets of $\Sigma_{b^*,i}^*$. Furthermore, $\|\xi_i^*\|_{b^*,C^0} < \infty$, since

$$\|\xi'_{k,i}\|_{b^*,C^0} \leq (1 + \epsilon_k)\|\xi_k\|_{v_k,C^0} \leq C.$$

(2) We will now show that $D_{b^*, u_i^*}\xi_i^* = 0$ weakly, i.e. $\langle \xi_i^*, D_{b^*, u_i^*}\eta \rangle_{b^*} = 0$ for any $\eta \in \Gamma^{0,1}(u_i^*)$. We have

$$\langle \xi_i^*, D_{b^*, u_i^*}\eta \rangle_{b^*} = \lim_{\delta \longrightarrow 0} \int_{S_{i,\delta}^*} \langle \xi_i^*, D_{b^*, u_i^*}\eta \rangle_{b^*} = \lim_{\delta \longrightarrow 0} \lim_{k \longrightarrow \infty} \int_{S_{i,\delta}^*} \langle \xi'_{k,i}, D_{b^*, u_i^*}\eta \rangle_{b^*}, \tag{5.30}$$

since $\xi'_{k,i} \rightarrow \xi_i^*$ in the C^0 -norm on $S_{i,\delta}$. By integration by parts,

$$\left| \int_{S_{i,\delta}^*} \langle \xi'_{k,i}, D_{b^*,u_i^*}^* \eta \rangle_{b^*} - \int_{S_{i,\delta}^*} \langle D_{b^*,u_i^*}^* \xi'_{k,i}, \eta \rangle_{b^*} \right| \leq C \int_{\partial S_{i,\delta}^*} |\xi'_{k,i}| |\eta| \leq C' \|\xi'_{k,i}\|_{b^*,C^0} \|\eta\|_{b^*,C^0}. \quad (5.31)$$

Since $\|D_{b^*,u_i^*}^* \xi'_{k,i}\|_{b^*,p} \rightarrow 0$ as $k \rightarrow \infty$ on $S_{i,\delta}^*$ and $\|\xi'_{k,i}\|_{b^*,C^0} \leq C$, by (5.30) and (5.31),

$$\langle \langle \xi_i^*, D_{b^*,u_i^*}^* \eta \rangle \rangle_{b^*} = 0 \quad \forall \eta \in \Gamma^{0,1}(u_i^*).$$

(3) Since $D_{b^*,u_i^*}^* \xi_i^* = 0$ weakly on S_i and $D_{b^*,u_i^*}^*$ is an elliptic operator, it follows that ξ_i^* is smooth and $D_{b^*,u_i^*}^* \xi_i^* = 0$. It will now be shown that $\xi_{\iota_h}^*(x_h^*) = \xi_h^*(\infty)$ for all $h \in \hat{I}$, i.e. $\xi^* \equiv \xi_I^* \in \Gamma(b^*)$. For each $h \in \hat{I}$, let $A_{h,\delta,k} \subset S$ denote the small cylinder connecting $q_{v_k}^{-1}(S_{h,\delta}^*)$ and $q_{v_k}^{-1}(S_{\iota_h,\delta}^*)$. Let $\epsilon > 0$ be any small number. Choose small $\delta > 0$ such that $u_h(B_{b^*,h}(\infty, 2\delta))$ and $u_{\iota_h}^*(B_{b^*,\iota_h}^*(x_h^*, 2\delta))$ lie in $B_{b^*}(u_h^*(\infty), \epsilon)$. Then we can write

$$u_{b^*}^*(z) = \exp_{b^*,u_{b^*}^*(x_h^*)} \bar{u}_{b^*}(z), \quad |\bar{u}_{b^*}(z)| < \text{inj } g_{V,b^*}; \quad \bar{\xi}'_k(z) \equiv \Pi_{b^*,\bar{u}_{b^*}(z)}^{-1} \xi'_k(z)$$

for $z \in B_{b^*,h}(\infty, \delta) \cup B_{b^*,\iota_h}^*(x_h^*, \delta)$. Similarly, put

$$\bar{\xi}_h^*(z) = \Pi_{b^*,\bar{u}_{b^*}(z)}^{-1} \xi_h^*(z) \quad \text{and} \quad \bar{\xi}_{\iota_h}^*(z) = \Pi_{b^*,\bar{u}_{b^*}(z)}^{-1} \xi_{\iota_h}^*(z)$$

for z in $B_{b^*,h}(\infty, \delta)$ and in $B_{b^*,\iota_h}^*(x_h^*, \delta)$, respectively. We can also assume that δ is so small that $|\bar{\xi}_h^* - \xi_h^*(\infty)|_{b^*}$ and $|\bar{\xi}_{\iota_h}^* - \xi_{\iota_h}^*(x_h^*)|_{b^*}$ do not exceed ϵ on $B_{b^*,h}(\infty, \delta)$ and on $B_{b^*,\iota_h}^*(x_h^*, \delta)$, respectively. Choose large k^* such that all $k > k^*$

$$\|\xi^* - \xi'_k\|_{C^0(S_{h,\delta}^* \cup S_{\iota_h,\delta}^*)} \leq \epsilon.$$

It can be assumed that $u_k(A_{h,2\delta,k})$ lies in $B_{b^*}(u^*(x_h^*); 2\epsilon)$ for $k > k^*$. Thus, we can write

$$u_k(z) = \exp_{b^*,u(x_h^*)} \bar{u}_k(z), \quad |\bar{u}_k(z)|_{b^*} < \text{inj } g_{V,b^*}; \quad \bar{\xi}_k(z) \equiv \Pi_{b^*,\bar{u}_k(z)}^{-1} \xi_k(z)$$

if $z \in A_{h,\delta,k}$. Pick points z_1 and z_2 , one on each component of the boundary of $A_{h,\delta,k}$. Then

$$\begin{aligned} |\xi_h^*(\infty) - \xi_{\iota_h}^*(x_h^*)|_{b^*} &\leq 2(\epsilon + |\bar{\xi}_h^*(q_{v_k}(z_1)) - \bar{\xi}_{\iota_h}^*(q_{v_k}(z_2))|_{b^*}) \\ &\leq 4(\epsilon + |\bar{\xi}'_{k,h}(q_{v_k}(z_1)) - \bar{\xi}'_{k,\iota_h}(q_{v_k}(z_2))|_{b^*}) \\ &\leq C(\epsilon + |\bar{\xi}_k(z_1) - \bar{\xi}_k(z_2)|_{b^*} + \|\zeta_k\|_{b^*,C^0(S_{h,\delta}^* \cup S_{\iota_h,\delta}^*)} \|\bar{\xi}_k\|_{b^*,C^0(A_{h,\delta,k})}). \end{aligned} \quad (5.32)$$

Since $A_{h,\delta,k}$ is uniformly equivalent to the union of two annuli with the larger radius bounded above by δ and the smaller radius less than half of the larger, by Lemma ?? in [Z1] and Holder's inequality,

$$|\bar{\xi}_k(z_1) - \bar{\xi}_k(z_2)|_{b^*} \leq C |\bar{\xi}_k(z_1) - \bar{\xi}_k(z_2)|_{b_k} \leq C' \delta^{\frac{2(p-2)}{p}} \|d\bar{\xi}_k\|_{v_k, L^p(A_{h,\delta,k})}. \quad (5.33)$$

By Corollary ?? in [Z1] and Proposition 5.11,

$$\|d\bar{\xi}_k\|_{v_k, L^p(A_{h,\delta,k})} \leq \|\xi_k\|_{v_k, p, 1} + \|du_{v_k}\|_{v_k, p} \|\xi_k\|_{v_k, C^0} \leq C. \quad (5.34)$$

Combining equations (5.32)-(5.34), we obtain

$$|\xi_h^*(\infty) - \xi_{\iota_h}^*(x_h^*)|_{b^*} \leq C(\epsilon + \delta^{\frac{2(p-2)}{p}} + \|\zeta_k\|_{b^*,C^0(S_{h,\delta}^* \cup S_{\iota_h,\delta}^*)}). \quad (5.35)$$

Since the last term in (5.35) tends to zero as $k \rightarrow \infty$ and ϵ and δ can be chosen to be arbitrarily small, it follows $\xi_h^*(\infty) = \xi_{\iota_h}^*(x_h^*)$.

Proposition 5.13 *For any simple bubble type $\mathcal{T}$, there exist $C, \delta \in C^\infty(\mathcal{M}_{\mathcal{T}}^{(0)}; \mathbb{R})$ such that for all $v \in F^{(0)}\mathcal{T}_\delta$ if $\mathcal{T}$ is regular and $v \in F^{(0)}\mathcal{T}_\delta$ if $\mathcal{T}$ is semiregular,*

$$\|\xi\|_{v,p,1} \leq C_p(b_v)\|D_v\xi\|_{v,p} \quad \forall \xi \in \Gamma_+(v) \quad \text{and} \quad \forall \xi \in \tilde{\Gamma}_+(v).$$

Proof: If not, we can choose a sequence $v_k \in F^{(0)}\mathcal{T}_\delta$, converging to some $b \in \mathcal{M}_{\mathcal{T}}^{(0)}$ and vector fields $\xi_k \in \Gamma_+(v_k)$ (or $\xi_k \in \tilde{\Gamma}_+(v_k)$) such that $\|\xi_k\|_{v_k,p,1} = 1$, while $\|D_{v_k}\xi\|_{v_k,p} \rightarrow 0$. If $\xi_k \in \Gamma_+(v_k)$, note that $\{\Gamma_-(v_k)\}$ C^0 -converges to $V \equiv \Gamma_-(b)$. If $\xi_k \in \tilde{\Gamma}_+(v_k)$, by Definition 3.11, a subsequence of $\{\tilde{\Gamma}_-(v_k)\}$ C^0 -converges to a subspace $V \subset L_1^p(b)$ such that $\pi_{b,-} : V \rightarrow \Gamma_-(b)$ is an isomorphism. In either case, by the first statement of Lemma 5.12, a subsequence of $\{\xi_k\}$ C^0 -converges to a vector field $\xi^* \in \Gamma_-(b)$. By the second statement of Lemma 5.12, ξ^* must be orthogonal V , since $\xi_k \in \Gamma_+(v_k)$ (or $\xi_k \in \tilde{\Gamma}_+(v_k)$). Thus, $\xi^* = 0$. On the other hand, by Proposition 5.11, there exists $\epsilon > 0$ such that $\|\xi_k\|_{v_k,p} \geq \epsilon$ for all k sufficiently large. However, by Lemma 5.12, $\|\xi_k\|_{v_k,C^0} \rightarrow 0$, which is a contradiction.

References

- [FO] K. Fukaya and K. Ono, *Arnold Conjecture and Gromov-Witten Invariants*, Topology 38 (1999), no. 5, 933–1048.
- [GH] P. Griffiths and J. Harris, *Principles of Algebraic Geometry*, 1994.
- [I] E. Ionel, *Genus-One Enumerative Invariants in $\mathbb{P}^n$ with Fixed j -Invariant*, Duke Math. J. 94 (1998), no. 2, 279–324.
- [LM] H. Lawson and M. Michelsohn, *Spin Geometry*, Princeton University Press, 1989.
- [LT] J. Li and G. Tian, *Virtual moduli cycles and Gromov-Witten invariants of general symplectic manifolds*, Topics in symplectic 4-manifolds, 47–83, First Int. Press Lect. Ser., I, Internat. Press, 1998.
- [MS] D. McDuff and D. Salamon, *J-holomorphic Curves and Quantum Cohomology*, American Mathematical Society, 1994.
- [RT] Y. Ruan and G. Tian, *A Mathematical Theory of Quantum Cohomology*, J. Diff. Geom. 42 (1995), no. 2, 259–367.
- [T] C. Taubes, *Self-dual Connections on 4-Manifolds with Indefinite Intersection Matrix*, J. Diff. Geom. 19 (1984), 517–560.
- [Z1] A. Zinger, *Basic Estimates of Riemannian Geometry Used in Gluing Pseudoholomorphic Maps*, notes.
- [Z2] A. Zinger, *Enumeration of Genus-Two Curves with a Fixed Complex Structure in $\mathbb{P}^2$ and $\mathbb{P}^3$* , preprint.